


Intramolecular Hydroamination of Alkynes Catalyzed by Pd(PPh₃)₄/Triphenylphosphine under Neutral Conditions

Gan B. Bajracharya, Zhibao Huo, and Yoshinori Yamamoto*

Department of Chemistry, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan

yoshi@yamamoto1.chem.tohoku.ac.jp


Received March 3, 2005



The intramolecular hydroamination of alkynes tethered with amino group **1** in the presence of catalytic amounts of Pd(PPh₃)₄ and PPh₃ in benzene at 100 °C proceeded smoothly without the use of any additional acid source to afford five- and six-membered nitrogen heterocycles **2** in good to excellent yields. A compulsory addition of carboxylic acid as a cocatalyst was not needed, and the reaction could be carried out under essentially neutral conditions.

Nonfunctionalized carbon–carbon multiple bonds are generally unreactive toward nucleophiles because of their electron-rich π -orbitals. However, the use of transition metal catalysts enables the addition of nucleophiles to such unactivated C–C multiple bonds.¹ Nitrogen-containing heterocycles are main building blocks for many biologically important compounds. The hydroamination reaction of alkynes using transition metal catalysts is an attractive route to reach such compounds,² since the reaction proceeds in an atom-economical manner. Several catalytic systems have been explored to promote the hydroamination of carbon–carbon multiple bonds.³ Recently, we reported that the palladium-mediated hy-

droamination of allenes,⁴ enynes,⁵ methylenecyclopropanes,⁶ and cyclopropene⁷ proceeded smoothly, giving the corresponding allylic amines in good to high yields. The presence of a carboxylic acid was needed as an unavoidable additive for the Pd-mediated inter- and intramolecular hydroamination of internal alkynes.⁸ In these reactions, the presence of carboxylic acid as a cocatalyst modifies the palladium catalyst to a more active hydridopalladium species via in situ oxidative addition of Pd⁰ into H-A. We also reported that an asymmetric version of the intramolecular hydroamination of alkynes proceeded smoothly by using a Pd₂(dba)₃·CHCl₃–PhCO₂H catalytic system in the presence of (*R,R*)-RENORPHOS as a chiral ligand to give various kinds of optically active five- and six-membered nitrogen heterocycles (eq 1).^{8c}



The use of an stoichiometric amount of an expensive chiral ligand [(*R,R*)-RENORPHOS] for obtaining a better yield and high enantioselectivity was a drawback for this reaction, and thus we further struggled in this area to find much milder and more efficient reaction conditions. While we were searching for an alternative catalytic system, we found that the intramolecular hydroamination of alkynes proceeded without using an acid source. We report that in the presence of 15 mol % Pd(PPh₃)₄ and 10 mol % PPh₃ in benzene at 100 °C the intramolecular hydroamination reaction of alkynes tethered with amino group **1** proceeded well to give the corresponding nitrogen heterocycles **2** in good to excellent yields under essentially neutral conditions.

After several attempts, we found that the reaction of **1a** in the presence of 15 mol % Pd(PPh₃)₄ together with 10 mol % PPh₃ in benzene at 100 °C for 66 h produced the corresponding hydroamination product **2a** in 86%

(1) Yamamoto, Y.; Radhakrishnan, U. *Chem. Soc. Rev.* **1999**, *28*, 199–207.

(2) Nakamura, I.; Yamamoto, Y. *Chem. Rev.* **2004**, *104*, 2127–2198.

(3) (a) Savoia, D. *Houben-Weyl*; Helmchen, G., Hoffmann, R. W., Mulzer, J., Schaumann E., Eds.; Georg Thieme Verlag: Stuttgart, 1995; Vol. E21e, pp 5356–5394. (b) Müller, T. E.; Beller, M. *Chem. Rev.* **1998**, *98*, 675–703 and references therein. (c) Taube, R. *Applied Homogeneous Catalysis with Organometallic Compounds*; Cornils, B., Herrmann, W. A., Eds.; Wiley-VCH: Weinheim, 2002; Vol. 1, Applications; pp 513–524. (d) Schaffrath, H.; Keim, W. *J. Mol. Catal. A: Chem.* **2001**, *168*, 9–14. (e) Nobis, M.; Driessen-Hölscher, B. *Angew. Chem., Int. Ed.* **2001**, *40*, 3983–3985. (f) Minami, T.; Okamoto, H.; Ikeda, S.; Tanaka, R.; Ozawa, F.; Yoshifuji, M. *Angew. Chem., Int. Ed.* **2001**, *40*, 4501–4503. (g) Straub, B. F.; Bergman, R. G. *Angew. Chem., Int. Ed.* **2001**, *40*, 4632–4635. (h) Utsunomiya, M.; Kuwano, R.; Kawatsura, M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2003**, *125*, 5608–5609. (i) Alonso, F.; Beletskaya, I. P.; Yus, M. *Chem. Rev.* **2004**, *104*, 3079–3160. (j) Shimada, T.; Bajracharya, G. B.; Yamamoto, Y. *Eur. J. Org. Chem.* **2005**, 59–62 and references therein.


(4) (a) Al-Masum, M.; Meguro, M.; Yamamoto, Y. *Tetrahedron Lett.* **1997**, *38*, 6071–6074. (b) Meguro, M.; Yamamoto, Y. *Tetrahedron Lett.* **1998**, *39*, 5421–5424.

(5) Radhakrishnan, U.; Al-Masum, M.; Yamamoto, Y. *Tetrahedron Lett.* **1998**, *39*, 1037–1040.

(6) Nakamura, I.; Itagaki, H.; Yamamoto, Y. *J. Org. Chem.* **1998**, *63*, 6458–6459.

(7) Nakamura, I.; Bajracharya, G. B.; Yamamoto, Y. *J. Org. Chem.* **2003**, *68*, 2297–2299.

(8) (a) Kadota, I.; Shibuya, A.; Lutete, L. M.; Yamamoto, Y. *J. Org. Chem.* **1999**, *64*, 4570–4571. (b) Lutete, L. M.; Kadota, I.; Shibuya, A.; Yamamoto, Y. *Heterocycles* **2002**, *58*, 347–357. (c) Lutete, L. M.; Kadota, I.; Yamamoto, Y. *J. Am. Chem. Soc.* **2004**, *126*, 1622–1623. (d) Patil, N. T.; Khan, N.; Yamamoto, Y. *Tetrahedron Lett.* **2004**, *45*, 8497–8499. (e) Patil, N. T.; Wu, H.; Kadota, I.; Yamamoto, Y. *J. Org. Chem.* **2004**, *69*, 8745–8750. For the other type of reactions using H-A-Pd, see: (f) Trost, B. M.; Schmidt, T. *J. Am. Chem. Soc.* **1988**, *110*, 2301–2303. (g) Trost, B. M.; Brieden, W.; Baringhaus, K. H. *Angew. Chem., Int. Ed. Engl.* **1992**, *31*, 1335–1336. (h) Trost, B. M.; Michellys, P. Y.; Gerusz, V. *J. Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 1750–1753. Review: (i) Trost, B. M. *Chem. Eur. J.* **1998**, *4*, 2405–2412.

TABLE 1. Optimization of Reaction Conditions for the Pd/Phosphine Additive Catalyzed Intramolecular Hydroamination of Alkynes^a

entry	catalyst (15 mol %)	phosphine additive (10 mol %)	additional additive (10 mol %)	time (h)	1b yield (%) ^b	2b yield (%) ^b
1	Pd(PPh ₃) ₄	PPh ₃		37	0	98 (87)
2		PPh ₃		37	100	0
3	Pd(PPh ₃) ₄			37	84	trace
4	Pd(PPh ₃) ₄	P(O)Ph ₃		41	18	41
5	Pd(PPh ₃) ₄	P(<i>o</i> -tolyl) ₃		41	5	49
6	Pd(PPh ₃) ₄	dppp		41	13	63
7	Pd(PPh ₃) ₄	dppf		41	22	55
8	Pd(PPh ₃) ₄	PPh ₃	CH ₃ CO ₂ H	16	trace	55
9	Pd(PPh ₃) ₄	PPh ₃	PhCO ₂ H	9	0	28
10 ^c	Pd(PPh ₃) ₄		PhCO ₂ H	overnight	0	94
11	Pd ₂ dba ₃ ·CHCl ₃	PPh ₃		37	37	trace
12	PdCl ₂ (PPh ₃) ₂	PPh ₃		37	88	0
13	Pd(OAc) ₂	PPh ₃		37	88	0
14	[(η^3 -C ₃ H ₅)PdCl] ₂	PPh ₃		37	80	0
15 ^d	Pd(PPh ₃) ₄	PPh ₃		37	0	77

^a The reaction of **1b** (0.1 mmol) was carried out in the presence of 15 mol % palladium catalyst and 10 mol % phosphine additive in benzene (0.1 M) at 100 °C under argon unless otherwise noted. ^b Yield was calculated from ¹H NMR integration using CH₂Br₂ as an internal standard; isolated yield is in parentheses. ^c 15 mol % PhCO₂H was used. ^d The reaction temperature was 120 °C.





FIGURE 1. Time profile of the cyclization of **1a** (0.1 mmol) in the presence of 15 mol % Pd(PPh₃)₄ and 10 mol % PPh₃ in benzene (0.1 M) at 100 °C.

yield (99% NMR yield) (eq 2). As shown in Figure 1, the



time profile of the reaction of **1a** monitoring by NMR indicated that the complete conversion of the substrate occurred within 66 h. In contrast, the reaction of **1a** in the presence of 15 mol % Pd(PPh₃)₄ only *without the presence of additional PPh₃* afforded only trace amounts of the product **2a** with the recovery of significant amounts of the starting material even after prolonged heating. The reaction of **1a** in the presence of 10 mol % PPh₃ only did not proceed at all, and the substrate was recovered. These results clearly indicate that combined use of Pd(PPh₃)₄ and PPh₃ is the key for cyclization of **1a**. Hartwig et al. apparently reported two examples of the Pd-catalyzed intermolecular hydroamination of cyclohexadiene without











use of an acid additive.⁹ In their report, Pd(PPh₃)₄ was generated in situ from [(η^3 -C₃H₅)PdCl]₂ and PPh₃, and in general the use of a catalytic amount of trifluoroacetic acid was needed as a cocatalyst for the other examples. To the best of our knowledge, the Pd(PPh₃)₄/PPh₃ catalytic system for the catalytic hydroamination of alkynes has not been reported in the past.

We observed that the reaction of trifluoromethanesulfonyl (Tf)-substituted substrate **1b** was faster than that of **1a** having the nonafluorobutanesulfonyl (Nf) group. Hence, for further optimization of the reaction conditions, we chose **1b** as a model substrate (eq 3), and the results are summarized in Table 1. The substrate **1b** was cyclized in the presence of both 15 mol % Pd(PPh₃)₄ and 10 mol % PPh₃ to afford the pyrrolidine **2b** within 37 h with excellent yield (87%) (entry 1). In the absence of either Pd(PPh₃)₄ or PPh₃, essentially no reaction was observed, indicating the importance of the combination of both Pd(PPh₃)₄ and PPh₃ (entries 2 and 3). The use of other monodentate [P(O)Ph₃, P(*o*-tolyl)₃] and bidentate ligands [dppp = 1,3-bis(diphenylphosphino)-propane, dppf = 1,1'-bis(diphenylphosphino)-ferrocene] afforded the product in moderate yields (entries 4–7). The use of additional carboxylic acids was not essential for this transformation, resulting in poor chemical yields (entries 8 and 9). Although the intramolecular hydroamination using the Pd(PPh₃)₄–PhCO₂H catalytic system without additional PPh₃ gave a high chemical yield in rather short reaction time (entry 10), we were motivated to carry out the present reaction under essentially neutral conditions. Other palladium catalysts such as Pd₂dba₃·CHCl₃, PdCl₂-(PPh₃)₂, Pd(OAc)₂, [(η^3 -C₃H₅)PdCl]₂, etc. were completely not effective (entries 11–14). Increasing reaction temperature up to 120 °C led to a moderate yield (entry 15).

The scope of the intramolecular hydroamination of alkynes is summarized in Table 2. Substrates **1c** and **1d**

(9) Löber, O.; Kawatsura, M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2001**, *123*, 4366–4367.

TABLE 2. Pd(PPh₃)₄/PPh₃-Catalyzed Intramolecular Hydroamination of Alkynes^a


Entry	Substrate	Time/h	Product	Yield/% ^b
1 ^c		38		75
2 ^c		100		70
3		71		80 ^d
4 ^c		72		61
5		96		80

^a The reaction of **1** (0.3 mmol) was carried out in the presence of 15 mol % Pd(PPh₃)₄ and 10 mol % PPh₃ in benzene (0.1 M) at 100 °C for the indicated reaction time under argon unless otherwise noted. ^b Isolated yield. ^c 30 mol % of Pd(PPh₃)₄ and 20 mol % of PPh₃ were used. ^d A mixture of *trans* and *cis* isomers (1.7:1) was obtained. The ratio was determined by ¹H NMR spectral analysis.

having methoxy and trifluoromethyl groups at the *para*-position of aromatic ring afforded the corresponding cyclized products **2c** and **2d**, respectively, in good yields, indicating that the substituents of aromatic ring did not exert a significant influence upon the cyclization (entries 1 and 2). In both cases, increased amounts of the catalyst were needed to obtain good chemical yields. The reaction of **1e**, having a pentyl group at the alkyne terminus, under the standard conditions proceeded smoothly to give a 1.7:1 mixture of *trans* and *cis* isomers of pyrrolidine **2e** in a good yield (entry 3). Not only Nf and Tf but also Bn could be used as a protecting group for the amine; **2f** was obtained from **1f** in 61% yield (entry 4). The cyclization of **1g** afforded the six-membered piperidine **2g** in a good yield (entry 5).

A proposed mechanism, which explains the role of triphenylphosphine as a cocatalyst in the present hydroamination reaction, is shown in Scheme 1, although it is highly speculative. Triphenylphosphine, acting as a Brønsted base, abstracts a proton from the nitrogen functionality at the early stage, producing **3** and a phosphonium ion [HPPH₃]⁺. Hydropalladation of alkyne with the cationic palladium hydride species generated from Pd⁰ and the phosphonium ion produces the vinylpalladium intermediate **4**.¹⁰ The vinylpalladium in-

SCHEME 1. A Plausible Mechanism



intermediate **4** then undergoes β -elimination to give the allene **5** coordinated with the cationic palladium hydride. A subsequent hydropalladation of **5** furnishes the π -allylpalladium species **6**. Nucleophilic attack of the amine on the resulting π -allyl moiety gives the final product **2** along with regeneration of Pd⁰. In our previous reports on the Pd/carboxylic acid catalyzed analogous reaction, we proposed the oxidative addition of Pd⁰ to RCOO–H bond of acid as the very first step to generate catalytically active hydridopalladium complex.^{8a–e} In contrast, formation of a phosphonium ion is the beginning step in the present transformation.

In summary we have developed a simple and acid free hydroamination protocol for the synthesis of various nitrogen heterocycles. This reaction provides a very

(10) In the palladium-catalyzed intramolecular hydrocarboxylation of ϵ -alkynyl malononitriles, we proposed a rapid intramolecular cyclization of carbanion to the triple bond of alkyne coordinated with a cationic palladium hydride generated from Pd(OAc)₂ and a carbon pronucleophile; see: Tsukada, N.; Yamamoto, Y. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2477–2480.

useful method for a catalytic synthesis of five- and six-membered nitrogen heterocycles. Now, it is clear that the catalytic intramolecular hydroamination of alkynes can be accomplished under essentially neutral conditions without using any acid source. We are further continuing our effort to develop a mild and an efficient asymmetric version of this hydroamination reaction and synthesizing various natural compounds by utilizing our methodology as the key step.¹¹

Experimental Section

Representative Procedure for Intramolecular Hydroamination. To a mixture of Pd(PPh₃)₄ (17.3 mg, 0.015 mmol, 15

(11) Recently we reported the synthesis of indolizidine (–)-209D based on an intramolecular hydroamination strategy; see: Patil, N. T.; Pahadi, N. K.; Yamamoto, Y. *Tetrahedron Lett.* **2005**, *46*, 2101–2103.

mol %), PPh₃ (2.6 mg, 0.01 mmol, 10 mol %), and *N*-triflyl-(6-phenyl-hex-5-ynyl)-amine **1b** (30.5 mg, 0.1 mmol) was added benzene (1.0 mL, 0.1 M) under argon atmosphere in a Wheaton microreactor. The mixture was stirred for 10 min at room temperature then heated at 100 °C. Progress of the reaction was monitored by TLC (hexane/ethyl acetate; 4/1). After complete consumption of the starting material, the reaction mixture was filtered through a short column of SiO₂ using diethyl ether as an eluent, and the resulting filtrate was concentrated. The residue was purified by column chromatography (silica gel, hexane/ethyl acetate; 20/1–5/1) to afford *N*-triflyl-2-[(*E*)-2-phenylethenyl]pyrrolidine **2b** in 87% yield (26.5 mg).

Supporting Information Available: Starting material preparation; characterization data for all new compounds; and ¹H and ¹³C NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO050412W