

Cluster Compounds

DOI: 10.1002/anie.200602203

Synthesis of Tellurido-Bridged IrPt₂, IrPd₂, and IrPtPd Clusters by Inserting Zero-Valent Pt and Pd Centers into Te–C Bonds**

*Takafumi Nakagawa, Hidetake Seino, Shoken Nagao, and Yasushi Mizobe**

Soluble transition-metal sulfido clusters have been investigated extensively because of their relevance to biological and industrial catalysts. In contrast, the chemistry of the analogues of tellurium, the heavier congener of sulfur, has been less-well developed.^[1] This difference is presumably because, at least in

[*] T. Nakagawa, Dr. H. Seino, Dr. S. Nagao, Prof. Dr. Y. Mizobe
Institute of Industrial Science
The University of Tokyo
Komaba, Meguro-ku, Tokyo, 153-8505 (Japan)
Fax: (+81) 3-5452-6361
E-mail: ymizobe@iis.u-tokyo.ac.jp


[**] This work was supported by Grant-in-Aid for Scientific Research on Priority Areas (No. 14078206, "Reaction Control of Dynamic Complexes") from the Ministry of Education, Culture, Sports, Science and Technology, Japan, and by CREST of JST.

part, good tellurido sources that can be used in organic media are scarce.^[2]

In the last decade, studies in our group have focused on the pursuit of rational methods to construct metal sulfido and selenido cluster cores with desired structures and atom compositions.^[3] In the course of these studies, we have demonstrated that newly prepared dinuclear μ -hydrosulfido and μ -hydroselenido complexes of the type $[\text{Cp}^*\text{MCl}(\mu\text{-EH})_2\text{MCp}^*\text{Cl}]$ ($\text{M} = \text{Ru}, \text{Ir}, \text{Rh}$; $\text{E} = \text{S}, \text{Se}$; $\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$) can serve as versatile building blocks to synthesize a variety of homo- and heterometallic sulfido, selenido, and sulfido-selenido clusters.^[3,4] However, preparation of the corresponding tellurido clusters is not yet feasible, as the hydrotellurido analogues of these μ -EH complexes are not accessible.

Formations of tellurido ligands from coordinated telluroates have been observed for isolated or in situ-generated complexes of Cr ,^[5] Mo ,^[6] Re ,^[7] Pd ,^[8] Ag ,^[9] and Cu .^[10] Furthermore, cleavage of the $\text{E}-\text{C}$ bonds of organochalcogen compounds has also been demonstrated in a well-defined manner for the reactions with low-valent late transition metals,^[11] which include the oxidative addition reactions of Ph_2Te with zero-valent Group 10 metal complexes.^[12] On the basis of these findings, we have attempted the synthesis of mixed-metal tellurido clusters by a new route, which involves the initial preparation of an Ir telluroato complex by using readily available RTeTeR compounds (in which $\text{R} = \text{aryl}$) as the Te source and subsequent transformation of this complex into the tellurido clusters through $\text{Te}-\text{C}$ bond cleavage of the telluroato ligands by using Group 10 noble metals.

Treatment of $[\text{Cp}^*\text{Ir}(\text{CO})_2]$ with one equivalent of TolTeTeTol ($\text{Tol} = p\text{-CH}_3\text{C}_6\text{H}_4$) in benzene at room temperature afforded a bis(telluroato) complex $[\text{Cp}^*\text{Ir}(\text{CO})(\text{TeTol})_2]$ (**1**) in high yield [Eq. (1)], which has been characterized by



single-crystal X-ray analysis (Figure 1).^[13] Complex **1** has an expected three-legged structure, for which the interatomic distances and angles are not unusual. Related thiolato and selenolato complexes $[\text{Cp}^*\text{Ir}(\text{CO})(\text{EPh})_2]$ ($\text{E} = \text{S}, \text{Se}$) are known, which were prepared from $[\text{Cp}^*\text{Ir}(\text{CO})_2]$ and





Figure 1. An ORTEP drawing of **1**. Hydrogen atoms are omitted for clarity (thermal ellipsoids shown at 30% probability). Selected bond lengths [Å] and angles [°]: Ir–Te(1) 2.6654(6), Ir–Te(2) 2.6579(8); Te(1)–Ir–Te(2) 79.46(2).

PhEEPh in THF by photo-irradiation. However, X-ray structures were not determined.^[14]

When **1** was treated with two equivalents of $[\text{Pt}(\text{PPh}_3)_3]$ in THF at room temperature, the mixed-metal tellurido cluster $[\{\text{Cp}^*\text{Ir}(\text{CO})\}(\mu_3\text{-Te})_2\{\text{PtTol}(\text{PPh}_3)_2\}]$ (**2**) was obtained. The NMR spectra of the residue obtained from the reaction mixture after removal of solvent by evaporation showed the presence of the two isomers, **2a** and **2b** (Scheme 1). After



Scheme 1. Synthetic routes for mixed-metal tellurido clusters.

crystallization of the residue from a mixture of benzene and hexane, only the major isomer, **2a**, was isolated as single crystals and fully characterized by X-ray diffraction.^[13] As shown in Figure 2, **2a** has a triangular IrPt_2 core with two capped tellurido ligands, for which, separations of Ir(1) from Pt(1) and Pt(2) by 3.8049(5)–3.8588(6) Å as well as Pt...Pt interatomic distances of 3.1539(6) and 3.2523(7) Å indicate the absence of any metal–metal bonding interactions. The Ir center has a three-legged geometry, while the geometry around two Pt atoms is square-planar. Two phosphine ligands are mutually *trans* with respect to the Pt–Pt vector, as are the two Tol ligands. The X-ray structure was not available for **2b**, but its NMR data are analogous to those of **2a** and suggest a structure with two mutually *cis*-phosphine ligands (shown in Scheme 1 for **2b**). Because of steric repulsion between the two phosphine ligands, **2b** is the less-favored isomer. Note that **2a** redissolved in C_6D_6 and converted readily into an equilibrium mixture of **2a** and **2b** at room temperature, the ratio of which was 4:1, thus indicating that the isomerization of **2a** into **2b** is fast for this IrPt_2 cluster.

Treatment of **1** with two equivalents of $[\text{Pd}(\text{PPh}_3)_4]$ gave the Pd analogue of **2**, $[\{\text{Cp}^*\text{Ir}(\text{CO})\}(\mu_3\text{-Te})_2\{\text{PdTol}(\text{PPh}_3)_2\}]$




Figure 2. An ORTEP drawing of **2a**. Only one of the two independent molecules is shown (thermal ellipsoids shown at 30% probability). For clarity all the hydrogen atoms are omitted and only the carbon atom bond to P is shown for the phenyl groups of the PPh₃ units. Selected bond lengths [Å] and angles [°]: Ir–Te 2.6769(7)–2.6985(7), Pt–Te, 2.6347(8)–2.7052(7); Ir–Te–Pt, 89.82(2)–92.69(2), Pt–Te–Pt, 72.39(2)–75.23(2), Te–Ir–Te 75.26(2), 75.60(2), Te–Pt–Te, 75.86(2)–76.44(2). In isostructural **3a**: Ir–Pd 3.784(1)–3.837(1), Pd–Pd 3.185(2), 3.089(1), Ir–Te, 2.666(1)–2.689(1), Pd–Te, 2.605(1)–2.702(2), Ir–Te–Pd 89.49(4)–92.91(3), Pd–Te–Pd, 71.31(4)–74.12(4), Te–Ir–Te 75.68(3), 75.91(3), Te–Pd–Te, 76.60(4)–76.98(4).

(**3**), which is present as a mixture of **3a** and **3b** under the reaction conditions (Scheme 1). Workup of the product solution afforded only **3a** as single crystals and the X-ray analysis disclosed its structure to be analogous to that of **2a**.^[13] When redissolved in C₆D₆, **3a** slowly converted into an equilibrium mixture of **3a** and **3b** to reach a molar ratio of 3:1. This isomerization of the IrPd₂ cluster is considerably slower than that of the IrPt₂ cluster, **2a**. As addition of 0.1 equivalents of PPh₃ to the solution of **3a** significantly decreases the rate of the formation of **3b**, this isomerization presumably proceeds through dissociation of PPh₃. The much-faster isomerization of **2a** than that of **3a** may be ascribable to the higher ability of PPh₃ to dissociate from a Pt center than from a Pd center.

The reaction of **1** with one equivalent of [Pt(PPh₃)₃] in THF at room temperature resulted in the isolation of the intermediate in the conversion of **1** into **2**. Workup of the reaction mixture afforded the tellurido- and tellurolato-bridged dinuclear complex [Cp*Ir(CO)(μ-Te)(μ-TeTol)PtTol(PPh₃)] (**4**), which was isolated as a mixture of isomers, **4a** and **4b** (Scheme 1). For the major isomer, **4a**, the structure has been determined unambiguously by X-ray analysis of single crystals obtained after repeated crystallization of this mixture (Figure 3).^[13] Two metal centers are connected by one μ-telluride and one μ-tellurolate, in which the tellurido and Tol ligands around Pt are mutually *cis*. Oxidative addition of a Pt⁰ center across one Te–Tol bond in **1** and coordination of a Te atom of the other intact TeTol ligand to Pt affords **4a**. There are no bonding interactions between metal centers in **4a**. The Pt–Te(1) bond at 2.6563(7) Å is considerably longer than the Pt–Te(2) bond at 2.6013(7) Å presumably because of the strong *trans* influence exerted by the Tol ligand. Around the Ir center, the Ir–μ-tellurolato bond (Ir–Te(1) 2.6243(6) Å) is shorter than the Ir–μ-tellurido bond (Ir–Te(2) 2.6934(8) Å;




Figure 3. An ORTEP drawing of **4a**. Hydrogen atoms are omitted for clarity (thermal ellipsoids shown at 30% probability). Selected bond lengths [Å] and angles [°]: Ir–Pt 3.9198(5); Ir–Te(1)–Pt 95.98(2), Ir–Te(2)–Pt 95.50(2), Te–Ir–Te 76.38(2), Te–Pt–Te 77.42(2).

Figure 3). Unfortunately, an X-ray crystal structure was not obtainable for **4b**. However, spectroscopic data of **4b** are consistent with the structure shown in Scheme 1, in which only the geometry around Pt differs from that of **4a**. Complex **4a** dissolved in C₆D₆ and converted into a 2.2:1 mixture of **4a** and **4b** in equilibrium after 30 h at room temperature. A similar isomerization was reported for the reaction of [Pt(PEt₃)₃] with *i*PrTePh, in which the initial product *cis*-[PtPh(Te*i*Pr)(PEt₃)₂] transformed smoothly into *trans*-[PtPh(Te*i*Pr)(PEt₃)₂].^[11] Treatment of an equilibrium mixture of **4a** and **4b** with one equivalent of [Pt(PPh₃)₃] afforded a mixture of **2a** and **2b**, which indicates that the complexes **4a** are the intermediates in the reaction course of **1** to **2**.

It is noteworthy that from the reaction of **1** with [Pd(PPh₃)₄], the telluride–tellurolato complex corresponding to **4** was not obtained. The reaction of equimolar **1** and [Pd(PPh₃)₄] resulted in the formation of a mixture of the trinuclear tellurido cluster, **3**, and residual **1**, as indicated by the NMR spectra of the reaction mixture recorded soon after mixing.

It was also found that from the reaction of the Ir–Pt complex, **4**, with [Pd(PPh₃)₄], another trinuclear cluster [(Cp*Ir(CO))(μ₃-Te)₂{PtTol(PPh₃)}{PdTol(PPh₃)}] (**5**) was available through the facile insertion of a Pd⁰ center into the remaining Te–Tol bond in complex **4** (Scheme 1). The NMR spectra of the reaction mixture showed the presence of **5a**. After removal of the solvent by evaporation, crystallization of the residue from a mixture of THF and hexane afforded **5a** as single crystals. The NMR spectra of **5a** in C₆D₆ indicated its rapid transformation into an equilibrium mixture of **5a** and **5b** in a 3:1 molar ratio. The X-ray crystal structure of **5a** is depicted in Figure 4.^[13] Cluster **5a** has the expected triangular core capped by two μ₃-tellurido ligands, in which the Pd atom occupies the vertex close to the CO ligand, presumably because the insertion of the Pd center into the Te–Tol bond occurs selectively at the sterically less-crowding side with respect to the IrPtTe₂ core in **4**, namely, from the direction opposite the Cp* ligand. Two PPh₃ ligands are mutually *trans* as observed for **2a** and **3a**, and similarly the structure of **5b** may be assigned as that with two PPh₃ ligands mutually *cis* around the Pt–Pd vector.




Figure 4. An ORTEP drawing of **5a**. Only one of the two independent molecules is shown (thermal ellipsoids shown at 30% probability). For clarity all the hydrogen atoms are omitted and only the carbon atom bound to P is shown for the phenyl groups of the PPh₃ units. Selected bond lengths [Å] and angles [°]: Ir...Pt, 3.8534(6), 3.8505(6), Ir...Pd 3.7821(8), 3.7879(7), Pt...Pd 3.2002(9), 3.1053(7), Ir-Te 2.6712(8)–2.6870(8), Pt-Te 2.6162(8)–2.6827(8), Pd-Te 2.6179(9)–2.718(1), Ir-Te-Pt 91.72(2)–93.52(2), Ir-Te-Pd 89.22(3)–91.47(3); Pt-Te-Pd, 71.39(2)–74.34(2), Te-Ir-Te 75.43(2), 75.85(2), Te-Pt-Te 76.44(2), 76.72(2), Te-Pd-Te 75.88(3), 76.55(2).

In conclusion, a versatile route for synthesizing mixed-metal tellurido clusters in solution has been developed, which involves the use of a readily available tellurolato complex as a metal–tellurido building block through Te–C bond scission with M⁰ centers (M = Pt, Pd). Note that the reactions of **1** with Pt^{II} and Pd^{II} complexes [MCl₂(cod)] (cod = cyclooctadiene) result in the formation of only the tellurolato-bridged dinuclear and trinuclear complexes [Cp*Ir(CO)(μ-TeTol)₂MCl₂] (M = Pt, Pd) and [(Cp*Ir(CO)(μ-TeTol))₂Pt], which will be described elsewhere. Application of the method reported herein to the synthesis of a wide range of mixed-metal tellurido clusters is now in progress.

Experimental Section

All manipulations were carried out under an atmosphere of N₂ by using Schlenk techniques. Complexes [Cp*Ir(CO)₂],^[15] [Pt(PPh₃)₃],^[16] [Pd(PPh₃)₄],^[17] and TolTeTeTol^[18] were prepared by reported methods.

1: TolTeTeTol (230 mg, 0.526 mmol) was added to a solution of [Cp*Ir(CO)₂] (191 mg, 0.498 mmol) in benzene (10 mL) at room temperature. The reaction mixture was stirred for 3 h at room temperature and the resultant deep red solution was concentrated in vacuo. Addition of hexane afforded **1** as deep red crystals (315 mg, 79% yield). IR (KBr): $\tilde{\nu}$ = 1982 cm⁻¹, ν (CO); ¹H NMR (C₆D₆): δ = 1.55 (s, 15H, Cp*), 2.06 (s, 6H, Me in Tol), 6.75, 7.96 ppm (d, *J* = 7.9 Hz, 4H each, C₆H₄); elemental analysis (%) calcd for C₂₅H₂₉OIrTe₂: C 37.87, H 3.69; found: C 37.51, H 3.53.

2: [Pt(PPh₃)₃] (101 mg, 0.103 mmol) was added to a THF solution (5 mL) of **1** (41 mg, 0.051 mmol) and the mixture was stirred for 26 h at room temperature. The resultant red solution was evaporated to dryness in vacuo. The ¹H NMR spectrum of this residue indicated the

presence of two products, **2a** and **2b**, in a molar ratio of 2:1. Crystallization of the residue from a mixture of benzene and hexane (1:10) afforded pure **2a**·0.25 C₆H₆ as red crystals (43 mg, 48% yield). Complex **2a** dissolved in C₆D₆ and converted into an equilibrium mixture of **2a** and **2b** in the ratio of 4:1. **2a:** IR (KBr): $\tilde{\nu}$ = 1979 cm⁻¹, ν (CO); ¹H NMR (C₆D₆): δ = 1.70 (s, 15H, Cp*), 2.15, 2.21 (s, 3H each, Me in Tol), 6.73, 6.85, 7.53, 7.60 (d, *J* = 7.6 Hz, 2H each, C₆H₄), 7.0–7.1 (m, 18H, Ph), 7.8–8.0 ppm (m, 12H, Ph); ³¹P{¹H} NMR (C₆D₆): δ = 7.1 (d with ¹⁹⁵Pt satellites, *J*_{P-P} = 5 Hz, *J*_{Pt-P} = 3790 Hz), 7.4 ppm (d with ¹⁹⁵Pt satellites, *J*_{P-P} = 5 Hz, *J*_{Pt-P} = 3700 Hz); elemental analysis (%) calcd for C_{62.5}H_{60.5}OIrP₂Pt₂Te₂: C 43.46, H 3.53; found: C 43.65, H 3.52. Data assignable to **2b:** ¹H NMR (C₆D₆): δ = 1.71 (s, 15H, Cp*), 2.14, 2.16 ppm (s, 3H each, Me in Tol); ³¹P{¹H} NMR (C₆D₆): 6.6 (d with ¹⁹⁵Pt satellites, *J*_{P-P} = 8 Hz, *J*_{Pt-P} = 3770 Hz), 6.9 ppm (d with ¹⁹⁵Pt satellites, *J*_{P-P} = 8 Hz, *J*_{Pt-P} = 3930 Hz).

3: [Pd(PPh₃)₄] (116 mg, 0.100 mmol) was added to a THF solution (5 mL) of **1** (40 mg, 0.050 mmol) and the resultant black solution was stirred at room temperature for 23 h. The NMR spectrum of the reaction mixture showed the presence of two isomeric products, **3a** and **3b**, in a 3:1 ratio. Hexane was added to the concentrated product solution and kept at –20 °C to yield black crystals of **3a**·0.25 C₆H₁₄ (32 mg, 41% yield). When redissolved in C₆D₆, **3a** was converted very slowly into an equilibrium mixture of **3a** and **3b**. The rate constants for the conversion of **3a** into **3b** and its reverse reaction at 293 K were calculated to be 4.5 × 10⁻⁵ s⁻¹ and 13 × 10⁻⁵ s⁻¹, respectively, based on the ¹H NMR spectral change. IR (KBr): $\tilde{\nu}$ = 1971 cm⁻¹, ν (CO); elemental analysis (%) calcd for C_{62.5}H_{62.5}OIrP₂Pd₂Te₂: C 48.37; H 4.06; found: C 47.64, H 3.83. Data assignable to **3a:** ¹H NMR (C₆D₆): δ = 1.71 (s, 15H, Cp*), 2.14, 2.23 (s, 3H each, Me in Tol), 6.73, 6.78, 7.40, 7.48 (d, *J* = 8.0 Hz, 2H each, C₆H₄), 7.0–7.1 (m, 18H, Ph), 7.8–8.0 ppm (m, 12H, Ph); ³¹P{¹H} NMR (C₆D₆): δ = 14.7 (d, *J* = 7 Hz), 14.9 ppm (d, *J* = 7 Hz). Data assignable to **3b:** ¹H NMR (C₆D₆): δ = 1.75 (s, 15H, Cp*), 2.13, 2.16 ppm (s, 3H each, Me in Tol); ³¹P{¹H} NMR (C₆D₆): δ = 14.7 (d, overlapping with the signal of **3a**), 15.5 ppm (d, *J* = 8 Hz).

4: A mixture of **1** (160 mg, 0.202 mmol) and [Pt(PPh₃)₃] (200 mg, 0.204 mmol) in THF (5 mL) was stirred at room temperature for one day and the resultant dark green solution was evaporated to dryness in vacuo. The ¹H NMR spectra of the residue showed the presence of the products **4a** and **4b** in a molar ratio of 3:1. The residue was extracted with benzene and the addition of hexane to the concentrated extract afforded a 3:1 mixture of **4a** and **4b** as dark green crystals (177 mg, 70% combined yield). Single crystals of pure **4a** were obtained by repeated fractional crystallization from a mixture of benzene and hexane. **4a** was dissolved in C₆D₆ and allowed to stand for 30 h at room temperature; the resulting solution contained a mixture of **4a** and **4b** in equilibrium in a ratio of 2.2:1 as confirmed by ¹H NMR spectroscopy. IR (KBr): $\tilde{\nu}$ = 1991 cm⁻¹, ν (CO); elemental analysis (%) calcd for C₄₃H₄₄OIrPPtTe₂: C 41.31, H 3.55; found: C 41.06, H 3.69. Data assignable to **4a:** ¹H NMR (C₆D₆): δ = 1.63 (s, 15H, Cp*), 1.97, 2.10 (s, 3H each, Me in Tol), 6.59, 7.19, 7.39, 7.71 ppm (d, *J* = 8.0 Hz, 2H each, C₆H₄); ³¹P{¹H} NMR (C₆D₆): δ = 9.5 ppm (s with ¹⁹⁵Pt satellites, *J*_{Pt-P} = 3910 Hz). Data assignable to **4b:** ¹H NMR (C₆D₆): δ = 1.68 (s, 15H, Cp*), 1.97, 2.12 ppm (s, 3H each, Me in Tol); ³¹P{¹H} NMR (C₆D₆): δ = 20.4 ppm (s with ¹⁹⁵Pt satellites, *J*_{Pt-P} = 3900 Hz).

5: [Pd(PPh₃)₄] (58 mg, 0.050 mmol) was added to a THF solution (5 mL) of **4a** and **4b** (65 mg, 0.052 mmol) and this mixture was stirred for 5 h at room temperature. The NMR spectra of the reaction mixture showed the presence of **5a**. Addition of hexane to the concentrated product solution afforded **5a**·0.25 C₆H₁₄ as red crystals (40 mg, 46% yield). When dissolved in C₆D₆, **5a** converted into an equilibrium mixture of **5a** and **5b** at room temperature, the ratio of which was 3:1. IR (KBr): $\tilde{\nu}$ = 1968 cm⁻¹, ν (CO); elemental analysis (%) calcd for C_{62.5}H_{62.5}OIrP₂PdPtTe₂: C 45.76, H 3.84; found: C 45.55, H 3.76. Data assignable to **5a:** ¹H NMR (C₆D₆): δ = 1.72 (s, 15H, Cp*), 2.13, 2.21 (s, 3H each, Me in Tol), 6.72, 6.86, 7.38, 7.65 (d, *J* =

7.6 Hz, 2H each, C₆H₄), 7.0–7.1 (m, 18H, Ph), 7.8–8.0 ppm (m, 12H, Ph); ³¹P{¹H} NMR (C₆D₆): δ = 11.0 (d with ¹⁹⁵Pt satellites, J_{P-P} = 6 Hz, J_{P-P} = 3650 Hz), 11.9 ppm (d, J_{P-P} = 6 Hz). Data assignable to **5b**: ¹H NMR (C₆D₆): δ = 1.69 (s, 15H, Cp*), 2.15, 2.23 ppm (s, 3H each, Me in Tol); ³¹P{¹H} NMR (C₆D₆): δ = 10.8 (d, J_{P-P} = 6 Hz; ¹⁹⁵Pt satellites could not be resolved), 11.4 ppm (d, J_{P-P} = 6 Hz).

X-ray crystallography: Single crystals of **1**, **2a**·0.25 C₆H₆, **3a**·0.25 C₆H₁₄, **4a**, and **5a**·0.25 C₆H₁₄ were sealed in glass capillaries under argon and mounted on a Rigaku Mercury-CCD diffractometer equipped with a graphite-monochromatized MoK_α source. All diffraction studies were carried out at room temperature.

Structure solution and refinements were carried out by using the CrystalStructure program.^[19] The positions of the non-hydrogen atoms were determined by Patterson methods (PATTY^[20]) and subsequent Fourier synthesis (DIRDIF 99^[21]). Non-hydrogen atoms were refined anisotropically by minimizing wR2 based on all data, for which full-matrix least-squares techniques are employed. All hydrogen atoms were placed at the calculated positions and included at the final stages of the refinements with fixed parameters.

Received: June 2, 2006

Revised: August 9, 2006

Published online: October 19, 2006

Keywords: cluster compounds · iridium · palladium · platinum · tellurium

2.048 g cm⁻³, R1 = 0.0360 (3428 data with I > 2σ(I)) and wR2 = 0.0970 (all 5887 unique data) for 291 variables, GOF = 1.031. **2a**·0.25 C₆H₆: M_r = 1727.21, 0.40 × 0.20 × 0.10 mm³, triclinic, space group P $\bar{1}$, a = 14.987(3), b = 18.459(4), c = 22.293(5) Å, α = 97.113(3), β = 102.993(3), γ = 96.124(3)°, V = 5905(2) Å³, Z = 4, room temperature, ρ_{calcd} = 1.943 g cm⁻³, R1 = 0.0380 (17619 data with I > 2σ(I)) and wR2 = 0.1290 (all 26675 unique data) for 1351 variables, GOF = 1.019. **3a**·0.25 C₆H₁₄: M_r = 1551.85, 0.20 × 0.10 × 0.05 mm³, triclinic, space group P $\bar{1}$, a = 14.966(6), b = 18.422(6), c = 22.259(9) Å, α = 97.023(5), β = 103.306(6), γ = 95.816(5)°, V = 5874(4) Å³, Z = 4, room temperature, ρ_{calcd} = 1.755 g cm⁻³, R1 = 0.0660 (10818 data with I > 2σ(I)) and wR2 = 0.1930 (all 25822 unique data) for 1395 variables, GOF = 1.008. **4a**: M_r = 1250.30, 0.50 × 0.15 × 0.02 mm³, monoclinic, space group P2₁/a, a = 17.857(3), b = 10.702(2), c = 22.276(4) Å, β = 104.8625(7)°, V = 4115(1) Å³, Z = 4, room temperature, ρ_{calcd} = 2.018 g cm⁻³, R1 = 0.0410 (6319 data with I > 2σ(I)) and wR2 = 0.1300 (all 9381 unique data) for 486 variables, GOF = 1.004. **5a**·0.25 C₆H₁₄: M_r = 1640.54, 0.30 × 0.25 × 0.15 mm³, triclinic, space group P $\bar{1}$, a = 14.939(2), b = 18.439(3), c = 22.229(3) Å, α = 96.981(2), β = 103.311(2), γ = 95.562(2)°, V = 5865(2) Å³, Z = 4, room temperature, ρ_{calcd} = 1.858 g cm⁻³, R1 = 0.0560 (16567 data with I > 2σ(I)) and wR2 = 0.1650 (all 25735 unique data) for 1355 variables, GOF = 1.027. CCDC-616845–616849 (**1**, **2a**, **3a**, **4a**, and **5a**, respectively) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

- [1] a) J. Wachter, *Eur. J. Inorg. Chem.* **2004**, 1367; b) D. Fenske, J. F. Corrigan, *Metal Clusters in Chemistry, Vol. 3* (Eds: P. Braunstein, L. A. Oro, P. R. Raithby), Wiley-VCH, Weinheim, **1999**, chap. 4.2; c) I. Dance, K. Fisher, *Prog. Inorg. Chem.* **1994**, *41*, 637; d) L. C. Roof, J. W. Kolis, *Chem. Rev.* **1993**, *93*, 1037; e) M. A. Ansari, J. A. Ibers, *Coord. Chem. Rev.* **1990**, *100*, 223.
- [2] Precedent tellurido clusters were obtained by using either the organotellurium reagents such as R₃P=Te and (Me₃Si)₂Te or the inorganic tellurium sources, for example, Te powder, Te_n²⁻ (n = 2, 4), NaTeH.^[1]
- [3] a) M. Hidai, Y. Mizobe, *Can. J. Chem.* **2005**, *83*, 358; b) M. Hidai, S. Kuwata, Y. Mizobe, *Acc. Chem. Res.* **2000**, *33*, 46.
- [4] a) W.-Y. Yeh, H. Seino, T. Amitsuka, S. Ohba, M. Hidai, Y. Mizobe, *J. Organomet. Chem.* **2004**, *689*, 2338; b) H. Seino, T. Masumori, M. Hidai, Y. Mizobe, *Organometallics* **2003**, *22*, 3424; c) S. Nagao, H. Seino, M. Hidai, Y. Mizobe, *J. Organomet. Chem.* **2003**, *669*, 124, and references therein.
- [5] L. Y. Goh, W. Chen, *J. Chem. Soc. Dalton Trans.* **1994**, 2697.
- [6] Y.-C. Shi, *Polyhedron* **2004**, *23*, 1663.
- [7] H. Egold, U. Flörke, *Z. Anorg. Allg. Chem.* **2001**, *627*, 2295.
- [8] R. Oilunkaniemi, R. S. Laitinen, M. Ahlgren, *J. Organomet. Chem.* **2001**, *623*, 169.
- [9] T. Langetepe, D. Fenske, *Z. Anorg. Allg. Chem.* **2001**, *627*, 820.
- [10] a) J. F. Corrigan, D. Fenske, *Angew. Chem.* **1997**, *109*, 2070; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 1981; b) D. Fenske, N. Zhu, *J. Cluster Sci.* **2000**, *11*, 135.
- [11] For example: a) H. Kuniyasu, A. Ohtaka, T. Nakazono, M. Kinomoto, H. Kurosawa, *J. Am. Chem. Soc.* **2000**, *122*, 2375; b) H. Kuniyasu, T. Kato, M. Inoue, J. Terao, N. Kambe, *J. Organomet. Chem.* **2006**, *691*, 1873; c) R. Hua, H. Takeda, S. Onozawa, Y. Abe, M. Tanaka, *J. Am. Chem. Soc.* **2001**, *123*, 2899; d) L. A. Goj, M. Lail, K. A. Pittard, K. C. Riley, T. B. Gunnoe, J. L. Petersen, *Chem. Commun.* **2006**, 982, and references therein.
- [12] L.-B. Han, N. Choi, M. Tanaka, *J. Am. Chem. Soc.* **1997**, *119*, 1795.
- [13] X-ray data for **1**: M_r = 792.92, 0.20 × 0.10 × 0.10 mm³, monoclinic, space group P2₁/n, a = 7.815(2), b = 12.696(3), c = 26.055(6) Å, β = 95.889(1)°, V = 2571(1) Å³, Z = 4, room temperature, ρ_{calcd} =
- [14] M. Herberhold, G.-X. Jin, A. L. Rheingold, *J. Organomet. Chem.* **1998**, *570*, 241.
- [15] J. W. Kang, K. Moseley, P. M. Maitlis, *J. Am. Chem. Soc.* **1969**, *91*, 5970.
- [16] R. Ugo, F. Cariati, G. La Monica, *Inorg. Synth.* **1968**, *11*, 105.
- [17] D. R. Coulson, *Inorg. Synth.* **1972**, *13*, 121.
- [18] W. S. Haller, K. J. Irgolic, *J. Organomet. Chem.* **1972**, *38*, 97.
- [19] D. J. Watkin, C. K. Prout, J. R. Carruthers, P. W. Betteridge, CRYSTALS Issue 10, CrystalStructure 3.6.0: Crystal Structure Analysis Package, Rigaku and Rigaku/MS, **2000–2004**, 9009 New Trails Dr., The Woodlands, TX 77381, USA.
- [20] P. T. Beurskens, G. Admiraal, G. Beurskens, W. P. Bosman, S. Garcia-Granda, R. O. Gould, J. M. M. Smits, C. Smykall, PATTY, The DIRDIF program system; Technical Report of the Crystallography Laboratory: University of Nijmegen, Nijmegen, The Netherlands, **1992**.
- [21] P. T. Beurskens, G. Admiraal, G. Beurskens, W. P. Bosman, R. de Gelder, R. Israel, J. M. M. Smits, DIRDIF99, The DIRDIF 99 Program System, Technical Report of the Crystallography Laboratory: University of Nijmegen, Nijmegen, The Netherlands, **1999**.