

A One-pot Synthesis of Cyclic Pyrido[1,2-*a*]quinoxaline Phosphate, a New Molecule of Biological Importance from a Quinoxaline Derivative of Sugar


Shyamaprosad Goswami* and Avijit Kumar Adak

Department of Chemistry, Bengal Engineering College (Deemed University), Howrah-711 103, West Bengal, India

(Received April 4, 2003; CL-030295)


A five membered cyclic pyrido[1,2-*a*]quinoxaline phosphate has been first synthesized in one step in a good yield by phosphorylation of the quinoxaline derivative of sugar **1a** with (PhO)P(O)Cl₂. Phosphorylation of the tetrols (**1a** and **1b**) and the diol acetals with other phosphorylating agents, however, affords a dehydration product 2-(2-furyl)quinoxaline.

Quinoxaline and their derivatives have found use in antitumor antibiotics,^{1a-c} potent bacteriocides,^{1d} molecular recognition research,^{1e} colorimetric sensors^{1f} and consequently they have become the subject of intense research for their biochemical significance.^{1g,h} 2-(D-arabino-Tetrahydroxybutyl)quinoxaline **1a** is a long known molecule² and has been used as a pteridine analogue³ for synthetic studies on the molybdenum cofactor.⁴ Synthetic phosphoric esters of 6-hydroxyquinoxaline and ester derivatives of 2-hydroxyquinoxaline have also found use in insecticidal preparations and anthelmintics.^{1g} Cyclic phosphates are important constituents and are present in a large number of biologically important molecules.⁵ However, no report has been made for the synthesis of a cyclic phosphate onto a quinoxaline system. We wish to report a simple and an efficient synthesis of such a new heterocyclic phosphate triester where the phosphate moiety is present in a five membered ring which is attached to the pyrido[1,2-*a*]quinoxaline system.



In continuation⁵ of our work, we have investigated the possibility of phosphorylation reaction with the tetrol **1a** to obtain an α -hydroxyphosphate where the phosphate would be present in a six membered ring under a variety of reaction conditions. Thus when the tetrol **1a** is subjected to reaction with (PhO)P(O)Cl₂, 16-phenoxy-13*H*-15,17-dioxo-6,9-diaza-16-phosphacyclopenta[*a*]phenanthrene-16-oxide **2**, an unprecedented five membered cyclic phosphate, was obtained in a good yield. This one step procedure thus efficiently creates both six and five membered fused heterocyclic rings onto a quinoxaline system resulting a novel cyclic pyrido[1,2-*a*]quinoxaline phosphate by ring closure followed by dehydration of the tetrol **1a** with (PhO)P(O)Cl₂ (Scheme 1).

However, treatment of **1a-b** with (EtO)₃P(O) or POCl₃, (PhO)P(O)Cl₂, (*p*-O₂N-C₆H₄O)P(O)Cl₂, 2-phenyl-bis-triazolo-




Scheme 1. Reagent and conditions: i. (PhO)P(O)Cl₂ neat, rt, 5h. ii. (EtO)₃P(O) neat or POCl₃/(PhO)P(O)Cl₂/(*p*-O₂N-C₆H₄O)P(O)Cl₂/2-phenyl-bis-triazoloyl phosphate/tris(1,2,4-triazoloyl) phosphate in dioxane/CH₃CN and Et₃N or in pyridine, rt, 4-6 h. iii. H⁺, rt, 4-6 h. iv. dry acetone, conc. H₂SO₄, rt, 12 h.

yl phosphate^{6a} and tris(1,2,4-triazoloyl) phosphate^{6b} with various solvents containing Et₃N such as dioxane, CH₃CN and also in pyridine resulted a product which is solely found to be 2-(2-furyl)quinoxaline **3**. Acid treatment (concentrated H₂SO₄, concentrated HCl, HCl in absolute alcohol as well as AcOH and H₃PO₄) of **1a-b** also led to the formation of **3**, previously which was however wrongly assigned as glucazidone **4**.⁷ We also first report here the complete spectral proof (¹H, ¹³C, MS, and IR data) in favour of the structure **3**.

In order to rationalize the role of these phosphorylating agents, we have tested the phosphorylation reactions on diol acetals (**5a** and **5b**) under the similar conditions and we isolated **3** as the sole product in each case in good yields. From the above findings (Scheme 1), we have reasoned that instead of phosphorylation, triethyl phosphate, triazolides and other phosphorus acid chlorides that are used, favoured the side chain dehydration of the tetrols (**1a** and **1b**) and their diol acetals (**5a** and **5b**) resulting the formation of furan ring attached to quinoxaline **3**. The five membered pyrido[1,2-*a*]quinoxaline phosphate **2** was synthesized in one step from the tetrol **1a** by reaction with neat (PhO)P(O)Cl₂ in a good yield (85%).

The structure **2** is established⁹ on the basis of detailed spectral analysis (FTIR, ¹H, ¹³C (normal & DEPT), ³¹P and MS) which is also supported from the following plausible mechanism (Scheme 2). Phenylphosphorodichloridate phosphorylates the tetrol **1a** while initially attacking the terminal alcohol^{6a} which follows cyclization to **6a**, through pyrazine ring nitrogen or cyclization followed by dehydrations to **3** through benzylic hydroxyl group. Dehydration of **6a** follows the stereoelectronically more favourable path **a** than path **b** (to **6b**) which, may give **2** exclusively.



Scheme 2. Plausible mechanism for the formation of **2**.

The ^1H NMR spectrum however shows a *cis* coupling constant ($J = 6.9$ Hz) for the vicinal olefin protons which is consistent with the structure **2**. Further, DEPT-135 experiment has also suggested that the double bonds are present between $\text{C}_{11}\text{-C}_1$ and $\text{C}_3\text{-C}_4$ atoms. The ESI MS shows an intense peak at m/z 353.9 (MH^+). Thus, careful analysis of all the spectral data confirms the structure **2** for the product.

X-ray crystallographic studies^{10a} of several five membered cyclic phosphate esters have established that alkoxy group on phosphorus is directed away from the ring. The large negative (-30.12 ppm) ^{31}P chemical shift value of **2** and those found in other five membered cyclic phosphate esters^{10b,c} suggest that the compound **2** exists as a single conformer at room temperature.

In conclusion, the first one-pot synthesis of a cyclic pyrido[1,2-*a*]quinoxaline phosphate **2**, a new heterocyclic phosphate triester of biological significance is reported by a simple and an efficient method. Interestingly, on reaction with acids and other phosphorylating agents, the tetrols and the diol acetals formed **3** as the sole product.

We acknowledge the Bose Institute, Kolkata for the paid facility of a Bruker 500 MHz NMR instrument and Professor Thomas Schrader of Philipps Universitat, Marburg, Germany for the mass spectra. AKA thanks CSIR, Govt. of India for a Senior Research Fellowship.

References and Notes

- a) M. J. Waring and K. R. Fox, in "Molecular Aspects of Anti-cancer Drug Action," ed. by S. Niedle and M. J. Waring, Macmillan, New York (1983), p 127. b) M. J. Waring, in "Antibiotic," ed. by F. E. Hahn, Springer-Verlag, New York (1979), Vol. 5, p 173. c) G. W. H. Cheeseman and R. F. Cookson, *Chem. Heterocycl. Compd.*, **35**, 1 (1979). d) A. R. Katritzky and A. J. Boulton, *Adv. Heterocycl. Chem.*, **22**, 367 (1978). e) J. S. Nowick, P. Ballester, F. Ebmeyer, and J. Rebek, Jr., *J. Am. Chem. Soc.*, **112**, 8902 (1990). f) C. B. Black, B. Andrioletti, A. C. Try, C. Ruiperez, and J. L. Sessler, *J. Am. Chem. Soc.*, **121**, 10438 (1999). g) N. Sato, in "Comprehensive Heterocyclic Chemistry," ed. by A. R. Katritzky, W. C. Rees, and F. V. E. Scriven, Pergamon, Oxford (1996), Vol. 6, p 236. h) A. E. A. Porter, in "Comprehensive Heterocyclic Chemistry," ed. by A. R. Katritzky and W. C. Rees, Pergamon, Oxford (1984), Vol. 3, p 157.
- a) P. Griess and G. Harrow, *Chem. Ber.*, **20**, 2811 (1887). b) D. Horton and M. J. Millar, *J. Org. Chem.*, **30**, 2457 (1965). c) S. Gerchakov, P. J. Whitman, and H. P. Schultz, *J. Med. Chem.*, **9**, 266 (1966).
- a) L. Larson, D. J. Rowe, C. D. Garner, and J. A. Joule, *J. Chem. Soc., Perkin Trans. 1*, **1989**, 2317. b) B. Bradshaw, A. Dinsmore, D. Collison, C. D. Garner, and J. A. Joule, *Chem. Commun.*, **1998**, 417. c) B. Bradshaw, D. Collison, C. D. Garner, and J. A. Joule, *Chem. Commun.*, **2001**, 123. d) For a recent paper in this series, see B. Bradshaw, A. Dinsmore, D. Collison, C. D. Garner, and J. A. Joule, *J. Chem. Soc., Perkin Trans. 1*, **2001**, 3232.
- a) J. A. Pateman, D. J. Coves, B. M. Rever, and D. B. Roberts, *Nature*, **201**, 58 (1964). b) R. Hille, *Chem. Rev.*, **96**, 2757 (1996). c) D. Collison, C. D. Garner, and J. A. Joule, *Chem. Soc. Rev.*, **1996**, 25. d) R. S. Pilato, K. A. Erickson, M. A. Greaney, E. I. Stiefel, S. P. Goswami, L. Kilpatrick, T. G. Spiro, E. C. Taylor, and A. L. Rhiengold, *J. Am. Chem. Soc.*, **113**, 9372 (1991). e) S. P. Goswami, *Heterocycles*, **35**, 1552 (1993). f) B. Bradshaw, A. Dinsmore, W. Ajana, D. Collison, C. D. Garner, and J. A. Joule, *J. Chem. Soc., Perkin Trans. 1*, **2001**, 3239. g) For the recent paper in this series, see B. Bradshaw, D. Collison, C. D. Garner, and J. A. Joule, *Org. Biomol. Chem.*, **1**, 129 (2003).
- S. P. Goswami and A. K. Adak, *Tetrahedron Lett.*, **43**, 503 (2002) and references cited therein.
- a) C. Broka, T. Hozumi, R. Arentzen, and K. Itakura, *Nucleic Acids Res.*, **8**, 5461 (1980). b) A. Kraszewski and J. Stawinski, *Tetrahedron Lett.*, **21**, 2935 (1980).
- a) A. G. Sanchez, M. Y. Antinolo, and A. G. Gonzalez, *An. R. Soc. Esp. Fis. Quim.*, **51B**, 431 (1954); *Chem. Abstr.*, **50**, 11078 (1958). b) A. G. Sanchez, A. G. Gonzalez, and M. Y. Antinolo, *An. R. Soc. Esp. Fis. Quim.*, **51B**, 659 (1955); *Chem. Abstr.*, **50**, 10108 (1958). c) K. Maurer, B. Schiedt, and H. Schroeter, *Ber.*, **68B**, 1716 (1935); *Chem. Abstr.*, **29**, 1822 (1935).
- Compounds **1** and **5** are prepared by the similar procedure reported in Refs. 2b and 3a respectively. Selected physical data for compound **5a**: mp 129-131 °C [lit^{3a} 145-146 °C]. ^1H NMR (CDCl_3 , 500 MHz): δ 9.0 (s, 1H, quinoxalin-2-yl), 8.16-8.14 (m, 2H), 8.07-8.05 (m, 2H), 5.24 (dd, 1H, $J = 2.0, 2.0$ Hz), 4.30-4.25 (m, 2H, OH and $\text{c}_4\text{-H}$), 4.19 (m, 1H), 4.09 (m, 2H), 3.07 (brs, OH), 1.46 and 1.38 ($2 \times$ s, 6H, $2 \times \text{CH}_3$). MS (FD): m/z (%): 290 (M^+ , 100), 275 (50), 189 (40). The spectroscopic data for compound **5b** is in accord with the structure given. For a recent preparation of **1a** and **1b** see: S. P. Goswami and A. K. Adak, *Tetrahedron Lett.*, **43**, 8371 (2002).
- A suspension of the tetrol **1a** (1.0 g, 3.98 mmol) and $(\text{PhO})\text{P}(\text{O})\text{Cl}_2$ (1.0 mL) was stirred under nitrogen atmosphere at ice bath (0-5 °C) for 1 h and then at rt for 4 h. Water (40 mL) was then added and the mixture was extracted with CH_2Cl_2 (2×50 mL). The organic layer was washed well with brine, dried (MgSO_4) followed by evaporation with rotary evaporator afforded a brown solid which was purified on silica gel (100-200 mesh) column chromatography eluting with petroleum ether : CH_2Cl_2 (3:2) gave **2** (0.88 g, 85%) as a brown crystalline solid. mp 84-86 °C. UV (CHCl_3) λ_{max} (log ϵ): 369 (4.09) and 284 (4.23) nm. FT-IR (KBr): 3137, 2925, 1612, 1588, 1552, 1497, 1297, 1189, 1083, 1058, 1002, 963, 917, 775, 758, 688, 594 cm^{-1} . ^1H NMR (CDCl_3 , 500 MHz): δ 9.23 (s, 1H, quinoxalin-2-yl), 8.09 (d, 1H, $J = 8.8$ Hz), 8.05 (d, 1H, $J = 8.8$ Hz), 7.75-7.70 (m, 2H), 7.67 (d, 1H, $J = 6.9$ Hz), 7.35-7.30 (m, 4H), 7.24 (d, 1H, $J = 8.4$ Hz), 7.20 (t, 1H, $J = 7.3$ Hz), 6.61 (qt, 1H, $J = 1.7, 1.7, 1.7$ Hz). ^{13}C NMR (CDCl_3 , 125 MHz): 152.00, 150.87 (d, $J = 7.3$ Hz), 145.49, 144.25, 142.48, 142.45, 141.68, 130.84, 130.25 (d, $J = 0.8$ Hz), 129.72, 129.61, 126.0 (d, $J = 1.2$ Hz), 120.54, 120.50, 112.90, 112.23. ^1H decoupled ^{31}P NMR (CDCl_3 , 202 MHz): -30.12. MS (ESI): m/z (%): 353.9 (MH^+ , 20), 349.0 (100), 327 (30), 196 (25). Calcd. for $\text{C}_{18}\text{H}_{13}\text{N}_2\text{O}_4\text{P}$: required C, 61.37; H, 3.72; N, 7.95%. Found C, 61.35; H, 3.78; N, 7.99%. Selected physical data for compound **3**: Yield: (80%); off-white solid; mp 76-78 °C. FT-IR (KBr): 3136, 3116, 1736, 1612, 1552, 1497, 1224, 1169, 1082, 916, 657, 594, 589 cm^{-1} . ^1H NMR (CDCl_3 , 500 MHz): δ 9.25 (s, 1H, quinoxalin-2-yl), 8.11 (d, 1H, $J = 8.3$ Hz), 8.07 (d, 1H, $J = 8.3$ Hz), 7.77-7.72 (m, 2H), 7.69 (d, 1H, $J = 7.3$ Hz), 7.33 (d, 1H, $J = 3.4$ Hz), 6.64 (qt, 1H, $J = 1.6, 1.7, 1.7$ Hz). ^{13}C NMR (CDCl_3 , 125 MHz): 151.97, 145.55, 144.23, 142.47, 142.43, 141.67, 130.93, 129.77, 129.61 (d, $J = 1.2$ Hz), 129.53, 112.88, 112.19. MS (ESI): m/z (%): 196 (M^+ , 100), 168 (30). Calcd. for $\text{C}_{12}\text{H}_8\text{N}_2\text{O}$: required C, 73.46; H, 4.11; N, 14.28%. Found C, 73.48; H, 4.15; N, 14.30%.
- a) M. G. Newton and B. S. Campbell, *J. Am. Chem. Soc.*, **96**, 7790 (1974) and references cited therein. b) F. Ramirez, A. V. Patwardhan, N. B. Dessai, and S. R. Heller, *J. Am. Chem. Soc.*, **87**, 549 (1965). c) G. M. Blackburn, J. S. Cohen, and L. Todd, *Tetrahedron Lett.*, **5**, 2873 (1964).