



A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

GDCh

International Edition

www.angewandte.org

Accepted Article

Title: Time-dependent Photodynamic Therapy for Multiple Targets: A Highly Efficient AIE-active Photosensitizer for Selective Bacterial Elimination and Cancer Cell Ablation

Authors: Ben Zhong Tang, Qiyao Li, Ying Li, Tianliang Min, Junyi Gong, Lili Du, David Lee Phillips, Junkai Liu, Jacky W. Y. Lam, Herman H. Y. Sung, Ian D. Williams, Ryan T. K. Kwok, Chun Loong Ho, Kai Li, and Jianguo Wang

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.201909706
Angew. Chem. 10.1002/ange.201909706

Link to VoR: <http://dx.doi.org/10.1002/anie.201909706>
<http://dx.doi.org/10.1002/ange.201909706>

RESEARCH ARTICLE

Time-dependent Photodynamic Therapy for Multiple Targets: A Highly Efficient AIE-active Photosensitizer for Selective Bacterial Elimination and Cancer Cell Ablation

Qiyao Li,^{†[a]} Ying Li,^{†[b][f]} Tianliang Min,^{†[c]} Junyi Gong,^[a] Lili Du,^[d] David Lee Phillips,^[d] Junkai Liu,^[a] Jacky W. Y. Lam,^[a] Herman H. Y. Sung,^[a] Ian D. Williams,^[a] Ryan T. K. Kwok,^[a] Chun Loong Ho,^[c] Kai Li,^[c] Jianguo Wang^{*[e]} and Ben Zhong Tang^{*[a][b][f]}

Abstract: Pathogen infection and cancer are the two major human health problems. In this work, we achieved an organic salt photosensitizer (PS), called 4TPA-BQ with aggregation-induced emission feature *via* one-step reaction. Owing to the aggregation-induced reactive oxygen species generation effect and sufficient small ΔE_{ST} , 4TPA-BQ shows a satisfactorily high 1O_2 generation efficiency of 97.8%. *In vitro* and *in vivo* experiments confirmed that 4TPA-BQ exhibited potent photodynamic antibacterial performance against ampicillin-resistant *Escherichia coli* with good biocompatibility in a short time (15 min). When the incubation time persisted long enough (12 h), cancer cells were ablated efficiently, leaving normal cells essentially unaffected. This is the first reported time-dependent fluorescence-guided photodynamic therapy in one individual PS for ordered and multiple targeting by varying the external conditions. This can update the design principle of efficient PSs in potential clinical applications.

Introduction

Pathogen infections cause severe illness and significant mortality globally. During the past decades, antibiotics have been employed predominately to treat diseases caused by bacteria since the first discovery of penicillin in 1928. However, after a long period abuse of antibiotics, the emergence of drug-resistant

bacteria has posed a severe threat to human health. A recent study reported that about 70% normal *E. coli* became drug resistant strains only after 3 h exposure to antibiotics.^[1] Currently, we are confronted with an intractable situation that new antibiotics development fails to keep pace with the generation of drug-resistant bacteria. Therefore, developing alternative antibacterial therapeutics is of utmost urgency and great importance to clinical application.

Cancer, meanwhile, is another major public health problem. Globally, about 1 in 6 deaths is due to cancer. In 2019, about 1,762,450 new cancer cases will be diagnosed and there will be 606,880 cancer deaths in the United States.^[2] To date, although various therapeutic modalities such as surgery, chemotherapy and radiotherapy have been exploited, they show different problems.^[3] Specifically, surgery operation is invasive to human body, taking chemotherapy drugs, such as Cisplatin can lead to the increased risk to get infection and drug resistance, and patients who receive radiotherapy treatment, may suffer from fatigue and loss of hair in the treatment area.

Recently, photodynamic therapy (PDT) has emerged as a promising method to combat pathogen bacteria as well as a powerful clinical protocol for cancer treatment.^[4] PDT employs photosensitizers (PSs) to produce destructive singlet oxygen (1O_2) or other reactive oxygen species (ROS) from endogenous molecular oxygen under light irradiation.^[5] Thus, combining the merits of PS and light irradiation, PDT exhibits several conspicuous advantages including noninvasiveness, no drug resistance, low cytotoxicity, selective targeting, spatiotemporal precision and synergistic effect over conventional therapeutic modalities. Among major achievements on PDT advance, fluorescence-guided PDT (FL-PDT), with the unique advantage of real-time monitoring, has attracted increasing attention. PS, which serves as the key component of FL-PDT, plays a decisive role in therapeutic effect.^[6] Most conventional organic PSs, however, encounter the problem of aggregation-caused quenching (ACQ) in biosystems, resulting in undesired weak emission and poor ROS generation. Luckily, the discovery of the aggregation-induced emission (AIE) phenomenon offers a convenient solution.^[7] Unlike ACQ luminophores, AIE luminogens are weakly emissive in dilute solutions. However, they show strong emission and ROS generation in the aggregate state, owing to the mechanism of restriction of intramolecular motion (RIM).^[8] Thus, AIE-based PSs hold promising potential in PDT applications. Although research on AIE-based PSs in anticancer and antibacterial treatments proceeds rapidly, various problems remain to be solved. One common problem is that to realize selective binding, specific targeting ligands are needed to introduce to PSs. Such a method is not only tedious in synthesis but also of high cost.^[9] Another big challenge is that for most

[a] Q. Li, J. Gong, J. Liu, Dr. J. W. Y. Lam, Dr. H. H. Y. Sung, Prof. I. D. Williams, Dr. R. T. K. Kwok, Prof. B. Z. Tang

Department of Chemistry, Hong Kong Branch of Chinese National Engineering Research Center for Tissue Restoration and Reconstruction, Institute for Advanced Study, Department of Chemical and Biological Engineering, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China
Email: tangbenz@ust.hk

[b] Dr. Y. Li, Prof. B. Z. Tang

Center for AIE Research, College of Materials Science and Engineering, Shenzhen University, Shenzhen 518060, China

[c] T. Min, Prof. K. Li, Prof. C. L. Ho

Department of Biomedical Engineering, Southern University of Science and Technology, Shenzhen, Guangdong 518055, China

[d] Dr. L. Du, Prof. D. L. Phillips

Department of Chemistry, The University of Hong Kong, Pokfulam Road, Hong Kong, China

[e] Prof. J. Wang

College of Chemistry and Chemical Engineering, Inner Mongolia University, Hohhot 010021, China
E-mail: wangjq@iccas.ac.cn

[f] Dr. Y. Li, Prof. B. Z. Tang

HKUST-Shenzhen Research Institute, No. 9 Yuexing First Road, South Area, Hi-tech Park, Nanshan, Shenzhen 518057, China


[†] These authors contributed equally to this work

Supporting information for this article is given via a link at the end of the document.

RESEARCH ARTICLE

existing PSs, it is quite difficult to manipulate their targets by merely adjusting the external conditions without changing their molecular structure. Thus, the target manipulation of a single PS without introducing a specific targeting group is worth exploring and is of great research significance. Based on the aforementioned situation, it is highly challenging but rewarding to develop a simple AIE-based PS with six conspicuous merits including: (1) facile preparation and stable chemical structure, (2) moderate water solubility, (3) high ROS efficiency with bright emission, (4) good biocompatibility, (5) high specificity to targets and (6) low dark toxicity but significant photo toxicity. More importantly, the achievement of multiple yet successive targeting in one individual structure-fixed PS is scarcely reported. That is to say: under a given condition, the PS can only recognize one specific target. However, by varying the condition, the target changes correspondingly. Thus, achieving the success to this target-manipulation by varying the external conditions in one individual PS not only opens a new door for PS design principle but also widens the applications for defending against pathogens and cancer.

In this contribution, motivated by anion- π interactions on triggering AIE effect,^[10] we designed and prepared an AIE-based PS, namely 4TPA-BQ via one-step synthesis. The inherent positive charge endows the molecule with modest water solubility for bioapplications. Simultaneously, 4TPA-BQ shows aggregation-induced ROS generation (AIROSG) effect, with a $^1\text{O}_2$ quantum yield of up to unity. Unprecedentedly, time-dependent PDT for multiple targets was achieved by controlling the incubation time of 4TPA-BQ with varied targets. In the presence of white light irradiation for 15 min, 4TPA-BQ could eliminate drug-resistant *Escherichia coli* infection efficiently, leaving host tissues unaffected. When the incubation time was lengthened to 12 h, 4TPA-BQ could kill cancer cells with slight harm to normal cells (Scheme 1).



Scheme 1. Illustration of 4TPA-BQ for time-dependent PDT.

Results and Discussion

The synthetic route of 4TPA-BQ was depicted in Figure 1A, 4,4'-(1,2-ethynediyl)bis[N,N-diphenylbenzenamine] (**1**) was synthesized via a modified Sonogashira coupling reaction according to the reported literature (Scheme S1) preliminarily,^[11] then a facile one-pot reaction of compound **1** and 2-methylallylamine produced 4TPA-BQ with a satisfactory yield of 81%.^[12] Detailed experimental procedures were provided in the Supporting Information. Both compound **1** and the product were fully characterized by ^1H NMR, ^{13}C NMR and high-resolution mass spectroscopies (Figures S1-S6).




Figure 1. (A) Synthetic route to 4TPA-BQ: (i) NaPF_6 , $[\text{Cp}^*\text{RhCl}_2]_2$, $\text{Cu}(\text{OAc})_2\cdot\text{H}_2\text{O}$, MeOH, 130°C , 6 h. (B) PL spectra of 4TPA-BQ in DMSO/water mixtures with different water fractions (f_w). (C) Plot of the relative emission intensity (I/I_0) versus f_w . Inset: fluorescence images of 4TPA-BQ in solution ($f_w = 0\%$) and aggregate ($f_w = 70\%$) states under irradiation from a hand-held UV lamp. Excitation wavelength: 380 nm; Concentration: 10 μM .

After confirming the structure of 4TPA-BQ, we investigated its optical properties. As shown in Figure S7, its UV-vis absorption spectrum presents two broad peaks at the wavelength of 375 nm and 440 nm in DMSO solution. Then photoluminescence (PL) spectrum was measured in DMSO/water mixtures to study its AIE property. As presented in Figures 1B, 1C and Table S1, when water fraction (f_w) is below 40%, the PL intensity is quite weak with negligible fluorescence quantum yield ($\Phi_{\text{Soln}} = 0.30\%$) and nearly no emission was observed by naked eyes under a hand-held UV light irradiation. However, upon increasing f_w to 50%, the emission intensity amplified sharply. An observation of bright orange fluorescence ($\Phi_{\text{Aggr.}} = 6.70\%$) was obtained at $f_w = 70\%$, in which emission intensity enhanced to about 40-fold compared to that of DMSO solution. The emission became weaker when f_w increased from 70%, which may be attributed to the effect of aggregate size (Figure S8). Generally, the larger the aggregate size, the brighter it is ($f_w = 70\%$, $D = 99$ nm; $f_w = 80\%$, $D = 76$ nm; $f_w = 90\%$, $D = 74$ nm; $f_w = 99\%$, $D = 71$ nm). Compared to the orange emission of forming aggregates, the solids exhibit a stronger and blue-shifted emission ($\Phi_{\text{Solid}} = 15.30\%$) (Figure S9), which is ascribed to the closely and ordered molecular packing, testified by powder X-ray

RESEARCH ARTICLE

diffraction (PXRD) analysis (Figure S10). Also, the UV-vis spectra of 4TPA-BQ in DMSO/water mixtures were measured and no obvious change was observed (Figure S11).

For in-depth understanding of the AIE feature of 4TPA-BQ, we obtained its single crystal in chloroform/hexane mixtures by slow evaporation (Table S2). It is noticed that anion- π^+ interactions between the fluorine atoms of hexafluorophosphate anions and the positively charged benzoquinoline core were found with the distance of 3.063 Å and calculated energy of -61.68 kJ/mol, which impedes π - π stacking in the crystal to avoid emission quenching (Figure 2A). Intramolecular and intermolecular hydrogen bonds of F-H interactions were also found in the crystal lattice, constraining rotation of the phenyl rings effectively, thus rigidifying structure in the crystal lattice (Figures 2A and 2C). Besides, the large dihedral angles between TPA moieties and benzoquinoline core were observed as 64.37° (θ_1), 82.61° (θ_2) and 66.49° (θ_3), further preventing detrimental π - π interactions (Figures 2B and S12). Consequently, anion- π^+ interactions together with highly twisted molecular conformation favors AIE characteristic of 4TPA-BQ, which agrees well with previously reported literature.^[10b]




Figure 2. (A) Anion- π^+ and F-H interactions, (B) dihedral angles, (C) intermolecular hydrogen bonding and (D) C-H... π interactions in the single crystal structure of 4TPA-BQ. E is defined as interaction energy between π^+ and anion calculated based on the single-crystal structure by single-point calculations using M062X/6-31+G (d,p) method.

Considering its typical D-A structure in favor of $^1\text{O}_2$ generation,^[13] preliminary experiments were performed to explore $^1\text{O}_2$ generation ability of 4TPA-BQ. Here, N,N'-di(2,3-dihydroxypropyl)-9,10-anthracenedipropanamide (DHPA) was utilized as $^1\text{O}_2$ indicator for the reason that (1) it can detect $^1\text{O}_2$ specifically, with an obvious decrease in absorbance; (2) it has no interaction with cationic PSs, thus this non-ionic chemical trap can evaluate $^1\text{O}_2$ production in a more accurate way.^[14] Firstly, we measured the UV-vis absorption spectra of DHPA in 4TPA-BQ aggregates and solution under white light irradiation, respectively (Figure S13). As presented in Figure 3A, it is worth noticing that the absorbance of DHPA decreased sharply in the presence of 4TPA-BQ aggregates, whereas irradiation on 4TPA-BQ solution did not lead to obvious DHPA consumption. This can be explained by that, on one hand, formation of aggregates can induce excited states energy splitting, thus promoting intersystem crossing (ISC) rate;^[15] on the other, according to RIM mechanism, aggregates will minimize the competitive nonradiative decay, which favors

ISC process. Thus, AIE feature can activate $^1\text{O}_2$ generation, which can be defined as AIROSG. Commercial available PSs including Chlorin E6 (Ce6) and Rose Bengal were also compared under the same circumstances (Figure S13). After 8 min exposure to white light, compared to more than 60% DHPA consumption in the presence of 4TPA-BQ aggregates, only 10% and 50% DHPA degraded in Ce6 and Rose Bengal, respectively (Figure 3A). The results reveal that 4TPA-BQ aggregates can generate $^1\text{O}_2$ more efficiently than Ce6 and Rose Bengal, both of which are commonly used and high $^1\text{O}_2$ production PSs for PDT.^[16] Furthermore, taking DHPA as an indicator and Rose Bengal as the standard PS, we calculated $^1\text{O}_2$ quantum yield of 4TPA-BQ in aqueous medium following the reported procedure.^[17] A high value of 97.8% was gained (Figures 3B and S14). The above results suggest that 4TPA-BQ in aqueous medium can produce $^1\text{O}_2$ at a high efficiency, rendering it a promising candidate for PDT application.




Figure 3. (A) Plot of relative absorbance of DHPA without and with different PSs under white light irradiation, where A_0 and A are the absorbance of DHPA at 378 nm before and after white light irradiation, respectively. [PS] = 9×10^{-6} M, [DHPA] = 4.5×10^{-5} M. (B) Decomposition rate of DHPA with 4TPA-BQ aggregates and RB, respectively. (C) Calculated energy level diagram between singlet and triplet states. (D) The natural transition orbitals (NTOs) of 4TPA-BQ.

For elucidation of its extremely high $^1\text{O}_2$ quantum yield, theoretical calculations were performed based on time-dependent density functional theory (TD-DFT) via Gaussian 09. As indicated in Figure 3C, notably, the energy gap between S_1 and T_4 state is only 0.03 eV, which is much smaller compared to most reported PSs.^[18] Apparently, for 4TPA-BQ, it is the sufficiently narrow ΔE_{ST} rather than SOC that plays the dominant role in enhancing ISC rate as well as high $^1\text{O}_2$ generation efficiency. Successively, natural transition orbitals (NTOs) were calculated for excited states analysis. As shown in Figure 3D, the hole NTOs are mainly localized on TPA moieties while the electron NTOs are mainly localized on benzoquinoline core. Thus, the highly separated charge distribution realized by incorporation of electron donor

RESEARCH ARTICLE

(TPA moieties) and acceptor (benzoquinoline core) contributes to the narrow energy gap between singlet and triplet states.^[19] Furthermore, to manifest the existence of triplet excited states which is pivotal for producing $^1\text{O}_2$, we performed the nanosecond transient absorption experiments (Figure S15). The determined triplet lifetime of 4TPA-BQ aggregates was 81 ns while no signal was detected for 4TPA-BQ in solution state, which again validates that formation of aggregates favors ISC process.

Since 4TPA-BQ is positively charged, it is expected to target both bacteria and cells *via* electrostatic interactions.^[20] Here, confocal laser scanning microscopy (CLSM) was utilized to verify the assumptions. *Staphylococcus epidermidis* (*S. epidermidis*) and *Escherichia coli* (*E. coli*) were selected as representatives for Gram-positive and Gram-negative bacteria, respectively. As illustrated in Figure S16, the bright orange fluorescence from 4TPA-BQ treated *E. coli* and *S. epidermidis* was readily observed with high contrast to background, indicating that 4TPA-BQ can bind to both gram-negative and gram-positive bacteria efficiently. Subsequently, normal mammalian cells including COS-7 and HLF cells were incubated with 4TPA-BQ under the same conditions (Figure S17). Unexpectedly, scarcely no fluorescence signals were detected under CLSM for these two kinds of cells, hinting that 4TPA-BQ cannot target normal cells. Thus, the bacterial and cell imaging results indicate that 4TPA-BQ has the potential for selectively recognizing and imaging bacteria over normal cells.

Inspired by the specific bacterial-targeting capability and high $^1\text{O}_2$ generation efficiency of 4TPA-BQ, we would like to study and compare its antibacterial activity and cytotoxicity, which were evaluated by plate count method and MTT assay, respectively. As presented in Figures 4A and 4B, 4TPA-BQ displays a dose-dependent antibacterial efficiency towards both *E. coli* and *S. epidermidis* under white light illumination. About 90% *E. coli* and *S. epidermidis* were killed at a low concentration of 2 μM ; and when 4TPA-BQ reached 5 μM , almost no bacterial colony could be observed on the agar for both *E. coli* and *S. epidermidis* (Figure 4D). Meanwhile, for groups incubated without irradiation, no obvious drop of survival rate was found with increased concentration of 4TPA-BQ, which suggests that the potent antimicrobial activity of 4TPA-BQ is entirely due to its intrinsic $^1\text{O}_2$ generation induced by white light irradiation. Considering its high killing efficiency on *E. coli* and *S. epidermidis*, we tried drug-resistant bacterial strains, ampicillin-resistant *E. coli*. As shown in Figure 4C, less than 30% ampicillin-resistant *E. coli* survived when treated with 5 μM 4TPA-BQ under white light irradiation. When increasing to 10 μM , the killing efficiency reached to almost 100%. Similarly, no dark toxicity to ampicillin-resistant *E. coli* was found. Therefore, when in the absence of irradiation, 4TPA-BQ can visualize bacteria *in situ*; When in the presence of irradiation, efficient $^1\text{O}_2$ generation ability endows 4TPA-BQ with powerful antibacterial capability, even including drug-resistant bacteria. To acquire solid evidence of its antibacterial activity, SEM analysis was conducted to visualize morphological changes of *E. coli* and *S. epidermidis* (Figure S18). For both control group and dark group, regular body shape and intact cell walls could be clearly observed. By contrast, for bacteria treated with both 4TPA-BQ and light, the walls were rigorously damaged, collapsed and fused. These direct information from SEM is in consistent with the antibacterial results. Subsequently, we investigated the toxicity of 4TPA-BQ towards normal cells. As illustrated in Figure 4E, for

both dark and illumination groups, no obvious drop of COS-7 cell viability was observed with increased 4TPA-BQ concentration. It is worth noticing that a highly active cell viability of 91% was maintained even under the high concentration of 10 μM , in sharp contrast to the antibacterial results. Analogous results were also achieved for HLF cells (Figure 4F). MTT results reveal that 4TPA-BQ displays excellent biocompatibility with normal cells. Thus, 4TPA-BQ can cause fatal damage to bacteria but negligible cytotoxicity to normal cells.




Figure 4. (A-C) Killing efficiency of 4TPA-BQ on *E. coli*, *S. epidermidis* and ampicillin-resistant *E. coli*. (D) Photographs of *E. coli* and *S. epidermidis* cultured on agar plate supplemented with 5 μM 4TPA-BQ. (E-F) Cell viability of COS-7 and HLF cells incubated with 4TPA-BQ for 15 min.




Figure 5. CLSM images of *E. coli* and HLF cells and *S. epidermidis* and HLF cells stained with PI after incubation with 5 μM 4TPA-BQ for 15 min under white light irradiation. The dash lines indicate HLF cells.

To further testify that 4TPA-BQ can realize the selective bacterial-targeting over normal cells, we cultured *E. coli* or *S. epidermidis* mixed with mammalian cells in the presence of 4TPA-BQ and light irradiation. Here, propidium iodide (PI) was utilized to mark the dead bacteria or cells. As shown in Figures 5 and S19, the detected red fluorescence of PI and orange fluorescence of 4TPA-BQ was from bacteria, whereas hardly no fluorescence was from mammalian cells, indicating that 4TPA-BQ killed the bacteria efficiently while normal cells were still alive. Hence, the co-culture experiments fully validated that 4TPA-BQ is competent to specifically and selectively recognize and kill bacteria upon light irradiation with normal mammalian cells essentially unaffected.

RESEARCH ARTICLE

In order to gain deep insight to selective bacterial targeting mechanism, zeta potential measurements, which offered the information of surface charges of bacteria and cells, were performed. The acquired data was summarized in Table S3. For untreated groups, the zeta potentials of *E. coli* and *S. epidermidis* are more negative than COS-7 cells and HLF cells, suggesting that selective binding is actuated by stronger electrostatic interactions with bacteria. After treated with 4TPA-BQ, the zeta potentials of cells and *S. epidermidis* barely changed, while *E. coli* become more positively surface charged. The unchanged surface charges of COS-7 and HLF cells revealed there is no interaction with 4TPA-BQ. The different zeta potential changes of *E. coli* and *S. epidermidis* are related to their cell wall architecture.^[21] For Gram-positive *S. epidermidis*, it has a thick but porous outer peptidoglycan wall and negatively charged teichoic acids inserting in the thick wall, thus 4TPA-BQ can be almost held in the pores without exposure to surface, leading to subtle change on *S. epidermidis* surface charge. As for Gram-negative *E. coli*, an additional membrane layer composed by negatively charged lipopolysaccharides (LPS) is located outside the peptidoglycan layer, making 4TPA-BQ stay on the surface, thus the surface charges of *E. coli* become more positive. The zeta potential data is in accordance with bacterial and cell imaging results.

Encouraged by its desired bacterial targeting and eliminating performance *in vitro*, we further evaluated the antibacterial ability of 4TPA-BQ against drug-resistant bacteria *in vivo*. We established a third-degree burn wounds infected with Ampicillin-resistant *E. coli* on mice model.^[22] After burn and infection, they were randomly assigned to 3 groups, treated with PBS only (Control group), 4TPA-BQ only (Dark group) and 4TPA-BQ with white light irradiation (PDT group), respectively. The macroscopic appearance of wound healing process was recorded at different time points. As shown in Figures 6A and 6B, at day 3 and 5 post burn, ulceration and purulence appeared on the wound surface in both Dark and Control groups, while the PDT group showed no such signs. At day 8, the sizes of Ampicillin-resistant *E. coli* infected wounds in the PDT group were significantly smaller in comparison to the other two groups. At day 10, the wounds in the PDT group had completely recovered with newly regenerated skins, while the wounds in the other two groups were relatively large with pus under the scab after uncovering. At day 5 and 10 post burn, we took the damaged skin tissues for bacterial culture and count (Figures 6C, S20 and S21A). It turned out that the number of bacteria in the PDT group was significantly less than that in the Control group and the Dark group. Similarly, at day 5 post burn, the liver and spleen were ground for bacteria counting. As presented in Figures 6C and S21B, there were almost no colonies in the organs from PDT group, displaying a sharp contrast to the Dark and Control groups. This result indicates that the PDT treatment had excellent germicidal efficacy and can effectively inhibit further infection of bacteria from the skin into deep organs.^[23] In addition, we further evaluated the wound healing efficacy of sectioned tissues by hematoxylin and eosin (H&E) staining (Figure 6D).^[24] At day 5, the intact and thick epidermis, as well as the hair follicles and blood vessels, were clearly observed in the neonatal skin of the PDT group. The neonatal structure recovered completely in the PDT group at day 10. However, in the Dark and Control groups, no epidermis appeared and almost no new hair follicles and blood vessels were

observed at day 5 and 10. Therefore, the *in vivo* experiments successfully demonstrated that 4TPA-BQ has excellent antimicrobial effect under white light irradiation that can significantly promote the wound healing process, consistent with *in vitro* tests.




Figure 6. Antibacterial activity against ampicillin-resistant *E. coli* infection *in vivo*. (A) Photographs of bacterial infected mice wounds treated with PBS only (Control group), 4TPA-BQ only (Dark group) and 4TPA-BQ+white light irradiation (PDT group), respectively. (B) Changes of the infected wound size subjected to different treatments. (C) Plate photographs of bacterial amount of skin wound and organs at day 5 in 100-fold dilution. (D) Hemotoxylin and eosin stain of the infected skin slices of day 5 and day 10. B: blood vessel; H: hair folliculus.




Figure 7. CLSM images of HeLa cells stained with PI upon incubation with 10 μ M 4TPA-BQ for 12 h under white light irradiation. Irradiation time: 30 min.

So far, the study appears finished, still we want to go deeper into the interaction between the positively charged PS and the negatively charged cells, and to see whether the interaction can be manipulated by regulating external conditions. Given the aforementioned cell staining time is only 15 min, we would like to see whether prolong staining time will make any difference (Figure S22). When COS-7 normal cells treated with 4TPA-BQ for

RESEARCH ARTICLE

4 h, almost no fluorescence signals could be detected, similar with CLSM images of 15 min. When extending incubation time to 8 h, weak orange fluorescence was observed. And more 4TPA-BQ accumulated into COS-7 cells after 12 h incubation. Thus, 4TPA-BQ can accumulate into normal cells under the sufficient long incubation time, however, the fluorescence intensity is too weak for effective visualization. Inspired by the previous report that cancer cells are more negatively surface charged than normal cells, HeLa cancer cells were utilized for cellular imaging under the same conditions.^[25] Intensified fluorescence of 4TPA-BQ was observed along with incubation time as well. Notably, in comparison to COS-7 cells, fluorescence of HeLa cells is remarkably stronger and clearer, hinting that the positively charged PS target cancer cells preferentially over normal cells. Taking advantage of its preference to cancer cells and efficient $^1\text{O}_2$ generation, both normal and cancer cells were employed for PDT application. As shown in Figures S23 and 7, the CLSM images of 4TPA-BQ are consistent with those of cellular imaging. For PI channel, no fluorescence was detectable at 4 h for both COS-7 and HeLa cells; After 8 h incubation, faint red fluorescence from HeLa cells could be noticed whereas COS-7 cells remained no red fluorescence; when incubation time further extended to 12 h, nearly all HeLa cancer cells in sight were stained with PI with enhanced red fluorescence, in sharp contrast to COS-7 cells, in agreement with MTT results (Figure S24). Another cancer cell, A549 cell was also utilized, presenting similar results (Figure S25). Hence, by regulating incubation time to a sufficient long period, 4TPA-BQ can eliminate cancer cells with no apparent toxicity towards normal cells through PDT process.

As demonstrated by above results, a time-dependent PDT towards varying targets was obtained by manipulating incubation time of PS. Virtually, cooperation of electrostatic attraction and hydrophobic effect plays a decisive role in this process.^[26] On the one hand, electrostatic interaction serves as the driving force for possible recognition. The positively charged 4TPA-BQ can target bacteria and cells in sequence, due to the fact that the absolute value of negative surface charges is ranked as bacteria>cancer cells>normal cells.^[20, 25] That is, the more negative the target is charged, the easier it can be bound to the positively charged PS. On the other, in the respect of differentiating bacteria and cells, hydrophobic 4TPA-BQ with ClogP=14 tends to target bacteria preferentially than cells.^[27] Consequently, cationic charge and hydrophobicity is well balanced in 4TPA-BQ. The positive charge endows it with modest water solubility desired for bio application, and more significantly, it is the prerequisite for recognizing varying targets including microbes and cells. Additionally, electrostatic interaction and hydrophobicity makes the multi-targeting in sequence. Essentially, it is the reasonable molecular structure design that realizes the multiple as well as ordered targeting therapeutic effect, which provide guidelines for the future development of antibacterial and anticancer materials.

Conclusion

In summary, a simple AIE-based phototherapeutic agent, namely TPA-BQ, with anion- π^+ interactions was designed and achieved successfully via one step synthesis. The inherent anion- π^+

interactions as well as the highly twisted conformation are responsible for its AIE feature, verified by single crystal analysis. An extremely high $^1\text{O}_2$ quantum yield of 97.8% was obtained in aggregate form thanks to the aggregation-induced ROS generation (AIROSG) effect, overwhelmingly surpassing commercial PSs. Theoretical calculations suggested that sufficient small ΔE_{ST} plays a dominate role in the ISC process, desired for $^1\text{O}_2$ generation. Prominently, 4TPA-BQ can bind to different targets by controlling incubation time. A broad-spectrum and potent antibacterial activity was attained after incubating 4TPA-BQ with pathogens for 15 min. Upon light irradiation, it targeted and killed both Gram-negative and Gram-positive bacteria at a low concentration of 2 μM , leaving normal cells unaffected. Both *in vitro* and *in vivo* experiments demonstrated its outstanding bacterial eliminating performance against ampicillin-resistant *E. coli* infection. When incubated with cells for 12 h, 4TPA-BQ specifically killed cancer cells with low toxicity to normal cells via PDT process. We proposed that this time-dependent PDT was achieved by collaboration of electrostatic attraction and hydrophobic effect between 4TPA-BQ and the corresponding targets. To the best of our knowledge, it is the first reported target-controllable PDT achieved by adjusting external conditions instead of changing the PS structure. Future work will focus on shifting the PS emission to near-infrared region. Therefore, the readily accessible strategy not only provides a platform for next generation PS, but also offers new insights for clinical applications of synergizing with conventional therapeutic modalities.

Experimental Section

Materials and methods, synthetic procedures, characterization, crystallographic data and computational details are available in the supporting information.

Acknowledgements

This work was partially supported by the National Natural Science Foundation of China Grant (21871060, 81271476, and 31870991), the University Grants Committee of Hong Kong (AoE/P-03/08), the Innovation and Technology Commission (ITC-CNERC14SC01 and ITCRD/17-9), the Research Grants Council of Hong Kong (16308016, C6009-17G and A-HKUST605/16), and the Science and Technology Plan of Shenzhen (JCYJ20160229205601482 and JCYJ20170818113348852).

Keywords: aggregation-induced emission • anion- π^+ interactions • time-dependent photodynamic therapy • multiple and ordered targeting

- [1] S. Nollins, J. Cayron, A. Dedieu, A. Page, F. Delolme, C. Lesterlin, *Science* **2019**, 364, 778-782.
- [2] R. L. Siegel, K. D. Miller, A. Jemal, *CA A Cancer J Clin.* **2019**, 69, 7-34.
- [3] a) C. W. Ng, J. Li, K. Pu, *Adv. Funct. Mater.* **2018**, 28, 1804688; b) X. Zhen, J. Zhang, J. Huang, C. Xie, Q. Miao, K. Pu, *Angew. Chem. Int. Ed.* **2018**, 57, 7804-7808.
- [4] a) M. H. Al-Afyouni, T. N. Rohrabough, K. F. Al-Afyouni, C. Turro, *Chem. Sci.* **2018**, 9, 6711-6720; b) Y. Gao, X. Wang, X. He, Z. He, X. Yang, S.


RESEARCH ARTICLE

- Tian, F. Meng, D. Ding, L. Luo, B. Z. Tang, *Adv. Funct. Mater.* **2019**, *29*, 1902673; c) H. Shi, X. Ma, Q. Zhao, B. Liu, Q. Qu, Z. An, Y. Zhao, W. Huang, *Adv. Funct. Mater.* **2014**, *24*, 4823-4830; d) D. Tao, L. Feng, Y. Chao, C. Liang, X. Song, H. Wang, K. Yang, Z. Liu, *Adv. Funct. Mater.* **2018**, *28*, 1804901; e) H. Yuan, H. Chong, B. Wang, C. Zhu, L. Liu, Q. Yang, F. Lv, S. Wang, *J. Am. Chem. Soc.* **2012**, *134*, 13184-13187.
- [5] T. J. Dougherty, C. J. Gomer, B. W. Henderson, G. Jori, D. Kessel, M. Korblick, J. Moan, Q. Peng, *J. Natl. Cancer Inst.* **1998**, *90*, 889-905.
- [6] a) A. S. Klymchenko, *Acc. Chem. Res.* **2017**, *50*, 366-375; b) J. F. Lovell, T. W. B. Liu, J. Chen, G. Zheng, *Chem. Rev.* **2010**, *110*, 2839-2857.
- [7] a) X. He, L.-H. Xiong, Z. Zhao, Z. Wang, L. Luo, J. W. Y. Lam, R. T. K. Kwok, B. Z. Tang, *Theranostics* **2019**, *9*, 3223-3248; b) F. Hu, S. Xu, B. Liu, *Adv. Mater.* **2018**, *30*, 1801350; c) C. Zhu, R. T. K. Kwok, J. W. Y. Lam, B. Z. Tang, *ACS Appl. Bio Mater.* **2018**, *1*, 1768-1786; d) J. Luo, Z. Xie, J. W. Y. Lam, L. Cheng, H. Chen, C. Qiu, H. S. Kwok, X. Zhan, Y. Liu, D. Zhu, B. Z. Tang, *Chem. Commun.* **2001**, 1740-1741; e) N. Alifu, X. Dong, D. Li, X. Sun, A. Zebibula, D. Zhang, G. Zhang, J. Qian, *Mater. Chem. Front.* **2017**, *1*, 1746-1753; f) Y. Huang, G. Zhang, F. Hu, Y. Jin, R. Zhao, D. Zhang, *Chem. Sci.* **2016**, *7*, 7013-7019; g) M. Wang, G. Zhang, D. Zhang, D. Zhu, B. Z. Tang, *J. Mater. Chem.* **2010**, *20*, 1858-1867; h) Y. Chen, W. Ai, X. Guo, Y. Li, Y. Ma, L. Chen, H. Zhang, T. Wang, X. Zhang, Z. Wang, *Small* **2019**, *15*, 1902352; i) Y. Li, Z. Liu, Y. Ma, Y. Chen, K. Ma, X. Wang, D. Zhang, Z. Wang, *Adv. Funct. Mater.* **2019**, *0*, 1903733; j) W. Yu, B. Guo, H. Zhang, J. Zhou, X. Yu, L. Zhu, D. Xue, W. Liu, X. Sun, J. Qian, *Sci. Bull.* **2019**, *64*, 410-416.
- [8] a) H. Gao, X. Zhang, C. Chen, K. Li, D. Ding, *Adv. Biosys.* **2018**, *2*, 1800074; b) J. Mei, N. L. Leung, R. T. Kwok, J. W. Lam, B. Z. Tang, *Chem Rev* **2015**, *115*, 11718-11940.
- [9] a) G. Feng, C. J. Zhang, X. Lu, B. Liu, *ACS Omega* **2017**, *2*, 546-553; b) M. Gao, F. Yu, C. Lv, J. Choo, L. Chen, *Chem. Soc. Rev.* **2017**, *46*, 2237-2271; c) W.-C. Huang, P.-J. Tsai, Y.-C. Chen, *Nanomedicine* **2007**, *2*, 777-787; d) X. Li, J. Kim, J. Yoon, X. Chen, *Adv. Mater.* **2017**, *29*, 1606857; e) X. Li, S. M. Robinson, A. Gupta, K. Saha, Z. Jiang, D. F. Moyano, A. Sahar, M. A. Riley, V. M. Rotello, *ACS Nano* **2014**, *8*, 10682-10686.
- [10] a) J. Wang, X. Gu, H. Ma, Q. Peng, X. Huang, X. Zheng, S. H. P. Sung, G. Shan, J. W. Y. Lam, Z. Shuai, B. Z. Tang, *Nat. Commun.* **2018**, *9*, 2963; b) J. Wang, X. Gu, P. Zhang, X. Huang, X. Zheng, M. Chen, H. Feng, R. T. K. Kwok, J. W. Y. Lam, B. Z. Tang, *J. Am. Chem. Soc.* **2017**, *139*, 16974-16979.
- [11] Z.-W. Chen, Y.-Z. Zhu, J.-W. Ou, Y.-P. Wang, J.-Y. Zheng, *J. Org. Chem.* **2014**, *79*, 10988-10998.
- [12] Y. R. Han, S.-H. Shim, D.-S. Kim, C.-H. Jun, *Org. Lett.* **2017**, *19*, 2941-2944.
- [13] S. Xu, Y. Yuan, X. Cai, C.-J. Zhang, F. Hu, J. Liang, G. Zhang, D. Zhang, B. Liu, *Chem. Sci.* **2015**, *6*, 5824-5830.
- [14] G. R. Martinez, F. Garcia, L. H. Catalani, J. Cadet, M. C. B. Oliveira, G. E. Ronsein, S. Miyamoto, M. H. G. Medeiros, P. D. Mascio, *Tetrahedron* **2006**, *62*, 10762-10770.
- [15] L. Yang, X. Wang, G. Zhang, X. Chen, G. Zhang, J. Jiang, *Nanoscale* **2016**, *8*, 17422-17426.
- [16] a) Z. Meng, B. Yu, G. Han, M. Liu, B. Shan, G. Dong, Z. Miao, N. Jia, Z. Tan, B. Li, W. Zhang, H. Zhu, C. Sheng, J. Yao, *J. Med. Chem.* **2016**, *59*, 4999-5010; b) J. Paczkowski, J. J. M. Lamberts, B. Paczkowska, D. C. Neckers, *Free Radic. Biol. Med.* **1985**, *1*, 341-351.
- [17] D. Wang, H. Su, R. T. K. Kwok, X. Hu, H. Zou, Q. Luo, M. S. Lee, W. Xu, J. W. Y. Lam, B. Z. Tang, *Chem. Sci.* **2018**, *9*, 3685-3693.
- [18] a) W. Wu, D. Mao, F. Hu, S. Xu, C. Chen, C.-J. Zhang, X. Cheng, Y. Yuan, D. Ding, D. Kong, B. Liu, *Adv. Mater.* **2017**, *29*, 1700548; b) S. Xu, W. Wu, X. Cai, C.-J. Zhang, Y. Yuan, J. Liang, G. Feng, P. Manghnani, B. Liu, *Chem. Commun.* **2017**, *53*, 8727-8730.
- [19] a) A. Endo, K. Sato, K. Yoshimura, T. Kai, A. Kawada, H. Miyazaki, C. Adachi, *Appl. Phys. Lett.* **2011**, *98*, 083302; b) H. Uoyama, K. Goushi, K. Shizu, H. Nomura, C. Adachi, *Nature* **2012**, *492*, 234-238; c) K. Schmidt, S. Brovelli, V. Coropceanu, D. Beljonne, J. Cornil, C. Bazzini, T. Caronna, R. Tubino, F. Meinardi, Z. Shuai, J.-L. Brédas, *J. Phys. Chem. A* **2007**, *111*, 10490-10499.
- [20] C. Zhu, Q. Yang, L. Liu, F. Lv, S. Li, G. Yang, S. Wang, *Adv. Mater.* **2011**, *23*, 4805-4810.
- [21] T. Maisch, R.-M. Szeimies, G. Jori, C. Abels, *Photochem. Photobiol. Sci.* **2004**, *3*, 907-917.
- [22] N. Bunyapraphatsara, S. Jirakulchaiwong, S. Thirawarapan, J. Manonukul, *Phytomedicine* **1996**, *2*, 247-251.
- [23] M. Zhao, M. Yang, E. Baranov, X. Wang, S. Penman, A. R. Moossa, R. M. Hoffman, *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 9814-9818.
- [24] Y. Li, S. Wang, R. Huang, Z. Huang, B. Hu, W. Zheng, G. Yang, X. Jiang, *Biomacromolecules* **2015**, *16*, 780-789.
- [25] B. Chen, W. Le, Y. Wang, Z. Li, D. Wang, L. Ren, L. Lin, S. Cui, J. J. Hu, Y. Hu, P. Yang, R. C. Ewing, D. Shi, Z. Cui, *Theranostics* **2016**, *6*, 1887-1898.
- [26] a) G.-j. Liu, S.-n. Tian, C.-y. Li, G.-w. Xing, L. Zhou, *ACS Appl. Mater. Interfaces* **2017**, *9*, 28331-28338; b) E. F. Palermo, K. Lienkamp, E. R. Gillies, P. J. Ragogna, *Angew. Chem. Int. Ed.* **2019**, *58*, 3690-3693.
- [27] a) R. W. Horobin, F. Rashid-Doubell, J. D. Pediani, G. Milligan, *Biotech Histochem* **2013**, *88*, 440-460; b) R. L. Phillips, O. R. Miranda, C.-C. You, V. M. Rotello, U. H. F. Bunz, *Angew. Chem. Int. Ed.* **2008**, *47*, 2590-2594.

RESEARCH ARTICLE

RESEARCH ARTICLE

A simple and high efficient AIE-based photosensitizer, namely 4TPA-BQ, was achieved. A broad-spectrum and potent antibacterial activity was attained after incubating 4TPA-BQ with pathogens for 15 min. When the incubation time was lengthened to 12 h, 4TPA-BQ specifically killed cancer cells with low toxicity to normal cells via PDT process.



Qiyao Li, Ying Li, Tianliang Min, Junyi Gong, Lili Du, David Lee Phillips, Junkai Liu, Jacky W. Y. Lam, Herman H. Y. Sung, Ian D. Williams, Ryan T. K. Kwok, Chun Loong Ho, Kai Li, Jianguo Wang* and Ben Zhong Tang*

Page No. – Page No.

Time-dependent Photodynamic Therapy for Multiple Targets: A Highly Efficient AIE-active Photosensitizer for Selective Bacterial Elimination and Cancer