

# The First Metal Complexes Derived from 3,5-Diethynylpyridine. X-ray Crystal Structure of [(AuPTo<sub>3</sub>)<sub>2</sub>{μ-(C≡C)<sub>2</sub>Py}] (Py = Pyridine-3,5-diyl; To = *p*-Tolyl)

José Vicente,<sup>\*,†</sup> María-Teresa Chicote,<sup>\*,†</sup> and Miguel M. Álvarez-Falcón

Grupo de Química Organometálica, Departamento de Química Inorgánica, Universidad de Murcia, Apartado 4021, Murcia 30071, Spain

Delia Bautista<sup>‡</sup>

SACE, Universidad de Murcia, Apartado 4021, Murcia 30071, Spain

Received July 2, 2004

The reactions of 3,5-diethynylpyridine (Py(C≡CH)<sub>2</sub>) with PPN[Au(acac)<sub>2</sub>] (2.5:1; PPN = Ph<sub>3</sub>P=N=PPh<sub>3</sub>) or with [AuCl(SMe<sub>2</sub>)] and NEt<sub>3</sub> (1:2:2) give respectively PPN[Au{C≡C(Py)-C≡CH}<sub>2</sub>] (**1**) and [Au<sub>2</sub>{μ-(C≡C)<sub>2</sub>Py}]<sub>n</sub> (**2**). Complex **2** reacts with monodentate ligands (1:2) or with 1,6-bis(diphenylphosphino)hexane (dpph, 1:1) to give neutral dinuclear complexes of the general formula [(AuL)<sub>2</sub>{μ-(C≡C)<sub>2</sub>Py}] (L = CN<sup>t</sup>Bu (**3**), PMe<sub>3</sub> (**4**), PPh<sub>3</sub> (**5**), PTo<sub>3</sub> (To = C<sub>6</sub>H<sub>4</sub>Me-4) (**6**); Au<sub>2</sub>L<sub>2</sub> = Au<sub>2</sub>(μ-dpph) (**7**)). The reactions of **6** with the complexes [MCl] and TlTfO (1:1:1) (TfO = CF<sub>3</sub>SO<sub>3</sub>) give the cationic trinuclear complexes [M{Py(C≡CAuPTo<sub>3</sub>)<sub>2</sub>}]TfO (M = AuPTo<sub>3</sub> (**8**), *cis*-PtCl(PPh<sub>3</sub>)<sub>2</sub> (**9**)). The crystal structure of complex **6** has been determined.

## Introduction

Metal complexes with bridging –C≡C(Ar)C≡C– spacers (Ar being an aromatic ring) have been shown to possess electrical conductivity, nonlinear optical, or liquid crystalline properties derived from the extended electronic conjugation they display.<sup>1</sup> The (arene)diethynylgold(I) complexes reported so far include the Ar groups C<sub>6</sub>H<sub>4</sub>-1,3,<sup>2</sup> C<sub>6</sub>H<sub>4</sub>-1,4,<sup>1</sup> (C<sub>6</sub>H<sub>3</sub>Me-5)-1,3, (C<sub>6</sub>H<sub>2</sub>Me<sub>2</sub>-2,5)-1,4,<sup>3</sup> (C<sub>6</sub>HMe<sub>3</sub>-2,4,6)-1,3, C<sub>6</sub>Me<sub>4</sub>-2,3,5,6,<sup>2</sup> and different fluorene-2,7-diyl<sup>4</sup> derivatives, many of which are luminescent. Among them only a few, described by Puddephatt<sup>5</sup> and by us,<sup>2,6</sup> are *m*-arene)diethynyl derivatives; four of these derivatives have been characterized by X-ray diffraction methods, namely [(AuL)<sub>2</sub>{(C≡C)<sub>2</sub>Ar-1,3}] (Ar = C<sub>6</sub>H<sub>4</sub>, C<sub>6</sub>HMe<sub>3</sub>-2,4,6, L = PPh<sub>3</sub>; Ar = C<sub>6</sub>H<sub>3</sub>Me-5, L = PPh<sub>2</sub>Me, P(OMe)<sub>3</sub>). Apart from these, the crystal structures of only three other arene)diethynyl complexes of any element (Ru,<sup>7</sup> Pt,<sup>8</sup> and Fe<sup>9</sup>) have been reported.

3,5-Diethynylpyridine has been used for the synthesis of some fully conjugated ligands containing pyridine-

diyndiyl subunits (Chart 1), from which some Re,<sup>10</sup> Ru,<sup>11</sup> and Pt<sup>12–14</sup> complexes have been obtained and shown to possess metallamacrocyclic structures and interesting physical properties. However, although it was first reported in 1978,<sup>15</sup> there is no previous report on any complex derived from 3,5-diethynylpyridine, which is rather surprising in view of the tempting possibilities this ligand offers as a potential building block for multimetallic complexes, including metallamacrocycles and polymers. Two platinum(II) complexes derived from the monoalkyne HC≡C(Py)C≡CSi<sup>t</sup>Pr<sub>3</sub> have been reported.<sup>14</sup> On the basis of the related 2,5- or 2,6-diethynylpyridine a series of monomeric and polymeric organic compounds<sup>16,17</sup> as well as some σ-acetylide complexes of Ru, Os,<sup>18</sup> and Pt<sup>17,19,20</sup> and some π-complexes

(7) Whittall, I. R.; Humphrey, M. G.; Houbrechts, S.; Maes, J.; Persoons, A.; Schmid, S.; Hockless, D. C. R. *J. Organomet. Chem.* **1997**, *544*, 277.

(8) Onitsuka, K.; Fujimoto, M.; Ohshiro, N.; Takahashi, S. *Angew. Chem., Int. Ed.* **1999**, *38*, 689.

(9) Weyland, T.; Costuas, K.; Toupet, L.; Halet, J. F.; Lapinte, C. *Organometallics* **2000**, *19*, 4228.

(10) Sun, S. S.; Lees, A. J. *Organometallics* **2001**, *20*, 2353.

(11) Campbell, K.; McDonald, R.; Tykwinski, R. R. *J. Org. Chem.* **2002**, *67*, 1133.

(12) Campbell, K.; Kuehl, J.; Ferguson, M. J.; Stang, P. J.; Tykwinski, R. R. *J. Am. Chem. Soc.* **2002**, *124*, 7266. Campbell, K.; McDonald, R.; Ferguson, M. J.; Tykwinski, R. R. *Organometallics* **2003**, *22*, 1353.

(13) Bosch, E.; Barnes, C. L. *Organometallics* **2000**, *19*, 5522.

(14) Campbell, K.; McDonald, R.; Ferguson, M. J.; Tykwinski, R. R. *J. Organomet. Chem.* **2003**, *683*, 379.

(15) Shvartsberg, M. S.; Moroz, A. A.; Kozhevnikova, A. N. *Izv. Akad. Nauk SSSR, Ser. Khim.* **1978**, *4*, 875.

(16) Naka, K.; Uemura, T.; Chujo, Y. *Macromolecules* **2000**, *33*, 4733.

(17) Bunten, K. A.; Kakkar, A. K. *Macromolecules* **1996**, *29*, 2885.

(18) Colbert, M. C. B.; Lewis, J.; Long, N. J.; Raithby, P. R.; Younus, M.; White, A. J. P.; Williams, D. J.; Payne, N. N.; Yellowlees, L.; Beljonne, D.; Chawdhury, N.; Friend, R. H. *Organometallics* **1998**, *17*, 3034.

<sup>†</sup> E-mail: jvs1@um.es (J.V.), mch@um.es (M.-T.C.). WWW: <http://www.um.es/>.

<sup>‡</sup> E-mail: dbc@um.es.

(1) Jia, G. C.; Puddephatt, R. J.; Scott, J. D.; Vittal, J. J. *Organometallics* **1993**, *12*, 3565.


(2) Vicente, J.; Chicote, M. T.; Álvarez-Falcón, M. M.; Abrisqueta, M. D.; Hernández, F. J.; Jones, P. G. *Inorg. Chim. Acta* **2003**, *347*, 67.

(3) Irwin, M. J.; Vittal, J. J.; Puddephatt, R. J. *Organometallics* **1997**, *16*, 6, 3541.

(4) Wong, W. Y.; Choi, K. H.; Lu, G. L.; Shi, J. X.; Lai, P. Y.; Chan, S. M.; Lin, Z. Y. *Organometallics* **2001**, *20*, 5446.

(5) MacDonald, M. A.; Puddephatt, R. J.; Yap, G. P. A. *Organometallics* **2000**, *19*, 2194.

(6) Vicente, J.; Chicote, M. T.; Abrisqueta, M. D.; Álvarez-Falcón, M. M. *J. Organomet. Chem.* **2002**, *663*, 40.

Chart 1<sup>a</sup>

<sup>a</sup> Legend: (A) X = C≡C, Y = Ph<sub>2</sub>C=C, -Z- = -[M]-; (B) X = C≡C, Y = C≡CC≡C, Y = C<sub>6</sub>H<sub>4</sub>-1,3 and X = C≡CC≡C, Y = R<sub>2</sub>C=C (R = Me, Ph).

of Co<sup>21</sup> have been described, which exhibit interesting optical and conductivity properties that in some cases are affected by quaternization of the pyridine nitrogen.<sup>17</sup> The crystal structure of a hexanuclear metallamacrocyclic complex of platinum containing C<sub>5</sub>H<sub>3</sub>N(C≡C)<sub>2</sub>-2,6 ligands<sup>20</sup> represents the only example of a diethynylpyridine complex to be fully characterized.

We have given different accounts on the “acac method”<sup>22</sup> which, on the basis of the ability of (acetylacetonato)gold(I) complexes of the types [Au(acac)<sub>2</sub>]<sup>-</sup> and [Au(acac)L] to deprotonate substrates containing at least an acidic hydrogen atom, led to a variety of complexes, including alkynylgold(I) derivatives.<sup>2,6,23–25</sup> In this paper we report on the use of 3,5-diethynylpyridine to prepare a family of complexes of the types [Au{C≡C(Py)C≡CH}<sub>2</sub>]<sup>-</sup>, [Au<sub>2</sub>{μ-(C≡C)<sub>2</sub>Py}]<sub>n</sub>, [(AuL)<sub>2</sub>{μ-(C≡C)<sub>2</sub>Py}], and [M{Py(C≡CAuPTO<sub>3</sub>)<sub>2</sub>}]<sup>+</sup> (M = AuPTO<sub>3</sub>, *cis*-PtCl(PPh<sub>3</sub>)<sub>2</sub>), which display all the potential coordination modes of the versatile 3,5-diethynylpyridine ligand.

## Experimental Section

IR spectroscopy, mass spectrometry (FAB<sup>+/−</sup>), elemental analyses, conductance measurements in acetone, and melting point determinations were carried out as described elsewhere.<sup>26</sup> The molar conductivity of the neutral complexes gave very low values (0–5 Ω<sup>−1</sup> cm<sup>2</sup> mol<sup>−1</sup>). The NMR spectra were measured on Bruker Avance 200, 300, and 400 MHz spectrometers. Chemical shifts are referred to TMS (<sup>1</sup>H, <sup>13</sup>C) or H<sub>3</sub>PO<sub>4</sub> (<sup>31</sup>P). Unless otherwise stated, the reactions were carried out at room temperature without any precautions to avoid oxygen or moisture. The syntheses of 3,5-diethynylpyridine,<sup>13</sup> PPN[Au(acac)]<sub>2</sub>,<sup>27</sup> and [AuCl(SMe<sub>2</sub>)]<sub>2</sub><sup>28</sup> were previously reported. [AuCl-

(PTO<sub>3</sub>)] and *cis*-[PtCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>] were prepared by adding 1 or 2 equiv of the appropriate phosphine to [AuCl(SMe<sub>2</sub>)] and [PtCl<sub>2</sub>(NPh)<sub>2</sub>],<sup>29</sup> respectively. TfO (TfO = CF<sub>3</sub>SO<sub>3</sub>) was obtained by reacting in water Tl<sub>2</sub>CO<sub>3</sub> and HtFO (1:2) after evaporation of the solvent and recrystallization of the residue from acetone and Et<sub>2</sub>O. All other chemicals were obtained from commercial sources and used as received. The solvents were distilled before use.

**Synthesis of PPN[Au{C≡C(Py)C≡CH}<sub>2</sub>] (1).** A solution of PPN[Au(acac)<sub>2</sub>] (676 mg, 0.72 mmol) in acetone (15 mL) was slowly dropped over a solution of 3,5-diethynylpyridine (23 mg, 1.8 mmol) in the same solvent (10 mL). The mixture was stirred for 8 h and filtered through Celite and the solution concentrated to dryness. The residue was stirred with Et<sub>2</sub>O/CH<sub>2</sub>Cl<sub>2</sub> (14:1, 3 × 45 mL), and the colorless microcrystalline solid was filtered and dried under reduced pressure (ca. 1 mbar) for 1 h to give 1·H<sub>2</sub>O·Et<sub>2</sub>O. Yield: 592 mg, 76%. Mp: 133 °C. Anal. Calcd for C<sub>58</sub>H<sub>50</sub>AuN<sub>3</sub>O<sub>2</sub>P<sub>2</sub>: C, 64.50; H, 4.67; N, 3.89. Found: C, 64.49; H, 4.28; N, 3.93. IR (cm<sup>−1</sup>): ν(CH) 3294 (w), 3232 (w); ν(C≡C) 2106 (s). Λ<sub>M</sub> (Ω<sup>−1</sup> cm<sup>2</sup> mol<sup>−1</sup>): 81. <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>): δ 8.51 (m, 2 H, py), 8.35 (m, 2 H, py), 8.23 (m, 2 H, py), 7.69–7.28 (m, 30 H, PPN), 3.5 (q, 4 H, CH<sub>2</sub>, Et<sub>2</sub>O, <sup>3</sup>J<sub>HH</sub> = 7 Hz), 3.11 (s, 2 H, CH), 1.66 (s, 2 H, H<sub>2</sub>O), 1.25 (t, 6 H, Me, Et<sub>2</sub>O, <sup>3</sup>J<sub>HH</sub> = 7 Hz). <sup>31</sup>P{<sup>1</sup>H} NMR (81 MHz, CDCl<sub>3</sub>): δ 21.1 (s, PPN). <sup>13</sup>C{<sup>1</sup>H} NMR (50 MHz, CDCl<sub>3</sub>): δ 152.6, 148.1, 141.5, 140.0, 133.8 (m, *p*-CH, PPN), 132.0 (m, *m*-CH, PPN), 129.6 (m, *o*-CH, PPN), 126.8 (m, *i*-C, PPN), 123.4, 117.9, 98.0 (C≡C), 80.6 (C≡CH), 79.7 (≡CH), 65.91 (CH<sub>2</sub>, Et<sub>2</sub>O), 15.20 (Me, Et<sub>2</sub>O). MS-FAB<sup>−</sup> (*m/z*, %): 449 (M<sup>−</sup>, 100%).

**Synthesis of [Au<sub>2</sub>{(C≡C)<sub>2</sub>Py}]<sub>n</sub> (2).** A mixture of 3,5-diethynylpyridine (198 mg, 1.6 mmol), [AuCl(SMe<sub>2</sub>)] (917 mg, 3.11 mmol), and NEt<sub>3</sub> (0.46 mL, 3.26 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was stirred for 0.5 h. The suspension was filtered and the solid washed successively with CH<sub>2</sub>Cl<sub>2</sub> (4 × 10 mL) and Et<sub>2</sub>O (10 mL) and dried under vacuum (ca. 1 mbar) for 1 h to give 2 as a bright yellow powder. Its insolubility in all common solvents impeded measuring its NMR spectra or recrystallizing it to improve the elemental analysis (see Results and Discussion). Yield: 802 mg, 97%. Mp: 115 °C dec. Anal. Calcd for C<sub>9</sub>H<sub>3</sub>Au<sub>2</sub>N: C, 20.83; H, 0.58; N, 2.70. Found: C, 22.15; H, 0.97; N, 3.06. IR (cm<sup>−1</sup>): ν(C≡C) 2114 (w).

**Synthesis of [(AuL)<sub>2</sub>{μ-(C≡C)<sub>2</sub>Py}] (L = <sup>t</sup>BuNC (3), PMe<sub>3</sub> (4), PPh<sub>3</sub> (5), PTO<sub>3</sub> (6); Au<sub>2</sub>L<sub>2</sub> = Au<sub>2</sub>(μ-dpph) (7)).** To a suspension of 2 (3, 286 mg, 0.55 mmol; 4, 213 mg, 0.41 mmol; 5, 120 mg, 0.23 mmol; 6, 277 mg, 0.53 mmol; 7, 243 mg, 0.47 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3, 10 mL; 4 and 5, 20 mL; 6, 30 mL; 7, 15 mL) was added the appropriate ligand (3, <sup>t</sup>BuNC 0.25 mL, 2.2 mmol; 4, PMe<sub>3</sub> 1 M in toluene, 2 mL, 2 mmol; 5, PPh<sub>3</sub> 127 mg, 0.49 mmol; 6, PTO<sub>3</sub> 325 mg, 1.07 mmol; 7, dpph 213 mg, 0.47 mmol). The reaction mixture was stirred for 1 h at room temperature (4, under an N<sub>2</sub> atmosphere) or refluxed for 4 h (7) and filtered through Celite. The solvent was removed under vacuum to 1 mL (4) or 3 mL (7) or to dryness (3, 5, 6). When *n*-hexane (3, 4, 10 mL) or Et<sub>2</sub>O (5, 7, 20 mL) or a 1:1 mixture of both solvents (6, 20 mL) was added and the mixture was stirred for a few minutes, a suspension resulted, which was filtered. The solid was washed with the same solvent (2 × 5 mL) and air-dried to give a bright yellow (3·0.5Et<sub>2</sub>O) or pale yellow (4, 5·H<sub>2</sub>O, 6) or white (7·H<sub>2</sub>O) powder.

3·0.5Et<sub>2</sub>O: yield 276 mg, 73%. Mp: 140 °C. Anal. Calcd for C<sub>21</sub>H<sub>26</sub>Au<sub>2</sub>N<sub>3</sub>O<sub>0.5</sub>: C, 34.92; H, 3.63; N, 5.82. Found: C, 34.80; H, 3.48; N, 6.35. IR (cm<sup>−1</sup>): ν(C≡N) 2230 (s); ν(C≡C) 2120 (w). <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>): δ 8.44 (d, 2 H, py, <sup>4</sup>J<sub>HH</sub> = 2 Hz), 7.69 (t, 1 H, py, <sup>4</sup>J<sub>HH</sub> = 2 Hz), 3.1 (q, 2 H, CH<sub>2</sub>, Et<sub>2</sub>O, <sup>3</sup>J<sub>HH</sub> = 7 Hz), 1.57 (s, 18 H, <sup>t</sup>Bu), 1.4 (t, 3 H, Me, Et<sub>2</sub>O, <sup>3</sup>J<sub>HH</sub> = 7 Hz). <sup>13</sup>C{<sup>1</sup>H} NMR (50 MHz, CDCl<sub>3</sub>): δ 150.3, 146.4 (CN), 141.6,

(19) Chawdhury, N.; Köhler, A.; Friend, R. H.; Younus, M.; Long, N. J.; Raithby, P. R.; Lewis, J. *Macromolecules* **1998**, *31*, 722.

(20) Leininger, S.; Schmitz, M.; Stang, P. J. *Org. Lett.* **1999**, *1*, 1921.

(21) Dana, B. H.; Robinson, B. H.; Simpson, J. J. *Organomet. Chem.* **2002**, *648*, 251.

(22) Vicente, J.; Chicote, M. T. *Coord. Chem. Rev.* **1999**, *193–195*, 1143.

(23) Vicente, J.; Chicote, M. T.; Abrisqueta, M. D. *J. Chem. Soc., Dalton Trans.* **1995**, 497. Vicente, J.; Chicote, M. T.; Abrisqueta, M. D.; Jones, P. G. *Organometallics* **1997**, *16*, 5628.

(24) Vicente, J.; Chicote, M.-T.; Abrisqueta, M.-D.; Guerrero, R.; Jones, P. G. *Angew. Chem., Int. Ed.* **1997**, *36*, 1203.

(25) Vicente, J.; Chicote, M. T.; Álvarez-Falcón, M. M.; Fox, M. A.; Bautista, D. *Organometallics* **2003**, *22*, 4792.

(26) Vicente, J.; Chicote, M. T.; Guerrero, R.; Saura-Llamas, I. M.; Jones, P. G.; Ramírez de Arellano, M. C. *Chem. Eur. J.* **2001**, *7*, 638.

(27) Vicente, J.; Chicote, M. T. *Inorg. Synth.* **1998**, *32*, 172.

(28) Tamaki, A.; Kochi, J. K. *J. Organomet. Chem.* **1974**, *64*, 411.

(29) Doyle, J. R.; Slade, P. E.; Jonassen, H. B. *Inorg. Synth.* **1960**, *6*, 128.

126.6, 121.0, 99.2 (C≡C), 59.0 (CH<sub>2</sub>, Et<sub>2</sub>O), 58.6 (CMe<sub>3</sub>), 29.6 (CMe<sub>3</sub>), 8.9 (Me, Et<sub>2</sub>O).

**4:** yield 191 mg, 69%. Mp: 293 °C dec. Anal. Calcd for C<sub>15</sub>H<sub>21</sub>Au<sub>2</sub>NP<sub>2</sub>: C, 26.84; H, 3.15; N, 2.09. Found: C, 26.88; H, 3.13; N, 2.08. IR (cm<sup>-1</sup>): ν(C≡C) 2098 (w). <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>): δ 8.47 (d, 2 H, py, <sup>4</sup>J<sub>HH</sub> = 2 Hz), 7.73 (t, 1 H, py, <sup>4</sup>J<sub>HH</sub> = 2 Hz), 1.55 (d, 18 H, PMe<sub>3</sub>, <sup>2</sup>J<sub>HP</sub> = 10 Hz). <sup>31</sup>P{<sup>1</sup>H} NMR (81 MHz, CDCl<sub>3</sub>): δ 1.36 (s). <sup>13</sup>C{<sup>1</sup>H} NMR (50 MHz, CDCl<sub>3</sub>): δ 150.5 (py), 141.5 (py), 121.2 (d, CCAu, <sup>3</sup>J<sub>CP</sub> = 5 Hz), 100.7 (d, CAu, <sup>2</sup>J<sub>CP</sub> = 30 Hz), 15.7 (d, PMe<sub>3</sub>, <sup>1</sup>J<sub>CP</sub> = 35 Hz).

**5**·H<sub>2</sub>O: yield 161 mg, 67%. Mp: 115 °C. Anal. Calcd for C<sub>45</sub>H<sub>35</sub>NAu<sub>2</sub>OP<sub>2</sub>: C, 50.91; H, 3.32; N, 1.32. Found: C, 50.79; H, 3.39; N, 1.27. IR (cm<sup>-1</sup>): ν(C≡C) 2112 (s). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.55 (d, 2 H, py, <sup>4</sup>J<sub>HH</sub> = 2 Hz), 7.83 (t, 1 H, py, <sup>4</sup>J<sub>HH</sub> = 2 Hz), 7.57–7.45 (m, 30 H, PPh<sub>3</sub>), 1.66 (s, 2 H, H<sub>2</sub>O). <sup>31</sup>P{<sup>1</sup>H} NMR (162 MHz, CDCl<sub>3</sub>): δ 42.61 (s). <sup>13</sup>C{<sup>1</sup>H} NMR (50 MHz, CDCl<sub>3</sub>): δ 150.7 (py), 141.5 (py), 137.5 (py), 134.3 (d, *o*-CH, PPh<sub>3</sub>, <sup>2</sup>J<sub>CP</sub> = 14 Hz), 131.6 (d, *p*-CH, PPh<sub>3</sub>, <sup>4</sup>J<sub>CP</sub> = 3 Hz), 129.6 (d, *i*-C, PPh<sub>3</sub>, <sup>1</sup>J<sub>CP</sub> = 56 Hz), 129.2 (d, *m*-CH, PPh<sub>3</sub>, <sup>3</sup>J<sub>CP</sub> = 11 Hz), 121.2 (d, CCAu, <sup>3</sup>J<sub>CP</sub> = 3 Hz), 100.2 (d, CAu, <sup>2</sup>J<sub>CP</sub> = 27 Hz).

**6:** yield 488 mg, 82%. Mp: 206 °C dec. Anal. Calcd for C<sub>51</sub>H<sub>45</sub>Au<sub>2</sub>NP<sub>2</sub>: C, 54.31; H, 4.02; N, 1.24. Found: C, 54.14; H, 4.30; N, 1.33. IR (cm<sup>-1</sup>): ν(C≡C) 2112 (w). <sup>1</sup>H NMR (200 MHz, CDCl<sub>3</sub>): δ 8.55 (d, 2 H, py, <sup>4</sup>J<sub>HH</sub> = 2 Hz), 7.82 (t, 1 H, py, <sup>4</sup>J<sub>HH</sub> = 2 Hz), 7.47–7.37 (m, 12 H, PT<sub>3</sub>), 7.25–7.21 (m, 12 H, PT<sub>3</sub>), 2.39 (s, 18 H, Me). <sup>31</sup>P{<sup>1</sup>H} NMR (81 MHz, CDCl<sub>3</sub>): δ 41.22 (s). <sup>13</sup>C{<sup>1</sup>H} NMR (50 MHz, CDCl<sub>3</sub>): δ 150.7, 141.9 (d, *p*-CH, PT<sub>3</sub>, <sup>4</sup>J<sub>CP</sub> = 2 Hz), 141.5, 134.1 (d, *o*-CH, PT<sub>3</sub>, <sup>2</sup>J<sub>CP</sub> = 14 Hz), 129.8 (d, *m*-CH, PT<sub>3</sub>, <sup>3</sup>J<sub>CP</sub> = 12 Hz), 126.6 (d, *i*-C, PT<sub>3</sub>, <sup>1</sup>J<sub>CP</sub> = 58 Hz), 121.2, 21.5 (d, Me, <sup>5</sup>J<sub>CP</sub> = 1 Hz).

**7**·H<sub>2</sub>O: yield 226 mg, 49%. Mp: 131 °C dec. Anal. Calcd for C<sub>39</sub>H<sub>37</sub>Au<sub>2</sub>NOP<sub>2</sub>: C, 47.24; H, 3.76; N, 1.41. Found: C, 47.05; H, 3.41; N, 1.52. IR (cm<sup>-1</sup>): ν(C≡C) 2106 (w). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.52 (d, 2 H, py, <sup>4</sup>J<sub>HH</sub> = 2 Hz), 7.79 (m, 1 H, py), 7.68–7.44 (m, 20 H, Ph), 2.37 (m, 4 H, CH<sub>2</sub>), 1.66 (s, 2 H, H<sub>2</sub>O), 1.59 (m, 4 H, CH<sub>2</sub>), 1.43 (m, 4 H, CH<sub>2</sub>). <sup>31</sup>P{<sup>1</sup>H} NMR (162 MHz, CDCl<sub>3</sub>): δ 37.35 (s). <sup>13</sup>C{<sup>1</sup>H} NMR (75 MHz, CDCl<sub>3</sub>): δ 150.5, 141.5, 133.3 (d, *m*-CH, Ph, <sup>3</sup>J<sub>CP</sub> = 8 Hz), 131.6 (m, *p*-CH, Ph), 130.3 (d, *i*-C, Ph, <sup>1</sup>J<sub>CP</sub> = 53 Hz), 129.2 (d, *o*-CH, Ph, <sup>2</sup>J<sub>CP</sub> = 15 Hz), 121.2 (s, CCAu), 100.1 (s, CAu), 30.3 (d, CH<sub>2</sub>, <sup>2</sup>J<sub>CP</sub> = 15 Hz), 27.8 (d, CH<sub>2</sub>, <sup>1</sup>J<sub>CP</sub> = 34 Hz), 25.3 (d, CH<sub>2</sub>, <sup>3</sup>J<sub>CP</sub> = 3 Hz). MS-FAB<sup>+</sup> (*m/z*, %): 974 (M<sup>+</sup>, 6%), 651 (Audp<sup>+</sup>, 36%).

**Synthesis of [(AuPT<sub>3</sub>)<sub>2</sub>{Py(C≡CAuPT<sub>3</sub>)<sub>2</sub>}]TfO (8).** A mixture of **6** (213 mg, 0.19 mmol), [AuClPT<sub>3</sub>] (107 mg, 0.20 mmol), and TfO (71 mg, 0.20 mmol) in dry THF (20 mL) was stirred under a nitrogen atmosphere for 0.5 h and then concentrated to dryness under reduced pressure. The residue was stirred with CHCl<sub>3</sub> (30 mL), and the suspension was filtered through Celite. The solution was concentrated to dryness, the residue stirred with *n*-pentane (2 × 20 mL), the suspension filtered, and the solid air-dried to give **8** as a microcrystalline bright yellow powder. Yield: 242 mg, 72%. Mp: 130 °C. Anal. Calcd for C<sub>73</sub>H<sub>66</sub>Au<sub>3</sub>F<sub>3</sub>NO<sub>3</sub>P<sub>3</sub>S: C, 49.31; H, 3.74; N, 0.79. Found: C, 49.53; H, 3.91; N, 0.66. IR (cm<sup>-1</sup>): ν(C≡C) 2114 (w). Λ<sub>m</sub> (Ω<sup>-1</sup> cm<sup>2</sup> mol<sup>-1</sup>): 105. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.49 (br, 2 H, py), 7.84 (t, 1 H, py, <sup>4</sup>J<sub>HH</sub> = 1.8 Hz), 7.41–7.22 (m, 36 H, Ar, PT<sub>3</sub>), 2.44 (s, 9 H, Me), 2.39 (s, 18 H, Me). <sup>31</sup>P{<sup>1</sup>H} NMR (162 MHz, CDCl<sub>3</sub>): δ 43.68 (s, CAuPT<sub>3</sub>), 36.85 (br s, NAuPT<sub>3</sub>). <sup>13</sup>C{<sup>1</sup>H}-APT NMR (100 MHz, CDCl<sub>3</sub>): δ 150.5, 143.3 (*p*-C, PT<sub>3</sub>), 142.3 (*p*-C, PT<sub>3</sub>), 133.9 (d, *o*-CH, PT<sub>3</sub>, <sup>2</sup>J<sub>CP</sub> = 14 Hz), 133.7 (m, *o*-CH, NAuPT<sub>3</sub>), 130.5 (m, *m*-CH, NAuPT<sub>3</sub>), 129.9 (d, *m*-CH, PT<sub>3</sub>, <sup>3</sup>J<sub>CP</sub> = 12 Hz), 125.7 (d, *i*-C, PT<sub>3</sub>, <sup>1</sup>J<sub>CP</sub> = 61 Hz), 124.2 (m, *i*-C, NAuPT<sub>3</sub>), 121.5, 21.5 (Me, PT<sub>3</sub>), 21.4 (Me, PT<sub>3</sub>). MS-FAB<sup>+</sup> (*m/z*, %): 1628 (M<sup>+</sup>, 27%).

**Synthesis of [(*cis*-PtCl(PPh<sub>3</sub>)<sub>2</sub>){Py(C≡CAuPT<sub>3</sub>)<sub>2</sub>}]TfO (9).** A mixture of **6** (230 mg, 0.2 mmol), *cis*-[PtCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>] (161 mg, 0.2 mmol), and TfO (72 mg, 0.2 mmol) in acetone (25 mL) was stirred for 1.25 h. The solvent was removed under

Table 1. Crystal Data for Complex 6

formula	C <sub>51</sub> H <sub>45</sub> Au <sub>2</sub> NP <sub>2</sub>
fw	1127.75
cryst size (mm)	0.34 × 0.28 × 0.06
cryst syst	monoclinic
space group	<i>P</i> 2 <sub>1</sub> / <i>c</i>
<i>a</i> (Å)	9.9851(6)
<i>b</i> (Å)	23.4212(19)
<i>c</i> (Å)	19.0934(15)
α (deg)	90
β (deg)	91.270(6)
γ (deg)	90
<i>V</i> (Å <sup>3</sup> )	4464.1(6)
<i>Z</i>	4
ρ <sub>calcd</sub> (Mg m <sup>-3</sup> )	1.678
λ (Å)	0.710 73
<i>T</i> (K)	173(2)
<i>F</i> (000)	2184
μ(Mo Kα) (mm <sup>-1</sup> )	6.671
θ range (deg)	3.05–25.00
abs cor	ψ scans
no. of rflns coll	8437
no. of indep rflns	7835
<i>R</i> <sub>int</sub>	0.0407
transmissn	0.997 05/0.449 49
no. of data/restraints/params	7835/18/505
<i>R</i> 1 ( <i>I</i> > 2σ( <i>I</i> )) <sup>a</sup>	0.0403
<i>wR</i> 2 (all rflns) <sup>b</sup>	0.0668

<sup>a</sup> *R*1 = Σ||*F*<sub>o</sub>| - |*F*<sub>c</sub>||/Σ|*F*<sub>o</sub>| for reflections with *I* > 2σ(*I*). <sup>b</sup> *wR*2 = [Σ(*w*(*F*<sub>o</sub><sup>2</sup> - *F*<sub>c</sub><sup>2</sup>)<sup>2</sup>)/Σ(*w*(*F*<sub>c</sub><sup>2</sup>)<sup>2</sup>)]<sup>0.5</sup> for all reflections; *w*<sup>-1</sup> = σ<sup>2</sup>(*F*<sup>2</sup>) + (*aP*)<sup>2</sup> + *bP*, where *P* = (2*F*<sub>c</sub><sup>2</sup> + *F*<sub>o</sub><sup>2</sup>)/3 and *a* and *b* are constants set by the program.

vacuum, and the residue was stirred with CHCl<sub>3</sub> (20 mL). The resulting suspension was filtered through Celite, the solution concentrated to dryness, and the solid residue stirred with *n*-pentane (25 mL). The suspension was filtered and the solid dried under reduced pressure (ca. 1 mbar) for 1 h to give a microcrystalline pale yellow powder. Yield: 365 mg, 90%. Mp: 142 °C dec. Anal. Calcd for C<sub>88</sub>H<sub>75</sub>Au<sub>2</sub>ClF<sub>3</sub>NO<sub>3</sub>P<sub>4</sub>TS: C, 52.02; H, 3.72; N, 0.69. Found: C, 52.08; H, 3.78; N, 0.89. IR (cm<sup>-1</sup>): ν(C≡C) 2114 (s); ν(PtCl), 318(s). Λ<sub>m</sub> (Ω<sup>-1</sup> cm<sup>2</sup> mol<sup>-1</sup>): 146. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.17 (m, 2 H, py), 7.48–7.14 (m, 55 H, PT<sub>3</sub> + PPh<sub>3</sub> + py), 2.42 (s, 18 H, Me). <sup>31</sup>P{<sup>1</sup>H} NMR (121 MHz, CDCl<sub>3</sub>): δ 40.22 (s, AuPT<sub>3</sub>), 16.67 (d, PtPPh<sub>3</sub>, <sup>1</sup>J<sub>PPt</sub> = 3655 Hz, <sup>2</sup>J<sub>PP</sub> = 18 Hz), 4.74 (d, PtPPh<sub>3</sub>, <sup>1</sup>J<sub>PPt</sub> = 3240 Hz, <sup>2</sup>J<sub>PP</sub> = 18 Hz). MS-FAB<sup>+</sup> (*m/z*, %): 1881 (M<sup>+</sup>, 26%).


**X-ray Structure Determination for Complex 6.** The crystals were mounted in inert oil on a glass fiber and transferred to the diffractometer (Siemens P4 with LT2 low-temperature attachment). Crystal data and refinement details are presented in Table 1. The unit cell parameters were determined from a least-squares fit of 88 accurately centered reflections (14.1° < 2θ < 25.8°). The structure was solved by the heavy-atom method and refined anisotropically on *F*<sup>2</sup> (program SHELX-97, G. M. Sheldrick, University of Göttingen, Göttingen, Germany). The largest residual peak of 1.62 e was 1.13 Å from Au2. Hydrogen atoms were included using a riding model.

## Results and Discussion


**Synthesis.** The reaction of 3,5-diethynylpyridine with PPN[Au(acac)<sub>2</sub>] to give the anionic complex PPN[Au{C≡C(Py)C≡CH}<sub>2</sub>] (1), in which gold coordinates to two monoanionic HC≡C(Py)C≡C<sup>-</sup> (Py = pyridine-3,5-diyl) ligands, requires the use of an excess of the dialkyne over the stoichiometric 2:1 molar ratio (Scheme 1) in order to prevent the formation of insoluble oligomers of the type (PPN)<sub>*n*</sub>[Au{(C≡C)<sub>2</sub>Py}]<sub>*n*</sub>. The complex [Au<sub>2</sub>{(C≡C)<sub>2</sub>Py}]<sub>*n*</sub> (2) was obtained by reacting 3,5-diethynylpyridine with [AuCl(SMe<sub>2</sub>)] and NEt<sub>3</sub>







**Figure 1.** Ellipsoid representation of **6** (50% probability). Selected bond lengths (Å) and angles (deg): Au(1)–C(10) = 2.011(10), Au(1)–P(1) = 2.269(2), C(9)–C(10) = 1.188(11), C(10)–Au(1)–P(1) = 175.3(2), Au(2)–C(8) = 1.995(8), Au(2)–P(2) = 2.269(2), C(7)–C(8) = 1.183(11), C(8)–Au(2)–P(2) = 173.3(2).



**Figure 2.** Two different views of the aurophilic contacts in **6** giving rise to infinite chains. The tolyl groups and the hydrogen atoms have been omitted. Selected bond lengths (Å) and angles (deg): Au(1)–Au(2)#1 = 3.2265(5), C(10)–Au(1)–Au(2)#1 = 76.6(2), P(1)–Au(1)–Au(2)#1 = 108.01(6), C(8)–Au(2)–Au(1)#2 = 76.1(2), P(2)–Au(2)–Au(1)#2 = 104.14(6).

**Crystal Structure of [(AuPTO<sub>3</sub>)<sub>2</sub>{μ-(C≡C)<sub>2</sub>Py}] (6).** The crystal structure of **6** (Figure 1, Table 1) shows a dinuclear complex in which two AuPTO<sub>3</sub> units are coordinated to the ethynyl fragments of the 3,5-pyridinediethynyl ligand, both gold atoms being in almost linear environments (C–Au–P = 175.3(2), 173.3(2)°). A search of the Cambridge Structural Data Base<sup>31</sup> reveals that the Au–C (2.011(10), 1.995(8) Å), Au–P (2.269(2) Å), and C≡C (1.183(11), 1.188(11) Å) bond distances found in **6** are similar to the mean values found for other alkynyl(phosphine)gold(I) dicoordinate complexes (Au–C = 2.004 Å, Au–P = 2.278 Å, C≡C = 1.195 Å). Additionally, the molecules of **6** are packed through intermolecular aurophilic contacts (Au...Au = 3.2265(5) Å) to give infinite chains (Figure 2). The molecules arrange in the chain so that the linear fragments C≡CAuPTO<sub>3</sub> of two vicinal molecules are almost perpendicular (dihedral angle PAu...AuP 89°). This type of “pseudopolymerization” in the solid state has been previously reported for other (alkynyl)gold(I) complexes.<sup>5,32–34</sup>

**NMR Spectra.** The heteroaromatic protons appear in the <sup>1</sup>H NMR spectra of complexes **1–9** in the ranges 8.44–8.55 ppm (d) and 7.69–7.83 ppm (t), only very slightly shielded with respect to their homologues in the free 3,5-diethynylpyridine ligand (at 8.65 and 7.85 ppm, respectively). Additionally, the <sup>1</sup>H NMR spectrum of **1** shows a singlet for the acetylenic protons at 3.11 ppm, slightly shifted to high field with respect to the free dialkyne (at 3.24 ppm), likely due to the anionic nature of **1**. The PTO<sub>3</sub> methyl protons in **8** give rise to two singlets. The singlet at 2.39 ppm corresponds to the C–AuPTO<sub>3</sub> fragments and is at higher field than that due to the N–AuPTO<sub>3</sub> moiety (at 2.44 ppm), in agreement with the greater electronegativity of nitrogen with respect to carbon.

The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of complex **1** shows a singlet at 21.1 ppm due to the PPN cation, while the spectra for the neutral complexes **3–7** display one singlet resonance (at 1.36 ppm (**4**, PMe<sub>3</sub>) or in the range 36–44 ppm (**5–7**, PAr<sub>3</sub>)) for the two equivalent phosphorus nuclei. In agreement with the <sup>1</sup>H NMR, the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of **8** shows two singlet resonances with a 2:1 intensity ratio. The most intense one (at 43.68 ppm) corresponding to two C–AuP fragments is shifted to low field with respect to that due to the N–AuP fragment (at 36.85 ppm), despite nitrogen being more electronegative than carbon. This can be attributed to the greater trans influence of carbon-donor with respect to nitrogen-donor ligands. In fact, complexes with the AuPPh<sub>3</sub> fragment coordinated to the nitrogen atom of different substituted pyridines or to alkynyl groups show the <sup>31</sup>P resonance at around 30 ppm<sup>35</sup> or in the range 42–56 ppm,<sup>25,36</sup> respectively. In the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of **9** the phosphine bonded to gold gives a singlet resonance (see Experimental Section), while those coordinated to platinum give two doublets with <sup>195</sup>Pt satellites, corresponding to the isomer containing a *cis*-PtCl(PPh<sub>3</sub>)<sub>2</sub> fragment. The spectrum of **9** shows a small singlet with <sup>195</sup>Pt satellites at 14.86 ppm (*J*<sub>PtP</sub> = 3677 Hz), compatible with the formation of a small amount (ca. 7%) of the isomer containing *trans*-PtCl(PPh<sub>3</sub>)<sub>2</sub>. Additionally, it shows a minor signal at 32.4 ppm (s) that grows with the acquisition time and is indicative of the formation of [AuClPTO<sub>3</sub>] resulting from some decomposition process, which impeded us measuring its <sup>13</sup>C NMR spectrum.

The <sup>13</sup>C{<sup>1</sup>H} NMR spectra of complexes **1** and **3–8** show the expected resonances (see Experimental Sec-

(31) CSD Version November 2003.

(32) Mohr, F.; Eisler, D. J.; McArdle, C. P.; Atieh, K.; Jennings, M. C.; Puddephatt, R. J. *J. Organomet. Chem.* **2003**, 670, 27.

(33) Mohr, F.; Jennings, M. C.; Puddephatt, R. J. *Eur. J. Inorg. Chem.* **2003**, 217. Irwin, M. J.; Jia, G.; Payne, N. C.; Puddephatt, R. J. *Organometallics* **1996**, 15, 51. Puddephatt, R. J. *Chem. Commun.* **1998**, 1055. Irwin, M. J.; Manojlovic-Muir, L.; Muir, K. W.; Puddephatt, R. J.; Yufit, D. S. *Chem. Commun.* **1997**, 219. Puddephatt, R. J. *Coord. Chem. Rev.* **2001**, 216, 313. Liau, R. Y.; Schier, A.; Schmidbaur, H. *Organometallics* **2003**, 22, 3199.

(34) McArdle, C. P.; Irving, M. J.; Jennings, M. C.; Vittal, J. J.; Puddephatt, R. J. *Chem. Eur. J.* **2002**, 8, 723.

(35) Casas, J. E.; Gimeno, M. C.; Jones, P. G.; Laguna, A.; Laguna, M. J. *Chem. Soc., Dalton Trans.* **1999**, 2819. Crespo, O.; Gimeno, M. C.; Laguna, A. *J. Organomet. Chem.* **1998**, 561, 13. Barranco, E. N.; Crespo, O.; Gimeno, M. C.; Jones, P. G.; Laguna, A.; Sarroca, C. *Dalton Trans.* **2001**, 2523.

(36) Lu, W.; Zhu, N. Y.; Che, C. M. *J. Am. Chem. Soc.* **2003**, 125, 16081. Hunks, W. J.; Macdonald, M. A.; Jennings, M. C.; Puddephatt, R. J. *Organometallics* **2000**, 19, 5063. Whittall, I. R.; Humphrey, M. G.; Houbrechts, S.; Persoons, A.; Hockless, D. C. R. *Organometallics* **1996**, 15, 5738.

tion). In those complexes containing  $\text{C}\equiv\text{C}-\text{AuPR}_3$  fragments the  $\alpha$ - and  $\beta$ -carbons of the  $\text{C}\equiv\text{C}$  subunits could be assigned when the  $J_{\text{CP}}$  coupling constants were observed, on the basis of the admitted fact<sup>23,34,37,38</sup> that they are greater for the  $\alpha$ -carbons. In all other cases, no assignment has been made, since controversial data can be found in the literature.<sup>38</sup>

NMR and analytical data prove that some complexes crystallize with  $\text{Et}_2\text{O}$  (**3**·0.5 $\text{Et}_2\text{O}$ ) or  $\text{H}_2\text{O}$  (**5**· $\text{H}_2\text{O}$ , **7**· $\text{H}_2\text{O}$ ).

**IR Spectra.** The infrared spectra of the ionic complexes show characteristic bands of the corresponding counterions (**1**, PPN;<sup>6</sup> **8**, **9**, TfO).<sup>39</sup> All complexes show one medium to weak band in the range 2098–2114  $\text{cm}^{-1}$  assignable to the  $\text{C}\equiv\text{C}$  stretching mode.<sup>40</sup> This band was expected<sup>32,41</sup> to appear at lower wavenumber in the polymeric complex **2** due to the  $(\text{C}\equiv\text{C})\rightarrow\text{Au}$  bond component, and its position should also change with the electron density around the metal center. However, very similar or identical values are found for complexes **1**, **2**, **7**, and **8** (2106, 2114, 2106, and 2114  $\text{cm}^{-1}$ , respectively), despite **1** being anionic, **2** polymeric, **7** neutral, and **8** cationic. In the spectrum of complex **1** two bands are observed (3294, 3232  $\text{cm}^{-1}$ ) for the  $\equiv\text{C}-\text{H}$  stretching

mode,<sup>40</sup> as observed in other alkynylgold complexes with two  $\text{C}\equiv\text{CH}$  moieties.<sup>6</sup> Additionally, complex **9** shows one  $\nu(\text{PtCl})$  band and **3–9** show the bands characteristic of the phosphine or isocyanide ligands they contain (see Experimental Section).

**FAB Mass Spectra.** The  $\text{FAB}^-$  (**1**) or  $\text{FAB}^+$  (**7–9**) mass spectra measured show in all cases the corresponding mass peak with relative intensity in the range of 6% (**7**) to 100% (**1**). The spectra of complexes **7–9** are dominated by the signals corresponding to the  $[\text{AuPR}_3]^+$ ,  $[\text{Au}(\text{PR}_3)_2]^+$ , and  $[\text{PtCl}(\text{PR}_3)_2]^+$  fragments.

## Conclusions

3,5-Diethynylpyridine has been used to prepare a family of mono- (**1**), di- (**2–7**), and trinuclear (**8**, **9**) homo- or heteronuclear complexes containing  $\sigma$  monoalkynyl ( $\text{C}\equiv\text{C}(\text{Py})\text{C}\equiv\text{CH}$ , **1**) or  $\sigma$  dialkynyl ( $\mu\text{-C}\equiv\text{C}(\text{Py})\text{C}\equiv\text{C}$ , **2–7**) ligands. Furthermore, complex **6** is a metalloligand from which homo- (**8**) or heterotrinuclear (**9**) complexes can be obtained after coordination of a third metal center to the pyridinic nitrogen atom.

**Acknowledgment.** We thank the Ministerio de Ciencia y Tecnología, FEDER (BQU2001-0133), for financial support and Fundación Séneca (Comunidad Autónoma de la Región de Murcia, Spain) for a grant to M.M.A.F.

**Supporting Information Available:** A crystallographic file in CIF format for complex **6**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM049509J

(37) McArdle, C. P.; Van, S.; Jennings, M. C.; Puddephatt, R. J. *J. Am. Chem. Soc.* **2002**, *124*, 3959. Eisler, D.; Hong, W.; Jennings, M. C.; Puddephatt, R. J. *Organometallics* **2002**, *21*, 3955.

(38) Naulty, R. H.; Cifuentes, M. P.; Humphrey, M. G.; Houbrechts, S.; Boutton, C.; Persoons, A.; Heath, G. A.; Hockless, D. C. R.; Luther-Davies, B.; Samoc, M. J. *Chem. Soc., Dalton Trans.* **1997**, 4167. Hong, X.; Cheung, K. K.; Guo, C.-X.; Che, C.-M. *J. Chem. Soc., Dalton Trans.* **1994**, 1867.

(39) Lawrance, G. A. *Chem. Rev.* **1986**, *86*, 17.

(40) Silverstein, R. M.; Bassler, G. C.; Morrill, T. C. *Spectrometric Identification of Organic Compounds*, 4th ed.; 1981, p 110.

(41) Mingos, D. M. P.; Yau, J.; Menzer, S.; Williams, D. J. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 1894.