

# Meglumine: A Novel and Efficient Catalyst for One-Pot, Three-Component Combinatorial Synthesis of Functionalized 2-Amino-4*H*-pyrans

Rui-Yun Guo, Zhi-Min An, Li-Ping Mo, Rui-Zhi Wang, Hong-Xia Liu, Shu-Xia Wang, and Zhan-Hui Zhang\*

College of Chemistry & Material Science, Hebei Normal University, Shijiazhuang 050024, China

**Supporting Information** 

**ABSTRACT:** An efficient one-pot synthesis of functionalized 2-amino-4*H*-pyrans by a meglumine-catalyzed three-component reaction has been developed. A broad range of substrates including aromatic and heteroaromatic aldehydes, isatin derivatives, and acenaphthenequinone are condensed with enolizable C–H activated compounds and alkylmalonates to give the desired products in high to excellent yields. This methodology provides an alternative approach for rapid access to construct a diversity-oriented library of 4*H*-pyrans.



KEYWORDS: one-pot synthesis, three-component reaction, functionalized 2-amino-4H-pyrans, meglumine

yran and its derivatives are an important class of heterocyclic compounds, which constitute the key core of various natural products<sup>1</sup> as well as photochromic materials.<sup>2</sup> They exhibit a wide range of biological activities such as anticancer, antimicrobial,<sup>4</sup> antioxidant,<sup>5</sup> and antiproliferative properties.<sup>6</sup> 4H-Pyran derivatives are also potential calcium channel antagonists, which are structurally similar to biologically active 1,4-dihydropyridines.<sup>7</sup> In addition, amino-4H-pyrans are often used in cosmetics and pigments, or are utilized as potentially biodegradable agrochemicals. The 4H-pyran derivatives bearing a nitrile functionality are also useful intermediates for the synthesis of various compounds such as pyridones, 1,4dihydropyridines, lactones, pyranopyrazoles, imidoesters, and aminopyrimidines.<sup>8</sup> Because of the important aforementioned properties of pyran derivatives, preparation of this heterocyclic nucleus has gained great importance in organic synthesis.<sup>9</sup> The most straightforward synthesis of this heterocyclic system involves a three-component coupling of an aldehyde with alkylmalonates and diverse enolizable C-H activated acidic compounds in the presence of homogeneous or heterogeneous pounds in the presence of nonogeneous of neterogeneous catalysts such as diammonium hydrogen phosphate,<sup>10</sup> *N*-methyl-imidazole,<sup>11</sup> 4-(dimethylamino)pyridine (DMAP),<sup>12</sup> lithium bromide,<sup>13</sup> hexamethylenetetramine,<sup>14</sup> 1,8-diazabicyclo[5.4.0]-undec-7-ene (DBU),<sup>15</sup> potassium phthalimide-*N*-oxyl,<sup>16</sup> li-pase,<sup>17</sup> Ce(SO<sub>4</sub>)<sub>2</sub>·4H<sub>2</sub>O,<sup>18</sup> cerium(III) chloride,<sup>19</sup> tetrabutyl-ammonium fluoride (TBAF),<sup>20</sup> or a basic ionic liquid.<sup>21</sup> This reaction proceeds in glycerol<sup>22</sup> or choline chloride-urea.<sup>23</sup> With widespread applications and bioactivity, the development of efficient, economically and environmentally benign synthetic

methodology for the preparation of these heterocyclic compounds is highly desirable.

Following the increasing environmental and economic considerations, the search for enviro-economic synthetic methods for organic reactions has received overwhelming attention. In this regard, efforts are being made to replace the expensive and hazardous catalysts with biodegradable materials, which are safe, inexpensive, harmless, and environmentally benign. Recently, Gu et al. introduced meglumine and gluconic acid aqueous solution as a promoting medium and catalyst for the multicomponent reaction of  $\beta$ -ketosulfones and formaldehyde.<sup>24</sup> Meglumine is an amino sugar derived from sorbitol with molecular formula C<sub>7</sub>H<sub>17</sub>NO<sub>5</sub> (Figure 1). In particular, the



low toxicity of meglumine allows its use in the formulation of pharmaceuticals as an excipient and in conjunction with iodinated compounds in contrast media such as diatrizoate meglumine and iodipamide meglumine.<sup>25</sup> Meglumine possesses environmentally benign properties such as biodegradability and physiological inertness. Additionally, meglumine is inexpensive,

```
Received:August 15, 2013Revised:September 18, 2013Published:October 18, 2013
```

ACS Publications © 2013 American Chemical Society

## **ACS Combinatorial Science**

noncorrosive, stable to air and moisture, and readily available in the market. It contains an amino group, primary and secondary hydroxyl groups that can activate the nucleophilic as well as electrophilic components of the reactions by hydrogen bonding and donation of a lone pair of electrons, respectively. Multicomponent coupling reactions (MCRs) have been acknowledged as powerful and bond-forming efficient tools for the construction of novel and structurally complex molecules with a minimum of reaction steps and simple workup procedures. Thus, the combination of MCRs, a biodegradable catalyst, and a green solvent will make the chemical reaction more environmentally as well as economically viable.<sup>26</sup>

Considering the above subjects and continuing our efforts on the development of new MCRs and environmental benign synthetic methodologies,<sup>27</sup> we wish to report a facile, one-pot, three-component process for the synthesis of functionalized 2-amino-4*H*-pyrans from the reaction of aldehydes, isatin derivatives, or acenaphthenequinone with enolizable C–H activated compounds and alkylmalonates in the presence of meglumine at room temperature (Scheme 1).

Scheme 1. Synthesis of Functionalized 2-amino-4*H*-pyrans Catalyzed by Meglumine



To explore the generality and feasibility of meglumine catalyzed MCRs, we investigated the optimization of the reaction conditions using benzaldehyde, 5,5-dimethylcyclohexane-1,3dione, and malononitrile as model reactants. Various catalysts were examined, and the results are summarized in Table 1. To justify the significance of catalyst in this three-component process, the reaction was first performed in the absence of the catalyst wherein only 40% yield of the target product was observed even at prolonged reaction time (Table 1, entry 1). In the presence of potassium carbonate, sodium acetate, or ammonium acetate, the reaction gave 55%, 45%, and 50% isolated yield of product after 1 h, respectively. In addition, sodium dodecylbenzenesulfonate, triethylamine, tetrabutylammonium bromide, and L-proline could also catalyze this transformation, but no improvement was observed (Table 1, entries 5-8). The reaction was then performed in the presence of metal oxides such as CuO, CeO<sub>2</sub>, ZnO, and MgO (Table 1, entries 9-12). However, these catalysts were less effective. We then tested the reaction in the presence of meglumine. Meglumine was found to be the best catalyst for this process, and the desired product 6{1,1,1}was formed in 97% yield in just 5 min (Table 1, entry 13). We think that in addition to the amino group, the presence of multiple hydroxyl groups in the structure of meglumine plays an important role for the formation of product because of the stabilization of the corresponding intermediates and transition states by hydrogen bonding.

Malononitrile <sup><i>a</i></sup>							
	$\begin{array}{c} CHO \\ \bullet \\ $		>NH₂ `CN }				
entry	catalyst	time (min)	yield (%) <sup>b</sup>				
1	no	60	40				
2	K <sub>2</sub> CO <sub>3</sub>	60	55				
3	CH <sub>3</sub> COONa	60	45				
4	CH <sub>3</sub> COONH <sub>4</sub>	60	50				
5	sodium dodecylbenzenesulfonate	60	65				
6	triethylamine	60	60				
7	tetrabutylammonium bromide	60	50				
8	L-proline	60	50				
9	CuO	60	45				
10	CeO <sub>2</sub>	60	40				
11	nano ZnO	60	45				
12	nano MgO	60	40				
13	meglumine	5	97				

Table 1. Influence of Different Catalysts on the Reaction

of Benzaldehyde, 5,5-Dimethylcyclohexane-1,3-dione, and

<sup>a</sup>Experimental conditions: benzaldehyde (1 mmol), 5,5-dimethylcyclohexane-1,3-dione, (1 mmol), malononitrile (1 mmol), catalyst (5 mol %), EtOH-H<sub>2</sub>O (1:1, 2 mL), room temperature. <sup>b</sup>Isolated yields.

Table 2. Optimization of Reaction Conditions<sup>a</sup>

entry	catalyst loading (mol %)	solvent	time (min)	yield (%) <sup>b</sup>
1	5	no	10	93
2	5	MeOH	20	80
3	5	EtOH	10	95
4	5	PEG 400	60	70
5	5	H <sub>2</sub> O	60	30
6	5	EtOH-H <sub>2</sub> O (1:1)	5	97
7	5	EtOH-H <sub>2</sub> O (2:1)	10	95
8	5	EtOH-H <sub>2</sub> O (1:2)	10	93
9	1	EtOH-H <sub>2</sub> O (1:1)	5	88
10	3	EtOH-H <sub>2</sub> O (1:1)	5	92
$11^{c}$	5	EtOH- $H_2O(1:1)$	5	96
$12^d$	5	EtOH-H <sub>2</sub> O (1:1)	5	96, 95, 93

<sup>*a*</sup>Experimental conditions: benzaldehyde (1 mmol), 5,5-dimethylcyclohexane-1,3-dione (1 mmol), malononitrile (1 mmol), solvent (2 mL), room temperature. <sup>*b*</sup>Isolated yields. <sup>*c*</sup>The reaction was carried out in 50 mmol scale. <sup>*d*</sup>Catalyst was reused three times.

Next, we screened the effect of the solvent and catalyst loading (Table 2). The reaction was not satisfactory in water, possibly because of incomplete homogeneity of the reaction mixture. Further studies showed that aqueous-ethanol (1:1 v/v) was the best choice of solvent for this transformation. The reaction was also carried out using different amounts of the catalyst, and the results showed that 5 mol % of catalyst was the best choice for completing the reaction in aqueous-ethanol. By decreasing the amount of catalyst to 3 mol % relative to substrate, the yield of product decreased (Table 2, entry 10). Consequently, the best result was obtained by using 5 mol % of meglumine as the catalyst and aqueous-ethanol (1:1 v/v) as the solvent (Table 2, entry 6).

Furthermore, when the reaction was scaled up to 50 mmol, an excellent result was obtained in the same time (Table 2, entry 11). For utility in large scale syntheses we demonstrated



Figure 2. Diversity of reagents.

the recyclability of meglumine for three consecutive cycles with almost the same catalytic activity (Table 2, entry 12). After completion of each reaction, the precipitated product was filtered and washed with aqueous ethanol. The crude product was purified by recrystallization from ethanol to afford the desired product. To recover the catalyst, the filtrate was dried under reduced pressure and the recovered catalyst was washed with diethyl ether and reused after drying.

To investigate the scope and limitation of this catalytic process, 29 aldehydes  $1\{1-29\}$ , 3 isatins  $2\{30-32\}$ , acenaphthenequinone 3{33}, 3 cyclic ketones 4{1-3}, 1 acyclic 1,3dicarbonyl compounds 4{4}, 2 naphthols 4{5-6}, 4-hydroxycoumarin  $4\{7\}$ , 2 barbituric acids  $4\{8-9\}$ , and 3 acetonitrile derivatives  $5\{1-3\}$  were chosen for library validation (Figure 2). We first examined the reaction of 5,5-dimethylcyclohexane-1,3dione  $4\{1\}$  and malononitrile  $5\{1\}$  with a diverse array of aldehydes. As evident from Table 3, this multicomponent reaction proceeded efficiently, and the desired products were obtained in high to excellent yields in relatively short times. The nature of the functional group on the aromatic ring of the aldehyde exerted a slight influence on the product yield and reaction time. It was observed that arylaldehydes carrying an electron-withdrawing group reacted faster with better yield than those having an electron-donating group. The reaction was also sensitive to the steric environment of the aromatic aldehyde with longer reaction times being required for benzaldehydes containing substituents at the 2-position. Heteroaromatic aldehydes, such as furan-2-carbaldehyde, thiophene-2-carbaldehyde, and pyridine-4-carbaldehyde, readily participated in this

transformation, affording the potentially biologically important 2-amino-4*H*-pyrans in high yields (Table 3, entries 27–29). To further demonstrate the scope of the substrates subject to meglumine catalysis, isatins were used as reactants with 5,5-dimethylcyclohexane-1,3-dione and malononitrile. Gratifyingly, it was found that the expected products were obtained in excellent yields, regardless of the effect of substitution on the isatins (Table 3, entries 30–32). In the case of acenaphthene-quinone  $3{33}$ , the desired spiro-4*H*-pyrans  $8{33,1,1}$  was also produced in 93% yield (Table 3, entry 33). It should be noted that the reaction with methyl cyanoacetate or ethyl cyano-acetate required longer reaction times than those with malono-nitrile, which was probably due to their lower reactivity (Table 3, entries 34-37).

Subsequently, a variety of enolizable C–H activated compounds such as carbonyl compounds possessing a reactive  $\alpha$ -methylene group cyclohexane-1,3-dione and cyclopentane-1,3-dione,  $\alpha$ -naphthol and  $\beta$ -naphthol, 4-hydroxycoumarin, and barbituric acids were evaluated in this three-component reaction. In all cases, the corresponding functionalized 2-amino-4H-pyrans were formed in high to excellent yields. Furthermore, the reaction may also be extended to acyclic 1,3-dicarbonyl compounds such as acetylacetone, although longer reaction times were required (Table 3, entries 44–46). The above results clearly indicate the present catalytic procedure is extendable to a wide variety of substrates to construct a diversity-oriented library of 4H-pyrans.

In summary, we have developed a simple and efficient one-pot, three-component, meglumine catalyzed synthesis of functionalized

# **ACS Combinatorial Science**

#### Table 3. One-Pot Synthesis of 2-Amino-4H-pyran Derivatives

			mp (°C)							mp (°C)		
entry	product	time (min)	yield (%) <sup>a</sup>	found	reported	entr	ry	product	time (min)	yield $(\%)^a$	found	reported
1	<b>6</b> {1,1,1}	5	97	225-226	224-226 <sup>22</sup>	41		6{23,2,1}	5	96	235-236	235-236 <sup>13</sup>
2	<b>6</b> {2,1,1}	15	92	200-201	203-205 <sup>17</sup>	42	5	<b>6</b> {18,3,1}	5	95	218-219	216-218 <sup>12</sup>
3	<b>6</b> {3,1,1}	13	94	190-191	$188 - 190^{18}$	43		<b>6</b> {26,3,1}	5	96	178-180	
4	<b>6</b> {4,1,1}	13	95	200-201	201-20316	44	ł	6{1,4,1}	150	92	165-166	164 <sup>32</sup>
5	<b>6</b> {5,1,1}	25	92	209-210	209-210 <sup>13</sup>	45		<b>6</b> {18,4,1}	120	93	155-156	154 <sup>32</sup>
6	<b>6</b> { <i>6</i> ,1,1}	20	91	177-178		46	<b>,</b>	<b>6</b> {23,4,1}	90	94	169-170	170 <sup>32</sup>
7	<b>6</b> {7,1,1}	20	93	194-195	$195 - 196^{20}$	47	,	6{2,5,1}	15	94	207-208	
8	<b>6</b> {8,1,1}	12	95	224-225	$223 - 225^{17}$	48	;	6{9,5,1}	15	94	206-207	$205 - 206^{15}$
9	<b>6</b> {9,1,1}	12	93	211-212	219-222 <sup>10</sup>	49	)	<b>6</b> {18,5,1}	15	95	245-246	$243 - 244^{12}$
10	<b>6</b> {10,1,1}	17	93	227-228	226-228 <sup>19</sup>	50	)	6{23,5,1}	10	97	235-236	236 <sup>35</sup>
11	<b>6</b> {11,1,1}	18	93	241-242	240-242 <sup>19</sup>	51		6{25,5,1}	15	95	218-220	
12	<b>6</b> {12,1,1}	14	93	209-210	$208 - 209^{21}$	52	5	<b>6</b> {9,6,1}	140	90	255-256	255 <sup>35</sup>
13	<b>6</b> {13,1,1}	13	93	205-206	204-205 <sup>10</sup>	53		<b>6</b> {16,6,1}	120	92	259-60	260 <sup>35</sup>
16	<b>6</b> {14,1,1}	20	93	215-216		54	ł	<b>6</b> {23,6,1}	110	93	187-188	187 <sup>35</sup>
15	<b>6</b> {15,1,1}	7	96	233-234	232-233 <sup>20</sup>	55		<b>6</b> {9,7,1}	20	93	254-255	255-256 <sup>22</sup>
16	<b>6</b> {16,1,1}	8	95	216-217	215-21611	56	5	<b>6</b> {15,7,1}	15	96	248-249	$247 - 249^{14}$
17	<b>6</b> {17,1,1}	5	96	235-236	236-23710	57	,	<b>6</b> {18,7,1}	17	93	265-266	$264 - 266^{12}$
18	<b>6</b> {18,1,1}	5	96	215-216	213-215 <sup>22</sup>	58	;	<b>6</b> {26,7,1}	13	96	208-210	
19	<b>6</b> {19,1,1}	10	94	193-195	192-195 <sup>10</sup>	59	)	<b>6</b> {2,8,1}	30	95	232-233	$231 - 232^{22}$
20	<b>6</b> {20,1,1}	5	93	229-230	$228 - 230^{28}$	60	)	<b>6</b> {18,8,1}	19	95	264-265	$263 - 264^{22}$
21	<b>6</b> {21,1,1}	10	95	241-242		61		6{23,8,1}	15	96	263-264	$262 - 263^{22}$
22	<b>6</b> {22,1,1}	5	96	215-216	213-217 <sup>10</sup>	62	5	6{9,9,1}	20	93	203-204	203 <sup>33</sup>
23	<b>6</b> {23,1,1}	5	96	185-186	$181 - 184^{10}$	63		<b>6</b> {18,9,1}	18	95	239-240	$238 - 240^{33}$
24	<b>6</b> {24,1,1}	10	93	227-228	226-228 <sup>13</sup>	64	ł	<b>6</b> {26,9,1}	15	97	>300	
25	<b>6</b> {25,1,1}	5	96	224-226		65		7{30,2,1}	15	94	298-299	$297 - 298^{22}$
26	<b>6</b> {26,1,1}	5	95	218-219	217-218 <sup>29</sup>	66	5	7{30,3,1}	25	97	>300	>300 <sup>34</sup>
27	<b>6</b> {27,1,1}	15	95	226-227	$225 - 226^{13}$	67	,	7{30,5,1}	30	95	221-222	$220 - 222^{22}$
28	<b>6</b> {28,1,1}	15	92	221-222	$222 - 223^{22}$	68	;	7{30,6,1}	35	93	235-236	$233 - 235^{22}$
29	<b>6</b> {29,1,1}	14	96	153		69	)	7{30,7,1}	20	95	>300	>300 <sup>22</sup>
30	7{30,1,1}	18	96	290-292	291-293 <sup>22</sup>	70	)	7{30,8,1}	30	95	298-299	296-298 <sup>22</sup>
31	7{31,1,1}	18	93	278-279	$277 - 278^{22}$	71		7{30,9,1}	25	97	229-230	$228 - 229^{36}$
32	7{32,1,1}	10	95	303-304	302-304 <sup>30</sup>	72	5	7{31,5,1}	40	93	>300	
33	<b>8</b> {33,1,1}	20	93	269-270	$268 - 270^{31}$	73		7{32,5,1}	15	95	>300	
34	<b>6</b> {18,1,2}	20	91	170-171	$171^{10}$	74		7{32,9,1}	20	95	225-227	
35	<b>6</b> {23,1,2}	18	92	187-188	$186 - 188^{10}$	75		<b>8</b> {33,2,1}	25	94	>300	>300 <sup>37</sup>
36	<b>6</b> {18,1,3}	22	90	150-151	150 <sup>10</sup>	76	5	<b>8</b> {33,3,1}	25	95	276-277	
37	<b>6</b> {23,1,3}	20	91	180-181	179-181 <sup>10</sup>	77	,	<b>8</b> {33,7,1}	30	80	>300	>300 <sup>31</sup>
38	<b>6</b> {9,2,1}	10	95	232-233	$233 - 235^{22}$	78	;	<b>8</b> {33,8,1}	35	90	>300	>300 <sup>31</sup>
39	<b>6</b> {18,2,1}	7	96	242-243	241-24312	79	)	<b>8</b> {33,9,1}	40	92	>300	>300 <sup>37</sup>
40	<b>6</b> {22,2,1}	6	93	202-203	$201 - 202^{17}$							
<i><sup>a</sup></i> Isolat	ed yield.											

2-amino-4*H*-pyrans using readily available starting materials. The salient features of this new methodology are broad substrate scope, room temperature reaction conditions, short reaction times, high yields of the products, operational simplicity, reusability of catalyst and the absence of hazardous organic solvents, which make it an applicable method for constructing diverse libraries of 4*H*-pyrans.

# EXPERIMENTAL PROCEDURES

**General Information.** All solvents and chemicals used in this work were obtained commercially and were used as received. Melting points were determined using an X-4 apparatus and are uncorrected. IR spectra were obtained as potassium bromide pellets with a Bruker-TENSOR 27 spectrometer. NMR spectra were recorded with a Bruker DRX-500 spectrometer at 500 MHz (<sup>1</sup>H) and 125 MHz (<sup>13</sup>C) using DMSO-*d*<sub>6</sub> or CDCl<sub>3</sub>

as the solvent with TMS as internal standard. Elemental analysis was performed by using a Vario EL III CHNOS elemental analyzer.

Typical Procedure for Synthesis of 2-Amino-7,7dimethyl-5-oxo-4-phenyl-5,6,7,8-tetrahydro-4H-chromene-3-carbonitrile 6 {1,1,1}. A mixture of benzaldehyde  $1\{1\}$  (0.11 g, 1 mmol)), 5,5-dimethylcyclohexane-1,3-dione  $2\{1\}$  (0.14 g, 1 mmol), malononitrile  $3\{1\}$  (0.07 g, 1 mmol), and meglumine (5 mol %, 0.01g) in EtOH-H<sub>2</sub>O (1:1, 2 mL) was stirred at room temperature for 5 min. The progress of the reaction was monitored by TLC (hexane-EtOAc, 5:1). After completion of the reaction, the precipitated product was filtered and washed with aqueous ethanol (5 mL). The crude product was purified by recrystallization from ethanol to afford the desired product. To recover the catalyst, the filtrate was dried under reduced pressure, and recovered catalyst was washed with diethyl ether (2 mL) twice and reused after drying. *Compound* **6**{6,1,1}. White solid; IR (KBr): 3545, 3327, 3215, 2937, 2193, 1683, 1653, 1606, 1508, 1367, 1251, 1211, 1174, 1035, 842 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.92 (t, *J* = 7.0 Hz, 3H, CH<sub>3</sub>), 1.03 (s, 3H, CH<sub>3</sub>), 1.10 (s, 3H, CH<sub>3</sub>), 1.33–1.44 (m, 4H, 2 × CH<sub>2</sub>), 1.75 (quin, *J* = 7.0 Hz, 2H, CH<sub>2</sub>), 2.19 and 2.24 (AB system, *J* = 16.5 Hz, 2H, CH<sub>2</sub>), 2.44 (s, 2H, CH<sub>2</sub>), 3.90 (t, *J* = 7.0 Hz, 2H, OCH<sub>2</sub>), 4.35 (s, 1H, CH), 4.49 (s, 2H, NH<sub>2</sub>), 6.80 (d, *J* = 8.5 Hz, 2H, HAr), 7.13 (d, *J* = 8.5 Hz, 2H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  14.0, 22.5, 27.6, 28.2, 28.9, 29.0, 32.2, 34.8, 40.7, 50.7, 63.1, 67.9, 114.2, 114.5, 115.6, 119.0, 128.6, 133.6, 135.4, 157.7, 158.2, 161.4, 196.2 ppm; Anal. Calcd for C<sub>23</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub>: C, 72.60; H, 7.42; N, 7.36. Found: C, 72.42; H, 7.26; N, 7.55.

*Compound* **6**{14,1,1}. White solid; IR (KBr): 3433, 3331, 3219, 2955, 2185, 1687, 1672, 1602, 1518, 1425, 1369, 1344, 1253, 1138, 1039, 852 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  1.03 (s, 3H, CH<sub>3</sub>), 1.11(s, 3H, CH<sub>3</sub>), 2.19 and 2.24 (AB system, *J* = 16.5 Hz, 2H, CH<sub>2</sub>), 2.44 (s, 2H, CH<sub>2</sub>), 4.37(s, 1H, CH), 4.50 (s, 2H, NH<sub>2</sub>), 5.13 (s, 2H, CH<sub>2</sub>), 6.87 (d, *J* = 8.5 Hz, 2H, HAr), 7.17 (d, *J* = 8.5 Hz, 2H, HAr), 7.59 (d, *J* = 8.5 Hz, 2H, HAr), 8.24 (d, *J* = 8.5 Hz, 2H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  27.9, 28.8, 31.3, 32.2, 34.4, 34.9, 40.7, 50.7, 63.9, 114.2, 118.8, 125.5, 127.0, 140.1, 149.7, 157.5, 161.5, 196.0 ppm; Anal. Calcd for C<sub>25</sub>H<sub>23</sub>N<sub>3</sub>O<sub>5</sub>: C, 67.41; H, 5.20; N, 9.43. Found: C, 67.22; H, 5.38; N, 9.62.

*Compound* **6**{21,1,1}. White solid; IR (KBr): 3421, 3336, 3217, 2960, 2191, 1700, 1660, 1480, 1430, 1365, 1251, 1211, 1138, 1039, 875 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  1.05 (s, 3H, CH<sub>3</sub>), 1.12 (s, 3H, CH<sub>3</sub>), 2.24 (s, CH<sub>2</sub>), 2.43 and 2.50 (AB system, *J* = 18.0 Hz, 2H, CH<sub>2</sub>), 4.31 (s, 1H, OH), 4.59 (s, 2H, NH<sub>2</sub>), 5.81 (s, 1H, CH), 7.31 (s, 2H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  27.9, 28.8, 31.3, 32.2, 34.4, 34.9, 40.7, 50.7, 63.9, 114.2, 118.8, 125.5, 127.0, 140.1, 149.7, 157.5, 161.5, 196.0 ppm; Anal. Calcd for C<sub>18</sub>H<sub>16</sub>Br<sub>2</sub>N<sub>2</sub>O<sub>3</sub>: C, 46.18; H, 3.44; N, 5.98. Found: C, 45.99; H, 3.60; N, 6.16.

*Compound* **6**{*25*,*1*,*1*}. White solid; IR (KBr): 3350, 3259, 3176, 2966, 2191, 1683, 1658, 1606, 1469, 1417, 1371, 1330, 1219, 1124, 1039, 823 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  1.03 (s, 3H, CH<sub>3</sub>), 1.12 (s, 3H, CH<sub>3</sub>), 2.19 and 2.24 (AB system, *J* = 16.5 Hz, 2H, CH<sub>2</sub>), 2.48 (s, 2H, CH<sub>2</sub>), 4.47 (s, 1H, CH), 4.60 (s, 2H, NH<sub>2</sub>), 7.42 (t, *J* = 7.5 Hz, 1H, HAr), 7.42 (s, 1H), 7.47 (d, *J* = 7.5 Hz, 1H, HAr), 7.48 (d, *J* = 7.5 Hz, 1H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  27.4, 29.0, 32.2, 35.6, 40.7, 50.6, 62.9, 113.5, 118.2, 124.1 (q, <sup>3</sup>*J*<sub>FC</sub> = 3.75 Hz), 125.2 (q, <sup>1</sup>*J*<sub>FC</sub> = 271.0 Hz), 129.0, 131.1 (q, <sup>2</sup>*J*<sub>FC</sub> = 32.0 Hz), 131.4, 144.2, 157.6, 161.9, 195.7 ppm; Anal. Calcd for C<sub>19</sub>H<sub>17</sub>F<sub>3</sub>N<sub>2</sub>O<sub>2</sub>: C, 62.98; H, 4.73; N, 7.73. Found: C, 63.12; H, 4.90; N, 7.55.

*Compound* **6**{29,1,1}. White solid; IR (KBr): 3419, 3394, 3216, 2956, 2187, 1683, 1672, 1600, 1421, 1373, 1251, 1205, 1138, 1037, 854 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  1.04 (s, 3H, CH<sub>3</sub>), 1.13 (s, 3H, CH<sub>3</sub>), 2.21 and 2.27 (AB system, *J* = 16.5 Hz, 2H, CH<sub>2</sub>), 2.48 (s, 2H, CH<sub>2</sub>), 4.40 (s, 1H, CH), 4.69 (s, 2H, NH<sub>2</sub>), 7.18 (d, *J* = 6.0 Hz, 2H, HAr), 8.54 (d, *J* = 6.0 Hz, 2H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  27.6, 28.9, 32.3, 35.1, 40.7, 50.5, 61.8, 112.9, 118.1, 122.8, 150.1, 151.6, 157.9, 162.3, 195.6 ppm; Anal. Calcd for C<sub>17</sub>H<sub>17</sub>N<sub>3</sub>O<sub>2</sub>: C, 69.14; H, 5.80; N, 14.23. Found: C, 68.98; H, 5.96; N, 14.05.

*Compound* **6**{*26,3,1*}. White solid; IR (KBr): 3404, 3333, 3211, 2937, 2193, 1670, 1643, 1597, 1442, 1375, 1329, 1234, 1166, 1111, 1066, 854 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  2.49–2.52 (m, 2H, CH<sub>2</sub>), 2.74–2.77 (m, 2H, CH<sub>2</sub>), 4.44 (s, 1H, CH), 4.79 (s, 2H, NH<sub>2</sub>), 7.40 (d, *J* = 8.0 Hz, 2H, HAr),

7.59 (d, J = 8.0 Hz, 2H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  25.0, 33.6, 35.7, 61.4, 117.9, 118.3, 125.7 (q, <sup>3</sup> $J_{FC} = 3.5$  Hz), 126.2 (q, <sup>1</sup> $J_{FC} = 270.6$  Hz), 129.9 (q, <sup>2</sup> $J_{FC} = 32.1$  Hz), 144.6, 159.0, 175.3, 200.8 ppm; Anal. Calcd for C<sub>16</sub>H<sub>11</sub>F<sub>3</sub>N<sub>2</sub>O<sub>2</sub>: C, 60.00; H, 3.46; N, 8.75. Found: C, 58.91; H, 3.52; N, 8.93.

*Compound* **6**{2,5,1}. White solid; IR (KBr): 3471, 3396, 3198, 2933, 2185, 1662, 1614, 1573, 1489, 1458, 1377, 1261, 1192, 1105, 1022, 756 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  3.87 (s, 3H, OCH<sub>3</sub>), 4.70 (s, 2H, NH<sub>2</sub>), 5.42 (s, 1H, CH), 6.86 (t, *J* = 7.5 Hz, 1H, HAr), 6.91 (d, *J* = 8.0 Hz, 1H, HAr), 7.02 (dd, *J* = 7.5, 1.5 Hz, 1H, HAr), 7.12 (d, *J* = 8.5 Hz, 1H, HAr), 7.20 (td, *J* = 7.5, 1.5 Hz, 1H, HAr), 7.48 (d, *J* = 8.5 Hz, 1H, HAr), 7.51 (d, *J* = 8.5 Hz, 1H, HAr), 7.56 (t, *J* = 7.5 Hz, 1H, HAr), 7.78 (d, *J* = 8.0 Hz, 1H, HAr), 8.16 (d, *J* = 8.0 Hz, 1H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  34.2, 55.7, 60.7, 111.1, 117.9, 120.0, 120.7, 121.1, 123.2, 124.5, 126.2, 126.5, 127.7, 128.4, 129.6, 132.8, 133.2, 143.6, 156.8, 159.8 ppm; Anal. Calcd for C<sub>21</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>: C, 76.81; H, 4.91; N, 8.53. Found: C, 76.66; H, 5.10; N, 8.38.

*Compound* **6**{25,5,1}. White solid; IR (KBr): 3473, 3338, 3198, 2935, 2195, 1670, 1602, 1572, 1460, 1408, 1379, 1261, 1188, 1114, 1050, 763 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.81 (s, 2H, NH<sub>2</sub>), 4.97 (s, 1H, CH), 6.98 (d, *J* = 8.0 Hz, 1H, HAr), 7.45–7.54 (m, 5H, HAr), 7.56 (d, *J* = 8.0 Hz, 1H, HAr), 7.60 (t, *J* = 8.0 Hz, 1H, HAr), 7.81 (d, *J* = 8.0 Hz, 1H, HAr), 8.19 (d, *J* = 8.0 Hz, 1H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  41.4, 60.4, 116.3, 119.5, 120.9, 123.2, 124.0 (q, <sup>1</sup>*J*<sub>FC</sub> = 271.0 Hz), 124.4 (q, <sup>3</sup>*J*<sub>FC</sub> = 3.8 Hz), 124.8 (q, <sup>3</sup>*J*<sub>FC</sub> = 3.8 Hz), 125.0, 125.8, 126.9, 127.1, 127.8, 129.4, 131.3 (q, <sup>2</sup>*J*<sub>FC</sub> = 32.1 Hz), 131.7, 133.5, 143.4, 145.4, 159.3 ppm; Anal. Calcd for C<sub>21</sub>H<sub>13</sub>F<sub>3</sub>N<sub>2</sub>O: C, 68.85; H, 3.58; N, 7.65. Found: C, 69.02; H, 3.75; N, 7.48.

*Compound* **6**{*26,7,1*}. White solid; IR (KBr): 3325, 3194, 3072, 2877, 2198, 1716, 1602, 1377, 1112, 765 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  4.60 (s, 1H, CH), 7.47–7.53 (m, 6H, HAr and NH<sub>2</sub>), 7.68 (d, *J* = 8.5 Hz, 2H, HAr), 7.73 (t, *J* = 7.0 Hz, 1H, HAr), 7.91(d, *J* = 7.5 Hz, 1H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>)  $\delta$  37.3, 57.6, 103.6, 113.4, 117.1, 119.5, 123.0, 125.2, 125.8 (q, <sup>1</sup>*J*<sub>FC</sub> = 270.4 Hz), 125.9 (q, <sup>3</sup>*J*<sub>FC</sub> = 3.8 Hz), 128.3 (q, <sup>2</sup>*J*<sub>FC</sub> = 31.2 Hz), 129.1, 133.6, 148.4, 152.7, 154.3, 158.5, 160.6 ppm; Anal. Calcd for C<sub>20</sub>H<sub>13</sub>F<sub>3</sub>N<sub>2</sub>O<sub>3</sub>: C, 62.18; H, 3.39; N, 7.25. Found: C, 61.99; H, 3.58; N, 7.43.

*Compound* **6**{*26,9,1*}. White solid; IR (KBr): 3441, 3304, 3183, 2193, 1705, 1608, 1323, 1126, 839 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  3.07 (s, 3H, CH<sub>3</sub>), 3.35 (s, 3H, CH<sub>3</sub>), 4.45 (s, 1H, CH), 7.44 (s, 2H, NH<sub>2</sub>), 7.49 (d, *J* = 8.0 Hz, 2H, HAr), 7.66 (d, *J* = 8.0 Hz, 2H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>)  $\delta$  28.1, 29.6, 36.9, 58.2, 88.4, 119.3, 125.2 (q, <sup>1</sup>*J*<sub>FC</sub> = 270.2 Hz), 125.7 (q, <sup>3</sup>*J*<sub>FC</sub> = 3.8 Hz), 127.9 (q, <sup>2</sup>*J*<sub>FC</sub> = 31.2 Hz), 128.0, 149.3, 150.5, 151.8, 158.2, 160.9 ppm; Anal. Calcd for C<sub>17</sub>H<sub>13</sub>F<sub>3</sub>N<sub>4</sub>O<sub>3</sub>: C, 53.97; H, 3.46; N, 14.81. Found: C, 54.15; H, 3.62; N, 15.00.

*Compound 7*{31,5,1}. White solid; IR (KBr): 3483, 3300, 3066, 2966, 2195, 1651, 1597, 1107, 692 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  2.20 (s, 3H, CH<sub>3</sub>), 6.57 (d, *J* = 9.0 Hz, 1H, HAr), 6.88 (d, *J* = 7.5 Hz, 1H, HAr), 6.90 (s, 1H, HAr), 7.09 (d, *J* = 7.5 Hz, 1H, HAr), 7.43 (s, 2H, NH<sub>2</sub>), 7.57 (d, *J* = 9.0 Hz, 1H, HAr), 7.60 (t, *J* = 7.5 Hz, 1H, HAr), 7.69 (t, *J* = 7.5 Hz, 1H, HAr), 7.90 (d, *J* = 8.5 Hz, 1H, HAr), 8.27 (d, *J* = 8.5 Hz, 1H, HAr), 10.6 (s, 1H, NH) ppm; <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>)  $\delta$  21.0, 51.4, 55.0, 110.3, 115.4, 119.1, 121.2, 123.3, 123.7, 125.0, 125.9, 127.5, 127.8, 128.2, 130.0, 132.2, 133.5, 135.5, 139.8

144.0, 161.5, 179.3 ppm; Anal. Calcd for  $C_{22}H_{15}N_3O_2$ : C, 74.78; H, 4.28; N, 11.89. Found: C, 74.60; H, 4.45; N, 12.03.

*Compound 7*{32,5,1}. White solid; IR (KBr): 3481, 3298, 3196, 2860, 2189, 1627, 1527, 1338, 1107, 750 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  6.64 (d, J = 9.0 Hz, 1H, HAr), 7.22 (d, J = 9.0 Hz, 1H, HAr), 7.59–7.65 (m, 4H, NH<sub>2</sub> and HAr), 7.71 (t, J = 8.0 Hz, 1H, HAr), 7.93 (d, J = 8.0 Hz, 1H, HAr), 8.02 (s, 1H, HAr), 8.29 (t, J = 7.0 Hz, 2H, HAr), 11.4 (s, 1H, NH) ppm; <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>)  $\delta$  51.6, 53.6, 111.1, 113.7, 118.8, 121.3, 121.4, 123.3, 123.4, 125.3, 127.1, 127.6, 128.0, 128.2, 133.8, 136.1, 143.6, 144.3, 148.8, 161.6, 179.7 ppm; Anal. Calcd for C<sub>21</sub>H<sub>12</sub>N<sub>4</sub>O<sub>4</sub>: C, 65.62; H, 3.15; N, 14.58. Found: C, 65.80; H, 2.96; N, 14.66.

*Compound 7*{32,9,1}. White solid; IR (KBr): 3396, 3186, 2968, 2362, 2202, 1743, 1506, 1390, 754 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  3.02 (s, 3H, CH<sub>3</sub>), 3.36 (s, 3H, CH<sub>3</sub>), 7.03 (d, *J* = 8.5 Hz, 1H, HAr), 7.77 (s, 2H, NH<sub>2</sub>), 8.17 (d, *J* = 8.5 Hz, 1H, HAr), 8.19 (s, 1H, HAr), 11.3 (s, 1H, NH) ppm; <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>)  $\delta$  28.1, 29.8, 48.0, 56.6, 86.5, 109.9, 117.1, 120.3, 126.5, 135.2, 143.0, 149.1, 150.1, 153.0, 158.9, 160.1, 178.7 ppm; Anal. Calcd for C<sub>17</sub>H<sub>12</sub>N<sub>6</sub>O<sub>6</sub>: C, 51.52; H, 3.05; N, 21.21. Found: C, 51.70; H, 2.91; N, 21.39.

*Compound* **8**{33,3,1}. Light yellow solid; IR (KBr): 3471, 3362, 3194, 2929, 2193, 1712, 1589, 1361, 1026, 781 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  2.34 (t, J = 4.5 Hz, 2H, CH<sub>2</sub>), 2.88 (t, J = 4.5 Hz, 2H, CH<sub>2</sub>), 7.48 (d, J = 7.0 Hz, 1H, HAr), 7.62 (s, 2H, NH<sub>2</sub>), 7.71 (t, J = 7.5 Hz, 1H, HAr), 7.87 (t, J = 8.0 Hz, 1H, HAr), 8.00 (t, J = 9.0 Hz, 2H, HAr), 8.35 (d, J = 8.0 Hz, 1H, HAr) ppm; <sup>13</sup>C NMR (125 MHz, DMSO- $d_6$ )  $\delta$  25.5, 33.6, 51.4, 57.6, 116.3, 118.1, 121.5, 122.5, 125.5, 129.2, 129.5, 130.4, 131.8, 132.8, 141.2, 141.8, 161.1, 178.3, 200.6, 203.6 ppm; Anal. Calcd for C<sub>20</sub>H<sub>12</sub>N<sub>2</sub>O<sub>3</sub>: C, 73.16; H, 3.68; N, 8.53. Found: C, 72.98; H, 3.85; N, 8.66.

# ASSOCIATED CONTENT

# Supporting Information

Copies of <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra of all new compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

## AUTHOR INFORMATION

#### **Corresponding Author**

\*E-mail: zhanhui@mail.nankai.edu.cn. Fax: +86-311-80787431.

# Funding

This work is financially supported by the National Natural Science Foundation of China (20872025 and 21272053), Nature Science Foundation of Hebei Province (B2011205031), and the Training Programs of Innovation and Entrepreneurship for Undergraduates of Hebei Province (201310094021).

#### Notes

The authors declare no competing financial interest.

## REFERENCES

(1) Wickel, S. M.; Citron, C. A.; Dickschat, J. S. 2H-pyran-2-ones from trichoderma viride and trichoderma asperellum. *Eur. J. Org. Chem.* **2013**, 2906–2913.

(2) Wang, S.; Qi, Q. Z.; Li, C. P.; Ding, G. H.; Kim, S. H. Photoswitching of bisthienylethene using  $2D-\pi$ -A type pyran-based fluorescent dye for rewritable optical storage. *Dyes Pigm.* **2011**, *89*, 188–192.

(3) Patil, S. A.; Wang, J.; Li, X. S.; Chen, J. J.; Jones, T. S.; Hosni-Ahmed, A.; Patil, R.; Seibel, W. L.; Li, W.; Miller, D. D. New substituted 4H-chromenes as anticancer agents. Bioorg. Med. Chem. Lett. 2012, 22, 4458-4461.

(4) Makawana, J. A.; Patel, M. P.; Patel, R. G. Synthesis and antimicrobial evaluation of new pyrano [4,3-*b*]pyran and pyrano[3,2-*c*]chromene derivatives bearing a 2-thiophenoxyquinoline nucleus. *Arch. Pharm.* **2012**, *345*, 314–322.

(5) Saundane, A. R.; Vijaykumar, K.; Vaijinath, A. V. Synthesis of novel 2-amino-4-(5'-substituted-2'-phenyl-1*H*-indol-3'-yl)-6-aryl-4*H*-pyran-3-carbonitrile derivatives as antimicrobial and antioxidant agents. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 1978–1984.

(6) Venkatesham, A.; Rao, R. S.; Nagaiah, K.; Yadav, J. S.; RoopaJones, G.; Basha, S. J.; Sridhar, B.; Addlagatta, A. Synthesis of new chromeno-annulated cis-fused pyrano[3,4-c]pyran derivatives via domino Knoevenagel-hetero-Diels-Alder reactions and their biological evaluation towards antiproliferative activity. *MedChemComm* **2012**, *3*, 652–658.

(7) Kang, S. S.; Cooper, G.; Dunne, S. F.; Luan, C. H.; Surmeier, D. J.; Silverman, R. B. Antagonism of L-type Ca<sup>2+</sup> channels Ca<sub>v</sub>1.3 and Ca<sub>v</sub>1.2 by 1,4-dihydropyrimidines and 4*H*-pyrans as dihydropyridine mimics. *Bioorg. Med. Chem.* **2013**, *21*, 4365–4373.

(8) (a) Kumar, D.; Reddy, V. B.; Sharad, S.; Dube, U.; Kapur, S. A facile one-pot green synthesis and antibacterial activity of 2-amino-4H-pyrans and 2-amino-5-oxo-5,6,7,8-tetrahydro-4H-chromenes. *Eur. J. Med. Chem.* 2009, 44, 3805–3809. (b) Litvinov, Y. M.; Shestopalov, A. M. Synthesis, structure, chemical reactivity, and practical significance of 2-amino-4H-pyrans. In *Advances in Heterocyclic Chemistry*; Katritzky, A. R., Ed.; Academic Press: New York, 2011; Vol. 103, pp 175–260. (9) Tian, J. J.; Guo, H. Y. New progress in the synthesis of pyrane derivatives via multicomponent reactions. *Chin. J. Org. Chem.* 2011, 31, 2009–2018.

(10) Balalaie, S.; Bararjanian, M.; Sheikh-Ahmadi, M.; Hekmat, S.; Salehi, P. Diammonium hydrogen phosphate: an efficient and versatile catalyst for the one-pot synthesis of tetrahydrobenzo[b]pyran derivatives in aqueous media. *Synth. Commun.* **2007**, *37*, 1097–1108. (11) Lian, X. Z.; Huang, Y.; Li, Y. Q.; Zheng, W. J. A green synthesis of tetrahydrobenzo[b]pyran derivatives through three-component condensation using N-methylimidazole as organocatalyst. *Monatsh. Chem.* **2008**, *139*, 129–131.

(12) Khan, A. T.; Lal, M.; Ali, S.; Khan, M. M. One-pot threecomponent reaction for the synthesis of pyran annulated heterocyclic compounds using DMAP as a catalyst. *Tetrahedron Lett.* **2011**, *52*, 5327–5332.

(13) Sun, W. B.; Zhang, P.; Fan, J.; Chen, S. H.; Zhang, Z. H. Lithium bromide as a mild, efficient, and recyclable catalyst for the one-pot synthesis of tetrahydro-4*H*-chromene derivatives in aqueous media. *Synth. Commun.* **2010**, *40*, 587–594.

(14) Wang, H. J.; Lu, J.; Zhang, Z. H. Highly efficient threecomponent, one-pot synthesis of dihydropyrano[3,2-c]chromene derivatives. *Monatsh. Chem.* **2010**, *141*, 1107–1112.

(15) Khurana, J. M.; Nand, B.; Saluja, P. DBU: a highly efficient catalyst for one-pot synthesis of substituted 3,4-dihydropyrano[3,2-c]chromenes, dihydropyrano[4,3-b]pyranes, 2-amino-4H-benzo[h] chromenes and 2-amino-4H benzo[g]chromenes in aqueous medium. *Tetrahedron* **2010**, *66*, 5637–5641.

(16) Dekamin, M. G.; Eslami, M.; Maleki, A. Potassium phthalimide-N-oxyl: a novel, efficient, and simple organocatalyst for the one-pot three-component synthesis of various 2-amino-4H-chromene derivatives in water. *Tetrahedron* **2013**, *69*, 1074–1085.

(17) Xu, J. C.; Li, W. M.; Zheng, H.; Lai, Y. F.; Zhang, P. F. One-pot synthesis of tetrahydrochromene derivatives catalyzed by lipase. *Tetrahedron* **2011**, *67*, 9582–9587.

(18) Islami, M. R.; Mosaddegh, E.  $Ce(SO_4)_2$ ·4H<sub>2</sub>O as a recyclable catalyst for an efficient, simple, and clean synthesis of 4H-Benzo[b]pyrans. *Phosphorus, Sulfur Silicon Relat. Elem.* **2009**, 184, 3134–3138.

(19) Sabitha, G.; Arundhathi, K.; Sudhakar, K.; Sastry, B. S.; Yadav, J. S. Cerium(III) chloride-catalyzed one-pot synthesis of tetrahydrobenzo[b]pyrans. Synth. Commun. 2009, 39, 433-442.

(20) Gao, S. J.; Tsai, C. H.; Tseng, C.; Yao, C. F. Fluoride ion catalyzed multicomponent reactions for efficient synthesis of 4*H*-chromene and *N*-arylquinoline derivatives in aqueous media. *Tetrahedron* **2008**, *64*, 9143–914.

(21) Ranu, B. C.; Banerjee, S.; Roy, S. A task specific basic ionic liquid, [bmIm]OH-promoted efficient, green and one-pot synthesis of tetrahydrobenzo[*b*]pyran derivatives. *Indian J. Chem.* **2008**, 47B, 1108–1112.

(22) Safaei, H. R.; Shekouhy, M.; Rahmanpur, S.; Shirinfeshan, A. Glycerol as a biodegradable and reusable promoting medium for the catalyst-free one-pot three component synthesis of 4*H*-pyrans. *Green Chem.* **2012**, *14*, 1696–1704.

(23) Revanna, C. N.; Swaroop, T. R.; Raghavendra, G. M.; Bhadregowda, D. G.; Mantelingu, K.; Rangappa, K. S. Practical and green protocol for the synthesis of substituted 4H-chromenes using room temperature ionic liquid choline chloride-urea. *J. Heterocycl. Chem.* **2012**, *49*, 851–855.

(24) Yang, J.; Li, H. Q.; Li, M. H.; Peng, J. J.; Gu, Y. L. Multicomponent reactions of  $\beta$ -ketosulfones and formaldehyde in a bio-based binary mixture solvent system composed of meglumine and gluconic acid aqueous solution. *Adv. Synth. Catal.* **2012**, *354*, 688–700.

(25) (a) Mattioni, B. E.; Kauffman, G. W.; Jurs, P. C.; Custer, L. L.; Durham, S. K.; Pearl, G. M. Predicting the genotoxicity of secondary and aromatic amines using data subsetting to generate a model ensemble. *J. Chem. Inf. Comput. Sci.* **2003**, *43*, 949–963. (b) Mozdarani, H.; Fadaei, S. Similar cytogenetic effects of sodium meglumine diatrizoate and sodium meglumine ioxithalamate in lymphocytes of patients undergoing brain CT scan. *Toxicol. Lett.* **1998**, *98*, 25–30.

(26) Gu, Y. L. Multicomponent reactions in unconventional solvents: state of the art. *Green Chem.* **2012**, *14*, 2091–2128.

(27) (a) Wang, H. J.; Mo, L. P.; Zhang, Z. H. Cerium ammonium nitrate-catalyzed multicomponent reaction for efficient synthesis of functionalized tetrahydropyridines. ACS Comb. Sci. 2011, 13, 181-185. (b) Zhang, X. N.; Li, Y. X.; Zhang, Z. H. Nickel chloride-catalyzed one-pot three-component synthesis of pyrazolophthalazinyl spirooxindoles. Tetrahedron 2011, 67, 7426-7430. (c) Deng, J.; Mo, L. P.; Zhao, F. Y.; Zhang, Z. H.; Liu, S. X. One-pot, three-component synthesis of a library of spirooxindole-pyrimidines catalyzed by magnetic nanoparticle supported dodecyl benzenesulfonic acid in aqueous media. ACS Comb. Sci. 2012, 14, 335-341. (d) Zhang, Z. H.; Zhang, X. N.; Mo, L. P.; Li, Y. X.; Ma, F. P. Catalyst-free synthesis of quinazoline derivatives using low melting sugar-urea-salt mixture as a solvent. Green Chem. 2012, 14, 1502-1506. (e) Deng, J.; Mo, L. P.; Zhao, F. Y.; Hou, L. L.; Yang, L.; Zhang, Z. H. Sulfonic acid supported on hydroxyapatite-encapsulated-\gamma-Fe2O3 nanocrystallites as a magnetically separable catalyst for one-pot reductive amination of carbonyl compounds. Green Chem. 2011, 13, 2576-2584. (f) Liu, Y. H.; Deng, J.; Gao, J. W.; Zhang, Z. H. Triflic acid-functionalized silica-coated magnetic nanoparticles as a magnetically separable catalyst for synthesis of gem-dihydroperoxides. Adv. Synth. Catal. 2012, 354, 441-447.

(28) Banerjee, S.; Horn, A.; Khatri, H.; Sereda, G. A green one-pot multicomponent synthesis of 4*H*-pyrans and polysubstituted aniline derivatives of biological, pharmacological, and optical applications using silica nanoparticles as reusable catalyst. *Tetrahedron Lett.* **2011**, *52*, 1878–1881.

(29) Balalaie, S.; Sheikh-Ahmadi, M.; Bararjanian, M. Tetra-methyl ammonium hydroxide: An efficient and versatile catalyst for the onepot synthesis of tetrahydrobenzo[b]pyran derivatives in aqueous media. *Catal. Commun.* **2007**, *8*, 1724–1728.

(30) Dandia, A.; Parewa, V.; Jain, A. K.; Rathore, K. S. Stepeconomic, efficient, ZnS nanoparticle-catalyzed synthesis of spirooxindole derivatives in aqueous medium via Knoevenagel condensation followed by Michael addition. *Green Chem.* **2011**, *13*, 2135–2145.

(31) Saeedi, M.; Heravi, M. M.; Beheshtiha, Y. S.; Oskooie, H. A. One-pot three-component synthesis of the spiroacenaphthylene derivatives. *Tetrahedron* **2010**, *66*, 5345–5348.

(32) Bhattacharyya, P.; Pradhan, K.; Paul, S.; Das, A. R. Nano crystalline ZnO catalyzed one pot multicomponent reaction for an

easy access of fully decorated 4*H*-pyran scaffolds and its rearrangement to 2-pyridone nucleus in aqueous media. *Tetrahedron Lett.* **2012**, *53*, 4687–4691.

(33) Kumar, B. S.; Srinivasulu, N.; Udupi, R. H.; Rajitha, B.; Reddy, Y. T.; Reddy, P. N.; Kumar, P. S. An efficient approach towards three component coupling of one pot reaction for synthesis of functionalized benzopyrans. *J. Heterocycl. Chem.* **2006**, *43*, 1691–1693.

(34) Elinson, M. N.; Ilovaisky, A. I.; Merkulova, V. M.; Zaimovskaya, T. A.; Nikishin, G. I. Electrocatalytic multicomponent assembling of aldehydes, *N*-alkyl barbiturates and malononitrile: an efficient approach to pyrano[2,3-d]pyrimidines. *Mendeleev Commun.* **2011**, *21*, 122–124.

(35) Wang, L. M.; Jiao, N.; Qiu, J.; Yu, J. J.; Liu, J. Q.; Guo, F. L.; Liu, Y. Sodium stearate-catalyzed multicomponent reactions for efficient synthesis of spirooxindoles in aqueous micellar media. *Tetrahedron* **2010**, *66*, 339–343.

(36) Raghuvanshi, D. S.; Singh, K. N. An efficient protocol for multicomponent synthesis of spirooxindoles employing *L*-proline as catalyst at room temperature. *J. Heterocycl. Chem.* **2010**, 47, 1323–1327.

(37) He, Y.; Guo, H. Y.; Tian, J. J. A simple three-component synthesis of spiro-pyran derivatives. J. Chem. Res. 2011, 528-530.