

Rhodium(III)-Cyclometalated-Imine Complexes: Solution Behavior and Reactivity with Molecular Hydrogen

Paolo Marcazzan, Brian O. Patrick, and Brian R. James*

Department of Chemistry, The University of British Columbia, 2036 Main Mall,
Vancouver, B.C. V6T 1Z1, Canada

Received November 12, 2004

Acetone solutions of *cis,trans,cis*-[Rh(H)₂(PPh₃)₂(acetone)₂]PF₆ and *cis*-[Rh(PPh₃)₂(acetone)₂]PF₆ react with the imines RN=CR'(R''), where R = PhCH₂, Ph, or C₆H₁₁; R' = H or Me; and R'' = Ph or *p*-OMe-C₆H₄, to form orthometalated complexes exemplified by [Rh(H){RN=CR'(o-C₆H₄)}(PPh₃)₂(acetone)]PF₆ that are generally characterized spectroscopically, and in one case (where R = PhCH₂, R' = Me, and R'' = *p*-OMe-C₆H₄) by X-ray crystallography. These complexes maintain their integrity in acetone solution and show no reaction toward 1 atm H₂ at ambient conditions. In MeOH, the complexes in which R' = Me simply change to the corresponding MeOH derivative, but when R' = H they are partially converted to the non-orthometalated, η^1 -N-imine species [Rh(PPh₃)₂(RN=CHR'')(MeOH)]PF₆ (**I**), and in MeOH, stoichiometric hydrogenation of the imine to the corresponding amine occurs via **I**. We have shown recently (*Inorg. Chem.* **2004**, *43*, 4820) that the catalyzed hydrogenation of PhCH₂N=C(H)Ph to dibenzylamine using *cis,trans,cis*-[Rh(H)₂(PPh₃)₂(MeOH)₂]PF₆ as catalyst precursor in MeOH proceeds via a cationic Rh-(η^1 -N-imine)(amine) intermediate (**II**), analogous to **I** but with the MeOH replaced by the amine that is formed via hydrolysis of the imine by the presence of adventitious water. Intermediate **I** (R = PhCH₂, R'' = Ph) is shown to be somewhat less active than **II** toward H₂, but both intermediates provide viable pathways for hydrogenation of the imine. The orthometalated imine-amine complex [Rh(H){PhCH₂N=C(Me)(o-C₆H₄)}(PPh₃)₂(NH₂CH₂Ph)]PF₆, where the benzylamine comes from imine hydrolysis, is also isolated and characterized crystallographically.

Introduction

Our recent studies on the homogeneous H₂-hydrogenation of the imine PhCH₂N=C(H)Ph catalyzed by the [Rh(COD)(PPh₃)₂]PF₆ (**1**) precursor (when, for example, imine:Rh = 100) have revealed that the active catalyst (the species that reacts with H₂) is a mixed Rh(I)-imine-amine species, the amine benzylamine being formed via an accompanying Rh-catalyzed hydrolysis of the imine by adventitious water.¹ Under stoichiometric conditions (imine:Rh = 1), we now observe quite different chemistry, which is reported here: the imine coordinates with orthometalation in an η^2 fashion via the imine-N and the ortho-C atom of a phenyl group, giving rise to a five-membered metallacycle within a hydrido-Rh(III)-o-metalated complex. Such intramolecular activation of a C–H bond of an aryl group of a donor ligand has long been known for platinum metal complexes of a variety of ligands, including N-donors,^{2–4} and there are many

early examples in Rh and Ir chemistry.^{5,6} Our group has recently reported on orthometalation within semicarbazone complexes of Rh (semicarbazone being a substituted imine)⁷ and within Ir species containing simple imines such as PhCH₂N=C(H)Ph,⁸ but to our knowledge this work is the first report of orthometalation of simple imines at Rh (specifically imines of the type RN=CR'(R''), where R = PhCH₂, Ph, or C₆H₁₁; R' = H or Me; and R'' = Ph or *p*-OMe-C₆H₄, containing solely alkyl or aryl substituents). Of more interest and relevance, however, are the solution chemistry of these new orthometalated species, their reaction with H₂, and the implications of the findings to catalyzed hydrogenation of the imines, a topic that continues to attract much interest.⁹

* To whom correspondence should be addressed. Tel: +1-(604)-822-6645. Fax: +1-(604)-822-2847. E-mail: brj@chem.ubc.ca.

(1) (a) Marcazzan, P.; Patrick, B. O.; James, B. R. *Organometallics* **2003**, *22*, 1177. (b) Marcazzan, P.; Abu-Gnim, C.; Seneviratne, K. N.; James, B. R. *Inorg. Chem.* **2004**, *43*, 4820. (c) Marcazzan, P.; Patrick, B. O.; James, B. R. *Inorg. Chem.* **2004**, *43*, 6838.

(2) (a) Parshall, G. W. *Acc. Chem. Res.* **1970**, *3*, 179. (b) Parshall, G. W. *Acc. Chem. Res.* **1975**, *8*, 113.

(3) (a) van Baar, J. F.; Vrieze, K.; Stufkens, D. J. *J. Organomet. Chem.* **1975**, *97*, 461. (b) van Baar, J. F.; Vrieze, K.; Stufkens, D. J. *J. Organomet. Chem.* **1975**, *85*, 249.

(4) Omae, I. *Chem. Rev.* **1979**, *79* (4), 287.

(5) Hughes, R. P. In *Comprehensive Organometallic Chemistry*; Wilkinson, G., Stone, F. G. A., Abel, E. W., Eds.; Pergamon Press: Oxford, 1982; Vol. 5, Chapter 34, p 392.

(6) Leigh, G. J.; Richards, R. L. In *Comprehensive Organometallic Chemistry*; Wilkinson, G., Stone, F. G. A., Abel, E. W., Eds.; Pergamon Press: Oxford, 1982; Vol. 5, Chapter 35, p 578.

(7) (a) Ezhova, M. B.; Patrick, B. O.; James, B. R.; Ford, M. E.; Waller, F. *J. Russ. Chem. Bull. Int. Ed.* **2003**, *52*, 2707 (M. Vol'pin Memorial Volume). (b) Ezhova, M. B.; Patrick, B. O.; Sereviratne, K. N.; James, B. R.; Waller, F. J.; Ford, M. E. *Inorg. Chem.*, in press.

(8) Marcazzan, P.; Patrick, B. O.; James, B. R. *Russ. Chem. Bull. Int. Ed.* **2003**, *52*, 2715 (M. Vol'pin Memorial Volume).

(9) (a) Kobayashi, S.; Ishitani, H. *Chem. Rev.* **1999**, *99*, 1069. (b) Tang, W.; Zhang, X. *Chem. Rev.* **2003**, *103*, 3029. (c) Blaser, H.-U.; Malan, C.; Pugin, B.; Spindler, F.; Steiner, H.; Studer, M. *Adv. Synth. Catal.* **2003**, *345*, 103. (d) Dorta, R.; Broggini, D.; Kissner, R.; Togni, A. *Chem. Eur. J.* **2004**, *10*, 4546.




Figure 1. Scheme showing reactions for formation of **4**.

Results and Discussion

Reaction of either *cis,trans,cis*- $[\text{Rh}(\text{H})_2(\text{PPh}_3)_2(\text{acetone})_2]\text{PF}_6$ (**2**) or *cis*- $[\text{Rh}(\text{PPh}_3)_2(\text{acetone})_2]\text{PF}_6$ (**3**)^{10,11} with $\text{PhCH}_2\text{N}=\text{C}(\text{H})\text{Ph}$ (Rh:imine = 1:1) in acetone at room temperature ($\sim 20^\circ\text{C}$) under Ar affords the Rh(III)-orthometalated complex $[\text{Rh}(\text{H})\{\text{PhCH}_2\text{N}=\text{CH}(\text{o-C}_6\text{H}_4)\}(\text{PPh}_3)_2(\text{acetone})]\text{PF}_6$ (**4**) (Figure 1), isolated as an off-white solid. Formation of orthometalated complexes via reaction of a simple imine with a Rh(I) species has not been reported in detail, although we have noted the existence of such species¹ and have reported on the Ir analogue of **4** that was isolated from reaction of the imine with the Ir analogue of **2**.⁸ The mechanism^{3,8} presumably involves displacement of an acetone ligand by imine when the *ortho*-CH group becomes close to the metal, this being accompanied, when **2** is the reactant, by reductive elimination of the hydride ligands as H_2 (detected in the ^1H NMR spectrum at δ_{H} 4.40 in acetone- d_6); oxidative addition of C–H then generates **4**. Detection of hydride (see below) versus deuteride in the product isolated from reaction of $\text{PhCH}_2\text{N}=\text{CHPh}$ with the precursor $[\text{Rh}(\text{D})_2(\text{PPh}_3)_2(\text{acetone-}d_6)_2]\text{PF}_6$ in acetone- d_6 is consistent with the hydrido ligand emanating from the orthometalation process. Of note, reaction of $[\text{RhCl}(\text{CO})_2]_2$ with $\text{MeN}=\text{C}(\text{H})\text{Ph}$ forms the η^1 -N-imine complex *cis*- $\text{RhCl}(\text{CO})_2(\text{MeN}=\text{CHPh})$, presumably because the presence of the electron-withdrawing CO ligands diminishes the tendency for oxidative addition.³

The IR data for **4** reveal bands for ν_{RhH} , $\nu_{\text{C=N}}$, and ν_{CO} for the coordinated acetone. The room-temperature $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum in acetone- d_6 shows a doublet (δ 40.40, $J_{\text{RhP}} = 116$), the J value being indicative of mutually *trans* phosphines.^{1c} The high-field hydride ^1H NMR resonance (δ -12.85 , $^2J_{\text{HP}} \approx J_{\text{RH}} = 13$) appears as a pseudo-quartet (ps q) instead of the expected doublet of triplets; such overlapping of triplets has been reported for other *trans*-Rh(PPh_3)₂-hydrido moieties.¹² Upfield-shifted δ_{H} resonances for the orthometalated ring protons and for the N=CH signal (δ_{H} 7.81 versus 8.50 for the free imine) are similar to those reported for the Ir analogue.⁸ Thus, in acetone solution, **4**

maintains the solid state structure shown in Figure 1 and confirmed crystallographically for a closely related imine complex (**5**, see below).

In CD_3OD , **4** shows different and more complex behavior (Figure 2) that was studied by variable-temperature NMR experiments (Figure S1 in the Supporting Information). The room-temperature $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **4** now indicates the presence of three species: $[\text{Rh}(\text{H})\{\text{PhCH}_2\text{N}=\text{CH}(\text{o-C}_6\text{H}_4)\}(\text{PPh}_3)_2(\text{CD}_3\text{OD})]\text{PF}_6$ (**4'**)¹³ (δ 40.52 d, $J_{\text{RhP}} = 116$), *cis*- $[\text{Rh}(\text{PPh}_3)_2(\text{CD}_3\text{OD})_2]\text{PF}_6$ (**3'**) (δ 57.02 d, $J_{\text{RhP}} = 207$),¹¹ and a fluxional species that gives rise to two broad, unresolved resonances centered at $\delta \sim 46$ and ~ 55 . These resonances resolve at 253 K into an AMX, 8-line pattern (δ 44.55 dd, $J_{\text{RhP}} = 164$, $^2J_{\text{PP}} = 54$; 54.74 dd, $J_{\text{RhP}} = 214$, $^2J_{\text{PP}} = 54$), indicative of a species containing two inequivalent *cis*-phosphines, each *trans* to a different ligand, and this is identified as the imino-solvento complex *cis*- $[\text{Rh}(\text{PPh}_3)_2(\text{PhCH}_2\text{N}=\text{CHPh})(\text{CD}_3\text{OD})]\text{PF}_6$ (**I**), shown in Figure 2. Free acetone (δ 2.10) is detected in the room-temperature ^1H NMR spectrum of **4** in CD_3OD (Figure S1), indicating its displacement by CD_3OD to give **4'**, while detection of free imine indicates that **4'** has undergone "reductive elimination" of this ligand, a premise confirmed by the presence of $[\text{Rh}(\text{PPh}_3)_2(\text{CD}_3\text{OD})_2]\text{PF}_6$ (**3'**) in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum (Figure S1) in an amount that corresponds to the amount of free imine. The room-temperature ^1H NMR spectrum also shows sharp singlets at δ 4.80 and 4.99 for the benzylic protons of free imine and the orthometalated imine (of **4'**), respectively, as well as a broad signal in the δ 4.7–5.3 region; at 253 K (Figure S1), this broad signal sharpens into a pair of doublets (δ 4.50 d, 5.35 d, $^2J_{\text{HH}} = 6$), showing diastereotopic inequivalence of the benzylic protons within **I**, presumably because of some restricted rotation of the imine moiety at lower temperatures. The $\text{CH}=\text{N}$ resonances of **4'** and **I** are not seen, presumably being masked by the aromatic proton signals. The room-temperature ^1H NMR spectrum also shows a downfield doublet (δ 9.65 d, 2H, $^3J_{\text{HH}} \approx 7$) that is not attributable to **3'** or to free imine; the corresponding ^1H – ^{13}C HETCOR NMR experiment reveals correlation of this signal with resonances corresponding to aromatic C atoms (δ 132), and so the signal is attributed to the *o*-protons of the imine-Ph moiety of **I** being involved in a π -arene interaction with Ph groups of the PPh_3 ligands that causes a shielding of this resonance. A similar shielding was observed for a Rh(I)-imine-amine complex recently reported by our group.^{1c} Neither **4'** nor **I** could be isolated pure from a solution mixture of **4'**, **3'**, and **I**, formed either from a solution of **4** in MeOH or from a MeOH solution of **4'** formed directly from reaction of *cis*- $[\text{Rh}(\text{PPh}_3)_2(\text{MeOH})_2]\text{PF}_6$ with 1 equiv of $\text{PhCH}_2\text{N}=\text{C}(\text{H})\text{Ph}$.

Thus in MeOH, the orthometalation process evident in **4** (and **4'**) is effectively reversed and affords **3'** and **I**, square planar species that can both undergo oxidative addition of H_2 (see below). Such behavior is relevant in that $[\text{Rh}(\text{COD})(\text{PPh}_3)_2]\text{PF}_6$ (**1**) is an effective precursor catalyst for hydrogenation of imines in MeOH solution but not in acetone. In the latter solvent, the integrity of the coordinatively saturated Rh(III) species **4** is

(10) (a) Schrock, R. R.; Osborn, J. A. *J. Am. Chem. Soc.* **1971**, *93*, 2397. (b) Haines, L. M.; Singleton, E. *J. Chem. Soc., Dalton* **1972**, 1891.

(11) Marcazzan, P.; Ezhova, M. B.; Patrick, B. O.; James, B. R. C. *R. Chim.* **2002**, *5*, 373 (J. A. Osborn Memorial Volume).

(12) (a) Al-Najjar, I. M.; El-Baih, F. E. M.; Abu-Loha, F. M.; Goma, Z. *Trans. Met. Chem.* **1994**, *19*, 325. (b) Ghedini, M.; Neve, F.; Lanfredi, A. M. M.; Uguzzoli, F. *Inorg. Chim. Acta* **1988**, *147*, 243. (c) Bhayat, I. I.; McWhinnie, W. R. *J. Organomet. Chem.* **1972**, *46*, 159. (d) Di Vaira, M.; Peruzzini, M.; Zanolini, F.; Stoppioni, P. *Inorg. Chim. Acta* **1983**, *69*, 37.

(13) A prime superscript (e.g., as in **4'**) refers to the $\text{CD}_3\text{OD}/\text{CH}_3\text{OH}$ analogue of the corresponding acetone/acetone- d_6 species (e.g., **4**).




Figure 2. Scheme showing the species generated in a CD_3OD solution of **4** (or **4'**).




Figure 3. Structural and ORTEP diagram of $[\text{Rh}(\text{H})\{\text{PhCH}_2\text{N}=\text{C}(\text{Me})(o\text{-C}_6\text{H}_3\text{-}p\text{-OMe})\}(\text{PPh}_3)_2(\text{acetone})]\text{PF}_6 \cdot \text{Me}_2\text{CO}$ (**5**) with 50% probability thermal ellipsoids.


maintained, and there is no reactivity toward H_2 . However, exposure of a MeOH solution of **4'** to 1 atm H_2 results in quantitative hydrogenation of the coordinated imine within 5 min to afford dibenzylamine and *cis,trans,cis*- $[\text{Rh}(\text{H})_2(\text{PPh}_3)_2(\text{CD}_3\text{OD})_2]\text{PF}_6$ (**2'**), as detected by $^{31}\text{P}\{^1\text{H}\}$ and ^1H NMR spectroscopies.^{10,11} Monitoring the H_2 -hydrogenation of **4** in MeOH (i.e. **4'**) by variable-temperature NMR spectroscopy showed no evidence for hydrido-imine intermediates. At 243 K, the spectra revealed the presence of **4'**, **2'** (formed from **3'**), **I**, free acetone, and a smaller amount of free imine; as the solution was warmed to 300 K, **I** and **4'** gradually disappeared with concomitant formation of dibenzylamine (δ_{CH_2} 4.80) and increased amounts of **2'** and **3'**. The dibenzylamine is, in fact, involved in a labile equilibrium with **3'**, and we have reported earlier on the NMR investigation of this equilibrium.^{1a} The only Rh species seen in the final solution at 300 K is **2'**, when all the imine has been converted to free amine.

Attempts to obtain a crystal of **4** suitable for X-ray analysis were unsuccessful, but the ketimine $\text{PhCH}_2\text{N}=\text{C}(\text{Me})(p\text{-MeOC}_6\text{H}_4)$ similarly underwent the orthometalation reaction with **2** in acetone to give $[\text{Rh}(\text{H})\{\text{PhCH}_2\text{N}=\text{C}(\text{Me})(o\text{-C}_6\text{H}_3\text{-}p\text{-OMe})\}(\text{PPh}_3)_2(\text{acetone})]\text{PF}_6$ (**5**), which was characterized crystallographically (Figure 3); selected structural parameters are given in Table 2. The complex crystallizes as a two-component twin with one acetone molecule in the asymmetric unit. The geometry at the Rh is essentially octahedral with *trans*-phosphines, the acetone being *trans* to the metalated *o*-carbon and the refined, located Rh-H hydrogen atom being *trans* to the imine-N atom. As for the related Ir complex $[\text{Ir}(\text{H})\{\text{PhCH}_2\text{N}=\text{CH}(o\text{-C}_6\text{H}_4)\}(\text{PPh}_3)_2(\text{acetone})]\text{PF}_6$, the large $\text{C}(53)\text{-O-Rh}$ angle of 137.9° and the $\text{P}(1)\text{-Rh-P}(2)$ angle of 166.8° indicate that the acetone-Me groups are subject to steric repulsions by the $\text{Rh}(\text{PPh}_3)_2$ moiety. The structure of **5** is generally very similar to that of the Ir complex noted above, with the

Table 1. Crystallographic Data for 5 and 9*

	5	9*
formula	$\text{C}_{58}\text{H}_{59}\text{NO}_3\text{F}_6\text{P}_3\text{Rh}$	$\text{C}_{59.50}\text{H}_{60}\text{N}_2\text{O}_{1.50}\text{F}_6\text{P}_3\text{Rh}$
fw	1127.88	1136.92
cryst color, habit	colorless, needle	red, irregular
cryst size (mm)	$0.30 \times 0.07 \times 0.07$	$0.40 \times 0.30 \times 0.20$
space group	$P2_1/c$	$P2_1/n$
<i>a</i> (Å)	10.060(1)	16.5613(5)
<i>b</i> (Å)	19.129(2)	17.8073(4)
<i>c</i> (Å)	27.912(3)	18.7853(5)
β (deg)	91.600(5)	107.822(2)
<i>V</i> (Å ³)	5369(1)	5274.2(2)
<i>Z</i>	4	4
<i>D</i> _{calc} (g cm ⁻³)	1.395	1.432
μ (mm ⁻¹)	0.473	0.481
total no. of reflns	157 997	48 378
no. of unique reflns	30 620	12 155
<i>R</i> _{int}	0.071	0.047
no. of variables	697	660
<i>R</i> ₁	0.044 (<i>I</i> > 2σ(<i>I</i>), 16 036 obs reflns)	0.032 (<i>I</i> > 3σ(<i>I</i>), 8601 obs reflns)
w <i>R</i> ₂	0.104 (all data)	0.090 (all data)
GOF	0.94 (all data)	1.12 (all data)

five-membered cyclometalated rings being essentially planar. The $\text{N}(1)\text{-C}(8)$ imine bond is 0.18 Å shorter than the $\text{N}(1)\text{-C}(10)$ single bond, while the Rh-O bond length of 2.19 Å likely reflects a *trans*-influence of the *ortho*-C ligand, although we have been unable to find any structural data on a Rh-acetone moiety (the Rh-O distances within related orthometalated Rh-semicarbazone complexes⁷ are in the range 2.207–2.310 Å). The solid state IR of **5** reveals the $\nu_{\text{Rh-H}}$, $\nu_{\text{C=N}}$, and $\nu_{\text{C=O}}$ bands. The room-temperature $^{31}\text{P}\{^1\text{H}\}$ and ^1H NMR spectra of **5** in acetone-*d*₆ correspond closely to those of **4**, with the former showing also the “extra” resonances for the imine Me and OMe protons. Interestingly, and in contrast with **4**, complex **5** upon dissolution in MeOH simply forms the MeOH -solvated analogue **5'** (δ_{P} 40.33 d, $J_{\text{RhP}} = 117.8$; $\delta_{\text{H}} -12.70$ ps q, $J_{\text{RhH}} \approx {}^2J_{\text{HP}} = 13$) and displays none of the behavior associated with the

**Figure 4.** Complexes 7–10.**Table 2.** Selected Bond Distances and Angles for [Rh(H){PhCH₂N=C(Me)(*o*-C₆H₃-*p*-OMe)}(PPh₃)₂-(acetone)]PF₆·Me₂CO (**5**) (estimated standard deviations in parentheses)

bond	length (Å)	bond	angle (deg)
Rh(1)–P(1)	2.3313(8)	P(1)–Rh(1)–C(2)	88.92(7)
Rh(1)–P(2)	2.3529(7)	P(2)–Rh(1)–N(1)	99.56(6)
Rh(1)–N(1)	2.1597(19)	P(2)–Rh(1)–O(2)	92.15(5)
Rh(1)–O(2)	2.1937(18)	P(2)–Rh(1)–H(1)	85.0(6)
Rh(1)–C(2)	2.001(2)	N(1)–Rh(1)–H(1)	171.4(7)
Rh(1)–H(1)	1.564(17)	N(1)–Rh(1)–C(2)	79.86(9)
N(1)–C(8)	1.296(3)	N(1)–Rh(1)–O(2)	92.11(8)
N(1)–C(10)	1.479(3)	N(1)–C(8)–C(1)	115.8(2)
O(2)–C(53)	1.223(3)	C(2)–C(1)–C(8)	117.0(2)
C(53)–C(54)	1.496(4)	P(2)–Rh(1)–C(2)	88.18(7)
C(53)–C(55)	1.498(4)	O(2)–C(53)–C(54)	122.6(3)
C(8)–C(1)	1.465(3)	O(2)–C(53)–C(55)	120.1(3)
C(8)–C(9)	1.521(3)	C(53)–O(2)–Rh(1)	137.93(19)
C(1)–C(2)	1.414(3)	Rh(1)–C(2)–C(1)	113.62(19)
C(10)–C(11)	1.527(3)	Rh(1)–C(2)–C(3)	128.52(19)
		Rh(1)–N(1)–C(8)	113.53(17)
bond	angle (deg)	Rh(1)–N(1)–C(10)	124.54(16)
P(1)–Rh(1)–P(2)	166.86(3)	C(53)–O(2)–Rh(1)	137.93(19)
O(2)–Rh(1)–C(2)	171.90(9)	Rh(1)–C(2)–C(1)	113.62(19)
P(1)–Rh(1)–N(1)	92.54(6)	Rh(1)–C(2)–C(3)	128.52(19)
P(1)–Rh(1)–O(2)	92.52(5)	Rh(1)–N(1)–C(8)	113.53(17)
P(1)–Rh(1)–H(1)	82.4(7)	Rh(1)–N(1)–C(10)	124.54(16)

chemistry of **4'** shown in Figure 2. Consistent with this, no reaction is observed when exposing a MeOH solution of **5** (or **5'**) to 1 atm H₂.

The orthometalated complex [Rh(H){CH₃N=CH(*o*-C₆H₄)}(PPh₃)₂(acetone)]PF₆ (**6**), formed from the imine MeN=C(H)Ph, was also isolated and characterized spectroscopically as for **4**, and its solution behavior is similar to that of **4**: in acetone, the solid state structure is maintained, while in CD₃OD, chemistry corresponding to that of Figure 2 is seen (i.e., formation of **6'**, the “reverse” of orthometalation, and reactivity toward H₂). Analogous orthometalated species [Rh(H){RN=CR'(*o*-C₆H₄)}(PPh₃)₂(acetone-*d*₆)]PF₆, **7**–**10**, were synthesized in situ in acetone-*d*₆ from the precursor imines PhN=C(H)Ph, C₆H₁₁N=C(H)Ph, PhCH₂N=C(Me)Ph, and PhCH₂N=CPh₂, respectively (Figure 4); their NMR spectra were assigned as for complexes **4**–**6**, and in the case of **7**, the high-field hydride ¹H resonance is seen as the expected doublet of triplets because of a sufficient difference in the *J*_{RhH} and ²*J*_{HP} values (16 and 13 Hz, respectively). Of key interest, species **9'** and **10'** (the MeOH derivatives of **9** and **10** that can be made in situ in MeOH from *cis*-[Rh(PPh₃)₂(MeOH)₂]⁺ and 1 equiv of the imine), like **5'** with the Me substituent at the imine-C atom, are essentially stable as such in MeOH (but see below for **9'**), while **7'** and **8'** (the MeOH analogues of **7** and **8**) that have the H atom at the

imine-C atom show the same complex behavior in MeOH as **4'** and **6'** (cf. Figure 2). Thus, the labile chemistry noted in MeOH is seen only for the imine systems that contain the N=CH moiety, but it is unclear why this governs the “reverse orthometalation”. There is no evidence for incorporation of deuterium into **4'**, **3'**, or **I** in the observed chemistry of Figure 2, and thus cleavage of the Rh–C bond by protons/deuterons is ruled out, implying that direct reductive elimination from the *cis*-Rh(H)–C moiety occurs (or alternatively the oxidative addition of the imine is less complete in MeOH). Possibly some H-bonding interaction between the MeOH and the H atom of the imine N=CH increases the acidity of the H atom, decreases the basicity of the imine, and thus decreases the propensity for oxidative addition to the Rh(I). A contributing factor is also likely to be the somewhat higher donor number of MeOH versus that of acetone (20 versus 17),¹⁴ a feature that has been demonstrated within the related *cis,trans,cis*-[Ir(H)₂-(PPh₃)₂(solvent)₂][BF₄] species.¹⁵

The labile chemistry noted in MeOH is also reflected in the catalytic hydrogenation of the different imines in this solvent. As noted above, hydrogenation of **4'** in MeOH gives dibenzylamine and *cis,trans,cis*-[Rh(H)₂-(PPh₃)₂(MeOH)₂][PF₆] (**2'**); this solution can then be treated with a further equivalent of PhCH₂N=CHPh, and under H₂ the imine is again reduced to the amine via the partial formation of **4'** and the chemistry of Figure 2. Thus, via a series of stoichiometric hydrogenations, the net process is a catalyzed imine hydrogenation via [Rh(PPh₃)₂(PhCH₂N=CHPh)(MeOH)]PF₆ (**I**). Of note, if excess imine is used under the same conditions in MeOH, catalytic hydrogenation occurs via [Rh(PPh₃)₂-(PhCH₂N=CHPh)(NH₂CH₂Ph)]PF₆ (**II**), the amine being formed via a Rh-catalyzed hydrolysis of the imine by adventitious water.¹ The rate (turnover)-determining step of the catalytic process is oxidative addition of H₂ to **II** with a rate constant that we will define as *k*_{II}, whose value at 30 °C has been determined^{1b} as ~45 M⁻¹ s⁻¹. An estimate of the corresponding *k*_I value for oxidative addition of H₂ to **I** at ~20 °C can be made as follows. Integration of the ³¹P{¹H} resonances of the species present in a solution of **4'** at this temperature ([Rh]_{total} = 0.022 M) shows a 4:1 ratio of **I**:**4'** and much smaller amounts of **3'** (see Figures 2 and S1), and thus the **I** is ~0.017 M; when the solution is shaken under 1 atm H₂ (which is effectively maintained at a constant solution concentration of ~1.5 × 10⁻³ M^{1b}), the imine is converted to amine with *t*_{1/2} ≈ 90 s, corresponding to a *k*_I value of ~8 M⁻¹ s⁻¹ at room temperature. Although the kinetic activation parameters have not been measured for these systems, it is almost certain that reactivity to give amine via **II** is more favorable than via **I** (probably by a factor of ~3, as rate constants for oxidative addition of H₂ to such systems typically double with a 10 °C increase in temperature).¹⁶ The difference in oxidative addition reactivity (see Figure 5) seems reasonable considering the higher basicity of benzylamine versus that of CD₃OD/MeOH. An important

(14) Huheey, J. E. *Inorganic Chemistry*, 3rd ed.; Harper Collins: New York, 1983; p 340.

(15) Crabtree, R. H.; Mellea, M. F.; Mihelcic, J. M.; Quirk, J. M. *J. Am. Chem. Soc.* **1982**, *104*, 107.

(16) James, B. R. *Homogeneous Hydrogenation*; Wiley: New York, 1973; Chapter XII.




Figure 5. Oxidative addition of H₂ to species **I** and **II**. The supposed dihydrido products have not been detected (see ref 1b).

result from this study is the demonstration that Rh-catalyzed imine hydrogenation can, in principle, occur *without* the need for hydrolysis of imine to amine. Under catalytic conditions in MeOH,¹ the ketimine PhCH₂N=C(Me)Ph was hydrogenated much more slowly than the aldimine PhCH₂N=C(H)Ph, with conversions of only ~4 and ~40% after 1 and 40 h, respectively. Ketimines are invariably hydrogenated more slowly than a corresponding aldimine because of increased steric bulk around the N=C moiety,¹⁷ but whether the hydrogenation of PhCH₂N=C(Me)Ph occurs via a solvent- η^1 -imine intermediate such as **I** or an η^1 -amine- η^1 -imine species such as **II** is unclear, particularly as an orthometalated species (**9***) containing also η^1 -benzylamine has been isolated (see below). With the more heavily substituted imine PhCH₂N=CPh₂, no catalytic hydrogenation is observed under ambient conditions in MeOH, while we have reported earlier^{1a} that the corresponding hydrogenation of PhN=C(H)Ph is inhibited by catalyst poisoning, the amine product PhNHCH₂Ph coordinating to the Rh via the N-phenyl group (η^4 in the solid state and η^6 in solution).

When the MeOH solution of [Rh(H){PhCH₂N=CMe-(*o*-C₆H₄)}(PPh₃)₂(acetone)]PF₆ (**9'**) was left at room temperature for about a week "under vacuo", a small crystal formed, and this proved by X-ray analysis (Figure 6) to be [Rh(H){PhCH₂N=C(Me)(*o*-C₆H₄)}(PPh₃)₂(NH₂CH₂Ph)]PF₆ (**9***), where the MeOH ligand of **9'** has been replaced by benzylamine, the amine hydrolysis product of the imine. We have reported elsewhere on this metal-catalyzed hydrolysis reaction,^{1,8} and indeed in the Rh-catalyzed hydrogenation of PhCH₂N=CHPh the active catalyst is **II**, as noted above. Whether **9*** is a catalyst precursor for hydrogenation of PhCH₂N=C(Me)Ph will remain unknown until a reliable synthetic method is realized for this complex. The structure of **9*** (Table 3) is similar to that of complex **5** but where the acetone has been replaced by the amine; the key bond lengths and angles are close to those in **5**, and even the Rh–N(2)_{amine} distance is the same as the Rh–O distance in **5**. We have noted recently the close correspondence in the structures of the Ir complexes [Ir(H){PhCH₂N=CH(*o*-C₆H₄)}(PPh₃)₂(L)]PF₆, where L = acetone or benzylamine.⁸




Figure 6. ORTEP diagram of the cation [Rh(H){PhCH₂N=C(Me)(*o*-C₆H₄)}(PPh₃)₂(NH₂CH₂Ph)]⁺ of complex **9*** with 50% probability thermal ellipsoids.

Table 3. Selected Bond Distances and Angles for [Rh(H){PhCH₂N=C(Me)(*o*-C₆H₄)}(PPh₃)₂(NH₂CH₂Ph)]PF₆ (**9***) (estimated standard deviations in parentheses)

bond	length (Å)	bond	angle (deg)
Rh(1)–P(1)	2.3499(6)	P(2)–Rh(1)–N(1)	92.29(7)
Rh(1)–P(2)	2.3344(6)	P(2)–Rh(1)–N(2)	91.49(5)
Rh(1)–N(1)	2.224(2)	P(1)–Rh(1)–C(18)	88.50(6)
Rh(1)–N(2)	2.162(2)	P(2)–Rh(1)–C(18)	86.79(6)
Rh(1)–C(18)	1.999(2)	N(2)–Rh(1)–H(54)	175(1)
Rh(1)–H(54)	1.43(2)	N(2)–Rh(1)–N(1)	101.70(8)
N(2)–C(15)	1.288(3)	N(2)–Rh(1)–C(18)	79.45(8)
N(2)–C(8)	1.478(3)	N(2)–C(15)–C(16)	124.9(2)
N(1)–C(1)	1.509(4)	N(2)–C(15)–C(17)	116.0(2)
C(1)–C(2)	1.509(4)	C(15)–C(17)–C(18)	116.4(2)
C(15)–C(16)	1.511(3)	C(8)–N(2)–C(15)	121.0(2)
C(15)–C(17)	1.467(3)	N(2)–C(8)–C(9)	115.1(2)
C(8)–C(9)	1.512(3)	N(1)–C(1)–C(2)	114.3(2)
C(17)–C(18)	1.411(3)	Rh(1)–C(18)–C(17)	114.3(2)
bond	angle (deg)	Rh(1)–C(18)–C(19)	128.0(2)
P(1)–Rh(1)–P(2)	169.30(2)	Rh(1)–N(2)–C(15)	113.8(1)
N(1)–Rh(1)–C(18)	178.55(9)	C(8)–N(2)–C(15)	121.0(2)
P(1)–Rh(1)–N(1)	92.22(7)	N(2)–C(8)–C(9)	115.1(2)
P(1)–Rh(1)–N(2)	97.08(5)	N(1)–C(1)–C(2)	114.3(2)
P(1)–Rh(1)–H(54)	85(1)	Rh(1)–C(18)–C(17)	114.3(2)
P(2)–Rh(1)–H(54)	86(1)	Rh(1)–C(18)–C(19)	128.0(2)
		Rh(1)–N(2)–C(15)	113.8(1)

Experimental Section

Unless stated otherwise, synthetic procedures were performed at room temperature (~20 °C) using standard Schlenk techniques under an atmosphere of dry Ar. The liquid imines PhCH₂N=CHPh and CH₃N=CHPh from Aldrich were purified by distillation; the solid imines PhCH₂N=C(Me)Ph, PhCH₂N=C(Me)-*p*-C₆H₄OMe, PhCH₂N=CPh₂, and PhN=CHPh, and the liquid C₆H₁₁N=CHPh were synthesized in this laboratory previously by Drs. D. E. Fogg, K. S. MacFarlane, and M. B. Ezhova.¹⁸ The [Rh(COD)(PPh₃)₂]PF₆ (**1**) and corresponding *cis*-, *trans*-, *cis*-[Rh(H)₂(PPh₃)₂(solvent)₂]PF₆ (**2**, **2'**) and *cis*-[Rh(PPh₃)₂(solvent)₂]PF₆ (**3**, **3'**) precursors (solvent = acetone, MeOH) were prepared according to literature procedures.^{9,10} Other reagents were purchased from commercial sources and used as supplied. Solvents were dried over the appropriate agents and distilled under N₂ prior to use. NMR spectra were recorded on Bruker AV 300 (300 MHz for ¹H, 121 MHz for ³¹P{¹H}), 75 MHz for ¹³C) and AV 400 (400 MHz for ¹H, 162 MHz for ³¹P-

(18) (a) Fogg, D. E. Ph.D. Dissertation, The University of British Columbia, Vancouver, 1994. (b) MacFarlane, K. S.; Ezhova, M. B. Unpublished results.

{¹H}, 100 MHz for ¹³C) spectrometers. Residual solvent proton (¹H, relative to external SiMe₄ δ 0.00) and external P(OMe)₃ (³¹P{¹H}, δ 141.00 versus external 85% aqueous H₃PO₄) were used as references. All *J* values are given in Hz. Infrared spectra were recorded on an ATLI Mattson Genesis Series FTIR spectrophotometer, and IR bands (KBr pellet) are reported in cm⁻¹. Elemental analyses were performed by Mr. P. Borda of this department using a Carlo Erba 1108 analyzer.

[*N*,*o*-C-(*N*-Benzylidenebenzylaminato)]hydrido(acetone)bis(triphenylphosphine)rhodium(III) Hexafluorophosphate, [Rh(H){PhCH₂N=CH(*o*-C₆H₄)}(PPh₃)₂(acetone)]PF₆ (4**).** To a pale yellow solution of [Rh(H)₂(PPh₃)₂(acetone)]PF₆ (**2**) (0.113 mmol) or a red solution of [Rh(PPh₃)₂(acetone)]PF₆ (**3**) (0.116 mmol) in acetone (3 mL) was added PhCH₂N=CHPh (22.0 μL, 0.116 mmol) under Ar and the mixture stirred for 2 h. The volume was then reduced under vacuum to ~1 mL to afford spontaneous precipitation of an off-white solid that was collected by filtration, washed with Et₂O (3 × 4 mL), and dried in vacuo. Yield: 0.090 g (75%). ³¹P{¹H} NMR (acetone-*d*₆): δ 40.40 (d, *J*_{RhP} = 116). ¹H NMR (acetone-*d*₆): δ -12.85 (ps q, 1H, Rh-*H*, *J*_{RhH} ≈ ²*J*_{HP} = 13), 2.10 (s, 6H, free CH₃COCH₃), 5.25 (s, 2H, CH₂), 6.52 (t, 1H, *p*-(*o*-C₆H₄), ³*J*_{HH} = 6.5), 6.72 (d, 1H, *o*-(*o*-C₆H₄), ³*J*_{HH} = 6.5), 6.95 (m, 2H, *m*-(*o*-C₆H₄), ³*J*_{HH} = 6.5), 7.01–7.79 (m, 35H, aromatics), 7.81 (s, 1H, =CH). IR (KBr): ν 2096 (Rh–H), 1660 (C=O), 1611, 1576 (C=N). Anal. Calcd for C₅₃H₄₉NOF₆P₃Rh: C, 62.06; H, 4.81; N, 1.37. Found: C, 62.0; H, 4.9; N, 1.5.

[*N*,*o*-C-(*N*-*p*-MeO-Phenylmethylbenzylaminato)]-hydrido(acetone)bis(triphenylphosphine)rhodium(III) Hexafluorophosphate, [Rh(H){PhCH₂N=C(Me)(*o*-C₆H₃-*p*-OMe)}(PPh₃)₂(acetone)]PF₆ (5**).** This complex was prepared as a light brown solid from a solution of [Rh(H)₂(PPh₃)₂(acetone)]PF₆ (**2**) (0.100 mmol) and the imine PhCH₂N=C(Me)C₆H₄-*p*-OMe (0.024 g, 0.100 mmol) according to the procedure given for **4**. Yield: 0.070 g (65%). ³¹P{¹H} NMR (acetone-*d*₆): δ 40.43 (d, *J*_{RhP} = 117). ¹H NMR (acetone-*d*₆): δ -12.51 (ps q, 1H, Rh-*H*, *J*_{RhH} ≈ ²*J*_{HP} = 13), 1.95 (s, 3H, C(CH₃)), 2.10 (s, 6H, free CH₃COCH₃), 2.99 (s, 3H, *p*-OCH₃), 5.40 (s, 2H, CH₂), 6.08 (s, 1H, *m*-(*o*-C₆H₃-), 6.60 (d, 1H, *o*-(*o*-C₆H₃-), ³*J*_{HH} = 7), 6.85 (d, 1H, *m*-(*o*-C₆H₃-), ³*J*_{HH} = 7), 6.95–7.70 (m, 35H, aromatics). IR (KBr): ν 2105 (Rh–H), 1670 (C=O), 1600, 1576 (C=N). Anal. Calcd for C₅₅H₅₃NO₂F₆P₃Rh: C, 61.75; H, 4.99; N, 1.31. Found: C, 61.35; H, 5.3; N, 1.7. An X-ray quality crystal of **5** was obtained by slow evaporation of an acetone/Et₂O solution (1:1 vol) of the compound.

[*N*,*o*-C-(*N*-Benzylidenemethylaminato)]hydrido(acetone)bis(triphenylphosphine)rhodium(III) Hexafluorophosphate, [Rh(H){CH₂N=CH(*o*-C₆H₄)}(PPh₃)₂(acetone)]PF₆ (6**).** To a solution of [Rh(PPh₃)₂(acetone)]PF₆ (**3**) (0.096 mmol) in acetone (3 mL) was added CH₂N=CHPh (12.0 μL, 0.097 mmol) under Ar and the mixture stirred for 2 h. The volume was then reduced to ~1 mL followed by addition of Et₂O (4 mL) to precipitate a creamy-white solid that was collected by filtration, washed with Et₂O (3 × 3 mL), and dried in vacuo. Yield: 0.065 g (70%). ³¹P{¹H} NMR (acetone-*d*₆): δ 42.10 (d, *J*_{RhP} = 116). ¹H NMR (acetone-*d*₆): δ -12.58 (ps q, 1H, Rh-*H*, *J*_{RhH} ≈ ²*J*_{HP} = 13), 2.10 (s, 6H, free CH₃COCH₃), 3.79 (s, 3H, CH₃N), 6.54 (t, 1H, *p*-(*o*-C₆H₄), ³*J*_{HH} = 6.5), 6.83–7.00 (m, 3H, *m*,*o*-(*o*-C₆H₄), ³*J*_{HH} = 6.5), 7.15–7.70 (m, 30H, aromatics), 7.74 (s, 1H, =CH). IR (KBr): ν 2101 (Rh–H), 1666 (C=O), 1618, 1578 (C=N). Anal. Calcd for C₄₇H₄₅NO F₆P₃Rh: C, 59.44; H, 4.78; N, 1.47. Found: C, 59.1; H, 4.9; N, 1.5.

In Situ Characterization of [Rh(H){RN=CR'(*o*-C₆H₄)}(PPh₃)₂(acetone-*d*₆)]PF₆. An NMR tube equipped with an airtight J Young PTFE valve was charged with the precursor *cis*-[Rh(PPh₃)₂(acetone-*d*₆)]PF₆ (~0.01 mmol) and a 1:1 mol equiv of the respective imine, and the solvent was added via vacuum-transfer. The resulting red solution was then analyzed by NMR spectroscopy. **R = Ph, R' = H (7):** ³¹P{¹H} NMR (acetone-*d*₆): δ 39.91 (d, *J*_{RhP} = 116). ¹H NMR (acetone-*d*₆): δ -13.12 (dt, 1H, Rh-*H*, *J*_{RhH} = 16, ²*J*_{HP} = 13), 6.40 (t, 1H, *p*-(*o*-

C₆H₄), ³*J*_{HH} = 7), 6.65 (d, 1H, *o*-(*o*-C₆H₄), ³*J*_{HH} = 7), 6.80 (m, 2H, *m*-(*o*-C₆H₄), 7.20–7.70 (m, 35H, aromatics), 8.00 (s, 1H, =CH). **R = C₆H₁₁, R' = H (8):** ³¹P{¹H} NMR (acetone-*d*₆): δ 38.40 (d, *J*_{RhP} = 117). ¹H NMR (acetone-*d*₆): δ -12.82 (ps q, 1H, Rh-*H*, *J*_{RhH} ≈ ²*J*_{HP} = 13), 1.20–1.90 (m, 10H, C₆(H)₁₀), 4.30 (pt, 1H, C₆(H)₁₀, ³*J*_{HH} = 10), 6.45 (t, 1H, *p*-(*o*-C₆H₄), ³*J*_{HH} = 7), 6.52 (d, 1H, *o*-(*o*-C₆H₄), ³*J*_{HH} = 7), 6.95 (t, 1H, *m*-(*o*-C₆H₄), ³*J*_{HH} = 7), 7.05–7.90 (m, 31H, *m*-(*o*-C₆H₄) and aromatics), 8.10 (s, 1H, =CH). **R = PhCH₂, R' = Me (9):** ³¹P{¹H} NMR (acetone-*d*₆): δ 40.33 (d, *J*_{RhP} = 117). ¹H NMR (acetone-*d*₆): δ -12.42 (ps q, 1H, Rh(*H*), *J*_{RhH} ≈ ²*J*_{HP} = 13), 1.95 (s, 3H, C(CH₃)), 5.45 (s, 2H, CH₂), 6.50 (t, 1H, *p*-(*o*-C₆H₄), ³*J*_{HH} = 7), 6.65 (d, 1H, *o*-(*o*-C₆H₄), ³*J*_{HH} = 7), 6.85 (m, 2H, *m*-(*o*-C₆H₄), ³*J*_{HH} = 7), 6.90–7.80 (m, 35H, aromatics). **9'**, the CD₃OD derivative of **9**, was made in situ in CD₃OD from *cis*-[Rh(PPh₃)₂(CD₃-OD)]⁺ and 1 equiv of PhCH₂N=C(Me)Ph (see text); after one week, a crystal that deposited from the solution was subjected to X-ray analysis and determined to have the structure [Rh-(H){PhCH₂N=C(Me)(*o*-C₆H₄)}(PPh₃)₂(NH₂CH₂Ph)]PF₆ (**9'**). There was insufficient material for further characterization. **R = PhCH₂, R' = Ph (10):** ³¹P{¹H} NMR (acetone-*d*₆): δ 38.54 (d, *J*_{RhP} = 118). ¹H NMR (acetone-*d*₆): δ -12.61 (ps q, 1H, Rh-*H*), *J*_{RhH} ≈ ²*J*_{HP} = 13), 5.13 (s, 2H, CH₂), 5.94 (d, 1H, *o*-(*o*-C₆H₄), ³*J*_{HH} = 7), 6.40 (t, 1H, *p*-(*o*-C₆H₄), ³*J*_{HH} = 7), 6.45 (m, 2H, *m*-(*o*-C₆H₄), ³*J*_{HH} = 7), 7.00–7.75 (m, 40H, aromatics).

Crystal Structure Determinations. Measurements were made at 173(0.2) K on a Bruker X8 APEX and on a Rigaku/ADSC CCD area detector diffractometer for **5** and **9'**, respectively, with graphite-monochromated Mo Kα radiation (0.71073 Å). Some crystallographic data for **5** and **9'** are shown in Table 1.

Data for **5** were collected and integrated using the Bruker SAINT¹⁹ software package and corrected for absorption effects using the multiscan technique (SADABS²⁰). The material crystallizes as a two-component twin, with the two components related by a 2.2° rotation about the (1,0,0) axis. The structure was solved by direct methods²¹ using an *hklf4* format data set including corrected twin overlaps, and final refinements were carried out using an *hklf5* format data set containing all reflections for both twin components. The ratio of both twin components is roughly 1:1. All non-hydrogen atoms were refined anisotropically. Four F atoms of the PF₆ anion were disordered and were modeled in two orientations. The Rh-H hydrogen atom was located in a difference map and refined isotropically, while all other hydrogen atoms were included in calculated positions but not refined. The material crystallizes with one molecule of acetone in the asymmetric unit. The final cycle of full-matrix least-squares refinement on *F*² was based on 30 620 reflections and 697 variable parameters (least-squares function minimized Σ*w*(*F*_o² - *F*_c²)², where *w* = 1/[σ²(*F*_o²) + (0.0359*P*)² + 0.00*P*] and *P* = (*F*_o² + 2*F*_c²)/3), and converged to R1 = 0.094, wR2 = 0.104, GOF = 0.94.

Data for **9'** were collected and processed using the d*TREK²² program. The structure was again solved by direct methods²¹ and expanded using Fourier techniques.²³ The material crystallizes with 1.5 molecules of MeOH in the asymmetric unit. All solvent atoms were refined isotropically, while the rest were refined anisotropically. The N-H and Rh-H hydrogen atoms were refined isotropically; the rest were included in fixed positions. Neutral atom scattering factors were taken from

(19) SAINT: Version 6.02; Bruker AXS Inc.: Madison, WI, 1999.

(20) SADABS: Bruker Nonius area detector scaling and absorption correction, Version 2.05; Bruker AXS Inc.: Madison, WI.

(21) SIR97: Altomare, A.; Burla, M. C.; Cammelli, G.; Cascarano, M.; Giacovazzo, C.; Guagliardi, A.; Moliterni, A. G. G.; Polidori, G.; Spagna, A. *J. Appl. Cryst.* **1999**, *32*, 115.

(22) d*TREK: Area Detector Software, version 4.13; Molecular Structure Corporation: The Woodlands, TX, 1996–1998.

(23) DIRDIF94: Beurskens, P. T.; Admiraal, G.; Beurskens, G.; Bosman, W. P.; de Gelder, R.; Israel, R.; Smits, J. M. M. The DIRDIF-94 program system. Technical Report of the Crystallography Laboratory; University of Nijmegen, The Netherlands, 1994.

Cromer and Waber,²⁴ and anomalous dispersion effects were included in F_{calc} ,²⁵ the values of $\Delta f'$ and $\Delta f''$ being those of Creagh and McAuley.²⁶ Values for the mass attenuation coefficients are those of Creagh and Hubbell,²⁷ and all calculations were performed using the teXsan program.²⁸ The final cycle of full-matrix least-squares refinement on F^2 was based on 11 765 observed reflections ($I > 0.00\sigma(I)$) and 660 variable parameters (using $\sum w(F_o^2 - F_c^2)^2$, where $w = 1/\sigma^2(F_o^2)$) and converged to $R1 = 0.056$, $wR2 = 0.090$, $GOF = 1.12$.

Conclusions

Simple imines of the type $\text{RN}=\text{C}(\text{H})\text{R}''$ or $\text{RN}=\text{C}(\text{R}')\text{R}''$, where R, R', and R'' are solely alkyl or aryl substituents, react in a 1:1 stoichiometry with cationic Rh(I)-(PPh₃)₂ species to form (via the usual $\text{C}_{\text{ortho}}\text{--H}$ activation) orthometalated complexes containing a five-membered ring, when a suitable phenyl substituent (R' or R'') is available at the imine-C atom. The complexes

(24) Cromer, D. T.; Waber, J. T. *International Tables for X-ray Crystallography*, Vol. IV; The Kynoch Press: Birmingham, England, Table 2.2 A, 1974.

(25) Ibers, J. A.; Hamilton, W. C. *Acta Crystallogr.* **1964**, *17*, 781.

(26) Creagh, D. C.; McAuley, W. J. In *International Tables for Crystallography*, Vol. C; Wilson, A. J. C., Ed.; Kluwer Academic Publishers: Boston, 1992; Table 4.2.6.8, pp 219–222.

(27) Creagh, D. C.; Hubbell, J. H. In *International Tables for Crystallography*, Vol. C; Wilson, A. J. C., Ed.; Kluwer Academic Publishers: Boston, 1992; Table 4.2.4.3, pp 200–206.

(28) *teXsan*: Crystal Structure Analysis Package; Molecular Structure Corporation: The Woodlands, TX, 1985 and 1992.

are stable as such in acetone solution (and show no reactivity toward 1 atm H₂), but in MeOH solution those that contain a H atom at the imine-C are converted partially to a cationic, mixed $\eta^1\text{-N}$ -imine/MeOH species that provides a pathway for reaction of the orthometalated complexes with H₂; this reaction generates amine in a stoichiometric reaction and is relevant in the Rh-catalyzed hydrogenation of the imines, where we have previously reported that a cationic, mixed $\eta^1\text{-N}$ -imine/amine species is an important catalytic intermediate. Characterized also is a cationic, Rh-orthometalated imine complex containing also an amine ligand, the amine being formed by hydrolysis of the imine. The studies derive from our ongoing efforts to gain a more detailed understanding of catalyzed imine hydrogenations, particularly on the role of the solvent.

Acknowledgment. We thank the Natural Sciences and Engineering Research Council of Canada (NSERC) for funding.

Supporting Information Available: ³¹P{¹H} and ¹H spectra of **4** in CD₃OD at 298 and 253 K (Figure S1), and X-ray crystallographic data for the structures of **5** and **9*** in CIF format. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM049121O