

Dalton Communications

Reaction of a Side-on Co-ordinated Carbene Ligand with the Wittig Reagent

Andrew J. Edwards, Jochem U. Köhler, Jack Lewis* and Paul R. Raithby

Department of Chemistry, Lensfield Road, Cambridge CB2 1EW, UK


The 1-methylene-4-phenylcyclohexadiene triosmium complex $[\text{Os}_3(\text{CO})_9(\mu_3\text{-}\eta^2\text{:}\eta^2\text{:}\eta^2\text{-H}_2\text{C}=\text{C}_6\text{H}_5\text{Ph})]$ has been prepared from the reaction of $[\text{Os}_3(\text{CO})_9(\mu_3\text{-}\eta^2\text{:}\eta^1\text{:}\eta^2\text{-C}_6\text{H}_5\text{Ph})]$, which contains a side-on co-ordinated carbene group, with the Wittig reagent $\text{H}_2\text{C}=\text{PPh}_3$; the crystal and molecular structure of the product has been determined by an X-ray diffraction study.

Arenes can adopt a $\mu_3\text{-}\eta^2\text{:}\eta^2\text{:}\eta^2\text{-face-capping}$ mode when they are co-ordinated to a triangular metal cluster core, and this mode has been observed in a variety of cluster compounds during the last few years.^{1,2} These compounds are of interest as model complexes for the adsorption of arenes on metal surfaces.³

During our investigations of the reactivity of the triosmium cluster $[\text{Os}_3(\text{CO})_9(\mu_3\text{-}\eta^2\text{:}\eta^2\text{:}\eta^2\text{-C}_6\text{H}_6)]$ **1**^{4,5} we recently synthesised a triosmium cluster complex **2** containing a carbene in an unusual side-on co-ordination mode (Scheme 1).⁶ The cyclohexadienone triosmium complex **3** is formed during the purification of **2** on TLC plates. Complex **3** can also be obtained easily from **2** in the presence of air and a Lewis base (e.g. NEt_3). These reactions give a first hint that carbene carbon may act as an electrophilic centre which is attacked by Lewis bases (NEt_3 or surface groups bearing free electron pairs or even water in the case of silica gel).⁶

We were interested in the reactivity of the carbene complex **2** towards nucleophilic reagents and to compare these results with those obtained in the case of mononuclear Fischer carbene complexes in which the carbene is co-ordinated in the usual end-on mode. Our first comparative reaction involved the Wittig reagent, $\text{H}_2\text{C}=\text{PPh}_3$. For example, this compound reacts readily with the mononuclear tungsten complex $[(\text{OC})_5\text{W}=\text{CPh}(\text{OMe})]$ to form two products. One product is $[\text{W}(\text{CO})_5(\text{PPh}_3)]$, which is simply formed by co-ordination of the PPh_3 group to the co-ordinatively unsaturated ' $\text{W}(\text{CO})_5$ ' fragment. The second product, $\text{H}_2\text{C}=\text{CPh}(\text{OMe})$, in which a double $\text{C}=\text{C}$ bond is formed between the carbene carbon and the methylene moiety of the Wittig reagent,⁷ may be viewed as a dimerisation of two carbenes.


In the comparative reaction with the novel triosmium carbene complex **2** a solution of $\text{H}_2\text{C}=\text{PPh}_3$ (0.07 mmol) in tetrahydrofuran (thf) (5 cm³) at -78°C was added to the orange solution of **2** (0.05 mmol). During the addition the solution became yellow. The reaction mixture was purified by TLC. Product **4** (Scheme 2) was obtained as orange-yellow microcrystals in 50% yield and initially characterised from spectroscopic data.[†] Its molecular and crystal structure has been established by X-ray crystallography.[‡] The asymmetric unit contains two independent but structurally similar molecules which are separated by normal van der Waals



Scheme 1 Synthesis of complexes **2** and **3**. (i) PhLi , thf, -78°C ; $[\text{N}(\text{PPh}_3)_2]\text{Cl}$, Pr^iOH ; (ii) $[\text{CPh}_3]\text{BF}_4$, CH_2Cl_2 , -78°C

[†] IR (CH_2Cl_2): $\nu(\text{CO})$ 2077m, 2033vs, 2024vs, 2002m, 1976m and 1950w cm^{-1} . NMR (CD_2Cl_2): ^1H , δ 7.3–7.6 (m, 5 H), 4.84 [t, $J(\text{HH}) = 2.5, 1 \text{ H}$], 3.51 [dd, $J(\text{HH}) = 8.0, 2.5, 2 \text{ H}$], 3.20 (s, 2 H) and 2.67 [d, $J(\text{HH}) = 8.0 \text{ Hz}, 2 \text{ H}$]. Positive-ion FAB mass spectrum: m/z 996 (M^+ , calc. 996).

[‡] Crystal data: $\text{C}_{22}\text{H}_{12}\text{O}_9\text{Os}_3$, $M = 990.92$, monoclinic, space group $P2_1/c$ (no. 14), $a = 18.811(4)$, $b = 14.155(3)$, $c = 17.463(3) \text{ \AA}$, $\beta = 95.21(3)^\circ$, $U = 4631(2) \text{ \AA}^3$, $Z = 8$, $D_c = 2.843 \text{ Mg m}^{-3}$, $\lambda(\text{Mo-K}\alpha) = 0.71073 \text{ \AA}$, $\mu(\text{Mo-K}\alpha) = 16.473 \text{ mm}^{-1}$, $F(000) = 3552$, crystal dimensions $0.25 \times 0.33 \times 0.38 \text{ mm}$. Data collection at room temperature on a Stoe four-circle diffractometer. Structure solved by direct methods and subsequent Fourier-difference syntheses. Hydrogen atoms were placed in idealised positions and allowed to ride on the relevant carbon atom. Refinement by full-matrix least squares on F_o^2 using SHELXTL PLUS⁸ and SHELXL 93⁹ to give $R1 = 0.079$ [$F > 4\sigma(F)$] and $wR2 = 0.237$ (all data), where $R1 = \Sigma||F_o| - |F_c||/\Sigma|F_o|$, $wR2 = \{\Sigma w(F_o^2 - F_c^2)^2/\Sigma[w(F_o^2)^2]\}^{1/2}$ and $w = 1/[\sigma^2(F_o^2) + (0.153P)^2 + 149.34P]$ where $P = (F_o^2 + 2F_c^2)/3$. Atomic coordinates, thermal parameters and bond lengths and angles have been deposited at the Cambridge Crystallographic Data Centre. See Instructions for Authors, *J. Chem. Soc., Dalton Trans.*, 1995, Issue 1, pp. xxv–xxx.



Scheme 2 Synthesis of complex 4. (i) thf, -78°C

distances. The molecular structure of one molecule is shown in Fig. 1.

In contrast to the reactions of the mononuclear complexes described above, the 1-methylene-4-phenylcyclohexadiene tris-osmium carbonyl product **4** contains a carbidic C=C double bond which remains in the co-ordination sphere of the tris-osmium cluster and the whole organic ligand still occupies a face-capping position. The ligand is co-ordinated formally by three C=C 'double bonds' which are slightly shorter than the non-co-ordinated C-C 'single bonds'. The co-ordinated part of the cyclohexadienyl ring is not strictly planar but is slightly bowed probably to enable a better interaction between the metals and the 'double bonds'. This is in contrast to the case of the cyclohexadienone complex **3** where the co-ordinated section of the ring remains planar.⁶ The sp^3 -hybridised carbon atom bearing the phenyl substituent is displaced out of the ring co-ordination plane so that the dihedral angle between the ring plane and the plane containing the two hinge carbons and this carbon is 24.8° compared to 19.5° in **3**.

The second product, which has not yet been characterised fully, is assumed to be an adduct between the phosphine and the carbene ligand forming a ylidic structure. The carbene complex **2** has in common with Fischer carbene complexes the reactivity towards this nucleophilic Wittig reagent. The product, however, is unique since the C=C double bond formed remains in a side-on co-ordination mode to the osmium cluster.

Acknowledgements

We thank the European Union for a Human Capital and Mobility Fellowship (for J. U. K.).




Fig. 1 Molecular structure of complex 4. Selected bond lengths (Å) and angles ($^{\circ}$) (parameters for the second molecule are given in square brackets): Os(1)–Os(2) 2.906(2) [2.920(2)], Os(1)–Os(3) 2.900(2) [2.892(2)], Os(2)–Os(3) 2.864(2) [2.888(2)], Os(1)–C(1) 2.30(3) [2.37(3)], Os(1)–C(11) 2.30(3) [2.31(3)], Os(2)–C(12) 2.32(3) [2.29(3)], Os(2)–C(13) 2.40(3) [2.31(3)], Os(3)–C(15) 2.30(3) [2.31(3)], Os(3)–C(16) 2.29(2) [2.28(3)], C(1)–C(11) 1.40(4) [1.36(4)], C(11)–C(12) 1.47(4) [1.48(4)], C(12)–C(13) 1.44(4) [1.39(4)], C(13)–C(14) 1.57(4) [1.52(4)], C(14)–C(15) 1.53(3) [1.51(4)], C(15)–C(16) 1.40(3) [1.27(4)] and C(11)–C(16) 1.50(4) [1.50(4)]; Os(3)–Os(1)–Os(2) 59.09(4) [59.59(4)], Os(3)–Os(2)–Os(1) 60.35(4) [59.73(4)], Os(2)–Os(3)–Os(1) 60.55(4) [60.67(4)], C(1)–C(11)–C(12) 122(2) [118(3)], C(1)–C(11)–C(16) 120(2) [124(3)], C(12)–C(11)–C(16) 118(2) [118(3)], C(11)–C(12)–C(13) 122(3) [116(3)], C(12)–C(13)–C(14) 119(3) [126(3)], C(13)–C(14)–C(15) 113(3) [106(3)], C(14)–C(15)–C(16) 120(2) [127(3)] and C(11)–C(16)–C(15) 122(2) [120(3)]

References

- H. Wadepohl, *Angew. Chem.*, 1992, **104**, 253; *Angew. Chem., Int. Ed. Engl.*, 1992, **31**, 247.
- D. Braga, P. J. Dyson, F. Grepioni and B. F. G. Johnson, *Chem. Rev.*, 1994, **94**, 1585.
- G. A. Somorjai, *J. Phys. Chem.*, 1990, **94**, 1013.
- M. P. Gomez-Sal, B. F. G. Johnson, J. Lewis, P. R. Raithby and A. H. Wright, *J. Chem. Soc., Chem. Commun.*, 1985, 1682.
- M. A. Gallop, B. F. G. Johnson, J. Lewis and A. H. Wright, *J. Chem. Soc., Dalton Trans.*, 1989, 481.
- A. J. Edwards, M. A. Gallop, B. F. G. Johnson, J. U. Köhler, J. Lewis and P. R. Raithby, *Angew. Chem.*, 1994, **106**, 1166; *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1093.
- C. P. Casey and T. J. Burkhardt, *J. Am. Chem. Soc.*, 1972, **94**, 6544.
- SHELXTL PLUS, Version 4.0, Siemens Analytical X-Ray Instruments Inc., Madison, WI, 1990.
- SHELXL 93, G. M. Sheldrick, Package for crystal structure refinement, University of Göttingen, 1993.

Received 2nd August 1995; Communication 5/05169J