Inorg. Chem. 2005, 44, 8561-8568



Synthesis and Structure of a Distorted Octahedral Palladium(II) Complex Coordinated with a Tetrathioether Ligand Tethered with Bulky Substituents

Nobuhiro Takeda, Daisuke Shimizu, and Norihiro Tokitoh*

Institute for Chemical Research, Kyoto University, Gokasho, Uji, Kyoto 611-0011, Japan

Received June 11, 2005

A new type of an *o*-phenylene-bridged tetrathioether ligand tethered with extremely bulky substituents, 2,4,6-tris-[bis(trimethylsilyl)methyl]phenyl (Tbt) groups, at its terminal sulfur atoms, TbtS[(*o*-phenylene)S]₃Tbt (1), was synthesized by taking advantage of the coupling reaction of thiols with iodobenzenes using Cu₂O in 2,4,6-trimethylpyridine. Complexation of 1 with Na₂PdCl₄ gave the corresponding dichloropalladium(II) complex, [PdCl₂(1)] (7). The X-ray structural analysis of 7 indicated that the central palladium metal is in a distorted octahedral environment, where the two inner sulfur atoms of 1 and the two chlorine atoms form a square planar arrangement around the palladium metal and the two terminal sulfur atoms of 1 weakly coordinate to the palladium center at the axial positions. In addition, a phenyl analogue of 1, PhS[(*o*-phenylene)S]₃Ph (2), was synthesized by a method similar to that for 1. Reaction of 2 with Na₂PdCl₄ gave the corresponding dichloropalladium(II) complex, [PdCl₂(2)] (8). X-ray crystallography of 8 showed a type of the structure different from the distorted octahedral structure in 7, i.e., a square planar arrangement around the central palladium atom with the one terminal sulfur atom of 2, its neighboring sulfur atom, and the two chlorine atoms. The results of the NMR studies on 8 in a CDCl₃ solution were not consistent with the results of the X-ray crystallography and suggested the coordination of the two inner sulfur atoms of 2 to the palladium metal, although a possibility of the existence of the rapid interconversion among isomers could not be excluded.

Introduction

Palladium(II) complexes generally have square planar geometry around the palladium atom, and six-coordinate complexes of palladium(II) are very rare. Cyclic poly-thioether (crown thioether) ligands such as 1,4,7-trithia-cyclononane (9S3),¹ 1,4,7,10,13,16-hexathiacyclooctadecane (18S6),² and 1,4,7-trithiacyclodecane (10S3)³ reportedly complexed with palladium(II) metals to give the corresponding five- or six-coordinate palladium(II) complexes, in which the four sulfur atoms are arranged in a square planar fashion

10.1021/ic050944v CCC: \$30.25 © 2005 American Chemical Society Published on Web 10/01/2005

around the palladium(II) atom and the remaining one or two sulfur atoms weakly coordinate to the palladium(II) atom [Pd···S: 2.96–3.27 Å] at the axial positions. In addition, the 9S3 ligand has been known to stabilize mononuclear palladium(III) species,⁴ which are generally known to be unstable. These unique properties of cyclic polythioether ligands are considered to be caused by their moderate, geometric preference of facial coordination and their ability to adjust the coordination modes and conformations to the geometric preference of the oxidation states of the metals.⁵ Acyclic polythioethers can more easily change their coor-

^{*} To whom correspondence should be addressed. E-mail: tokitoh@ boc.kuicr.kyoto-u.ac.jp.

 ⁽a) Wieghardt, K.; Küppers, H.-J.; Raabe, E.; Krüger, C. Angew. Chem., Int. Ed. Engl. 1986, 25, 1101–1103. (b) Blake, A. J.; Holder, A. J.; Hyde, T. I.; Roberts, Y. V.; Lavery, A. J.; Schröder, M. J. Organomet. Chem. 1987, 323, 261–270.

⁽²⁾ Blake, A. J.; Gould, R. O.; Lavery, A. J.; Schröder, M. Angew. Chem., Int. Ed. Engl. 1986, 25, 274–276.

^{(3) (}a) Grant, G. J.; Sanders, K. A.; Setzer, W. N.; VanDerveer, D. G. Inorg. Chem. 1991, 30, 4053–4056. (b) Chandrasekhar, S.; McAuley, A. Inorg. Chem. 1992, 31, 2663–2665.

⁽⁴⁾ Blake, A. J.; Holder, A. J.; Hyde, T. I.; Schröder, M. J. Chem. Soc., Chem. Commun. 1987, 987–988.

⁽⁵⁾ For reviews on crown thioethers, see: (a) Cooper, S. R. Acc. Chem. Res. 1988, 21, 141–146. (b) Blake, A. J.; Schröder, M. Adv. Inorg. Chem. 1990, 35, 1–80. (c) Cooper, S. R.; Rawle, S. C. Struct. Bonding (Berlin) 1990, 72, 1–72. (d) Kellogg, R. M. In Crown Compounds: Toward Future Applications, Cooper, S. R., Ed.; Wiley VCH: New York, 1992. (e) Levason, W.; Reid, G. In Comprehensive Coordination Chemistry II, from Biology to Nanotechnology; Lever, A. B. P., Ed.; Elsevier Ltd.: Oxford, U.K., 2004; Vol. 1, pp 399–410.



dination modes and conformations; however, the coordination chemistry of acyclic polythioether ligands has been less investigated probably due to their relatively low coordination ability to metals.^{5b,c,6}

On the other hand, we have succeeded in the syntheses of a variety of highly reactive low-coordinate species containing main group metals by taking advantage of effective steric protection groups, 2,4,6-tris[bis(trimethylsilyl)methyl]phenyl (Tbt) and 2,6-bis[bis(trimethylsilyl)methyl]-4-[tris(trimethylsilyl)methyl]phenyl (Bbt) groups.⁷ In addition, we have recently succeeded in the application of the Tbt and Bbt groups to the stabilization of transition metal complexes, i.e., the synthesis of the first platinum–disulfido and –diselenido complexes by the use of extremely bulky aryldimethylphosphines bearing the Tbt or Bbt group.⁸

In this paper, we report the synthesis of a new bulky tetrathioether ligand 1 (Chart 1) tethered with two Tbt substituents and its application to the synthesis of a palladium(II) complex.⁹ It is expected that the introduction of bulky substituents to an acyclic tetrathioether ligand leads to the moderate restriction of the conformation of the ligand and the kinetic stabilization of the metal center. To investigate the effect of the bulky substituents, synthesis of the phenyl analogue of 1 (2) and its complexation with a palladium metal are also described.

Results and Discussion

Synthesis of Tetrathioether Ligands 1 and 2. The synthesis of tetrathioether ligand 1 was examined by successive coupling reactions of thiols with iodobenzenes. Tbt-substituted thiol 3 was synthesized by the reaction of TbtLi, prepared from TbtBr and *tert*-butyllithium, with elemental sulfur, followed by the treatment of the resulting TbtSLi with H_2O (Scheme 1).

Coupling reaction of **3** with 1,2-diiodobenzene using Cu_2O in refluxing 2,4,6-trimethylpyridine¹⁰ afforded sulfide **5** in

- (7) (a) Okazaki, R.; Tokitoh, N. Acc. Chem. Res. 2000, 33, 625–630. (b) Tokitoh, N. Acc. Chem. Res. 2004, 37, 86–94. (c) Tokitoh, N. Bull. Chem. Soc. Jpn. 2004, 77, 429–441.
- (8) (a) Nagata, K.; Takeda, N.; Tokitoh, N. Angew. Chem., Int. Ed. 2002, 41, 136–138. (b) Nagata, K.; Takeda, N.; Tokitoh, N. Chem. Lett. 2003, 32, 170–171. (c) Nagata, K.; Takeda, N.; Tokitoh, N. Bull. Chem. Soc. Jpn. 2003, 76, 1577–1587.
- (9) A part of this paper had been preliminarily reported: Tokitoh, N.; Shimizu, D.; Takeda, N.; Sasamori, T. *Phosphorus, Sulfur, Silicon* 2005, 180, 1241–1245.
- (10) The synthesis of diaryl ethers by the reaction of phenols with aryl halides using Cu₂O catalyst in 2,4,6-trimethylpyridine has been reported; see: Bacon, R. G. R.; Stewart, O. J. J. Chem. Soc. 1965, 4953-4961.

Scheme 1. Synthesis of Tetrathioether Ligands 1 and 2



52% yield. Further coupling reaction of **5** with 0.5 equiv of 1,2-benzenedithiol under conditions similar to the above coupling reaction using Cu_2O gave the expected tetrathioether **1** in a good yield.

To study the effects of bulky substituents, we examined the synthesis of a ligand bearing phenyl groups instead of Tbt groups. The phenyl-substituted ligand 2 was synthesized by the method similar to that for 1, though in a relatively low yield (Scheme 1).

Structure of Tetrathioether Ligands 1 and 2. Tetrathioether ligands 1 and 2 were characterized by the NMR spectra, MS, and elemental analysis, and their molecular structures in the crystalline states were definitively determined by X-ray structural analyses. Figures 1 and 2 show the ORTEP drawings of 1 and 2, respectively, and the selected bond lengths, angles, and torsion angles are summarized in Table 1. While tetrathioether 2 has a conformation close to C_2 symmetry, in the case of 1 the conformations around the S2 and S3 atoms are very different from each other. All the C-S bond lengths of 1 and 2 [1.76-1.78 Å] are usual as those of common diaryl sulfides [1.762–1.774 Å].¹¹ It is noteworthy that each of the C1-S1-C28 and C45-S4-C46 planes in 1 is nearly perpendicular to the benzene ring of its connecting Tbt group $(C28-S1-C1-C6 = 91.3(5)^{\circ})$ and C45-S4-C46-C47 = $80.6(5)^\circ$) and is almost parallel to its neighboring *o*-phenylene ring (C1-S1-C28-C33 = $178.0(4)^{\circ}$ and C46-S4-C45-C40 = 175.1(5)°). This conformation in 1 can be explained in terms of the steric hindrance caused by the Tbt groups.



Figure 1. ORTEP drawing of **1** with thermal ellipsoids (50% probability). Hydrogen atoms, trimethylsilyl groups, and a solvent molecule were omitted for clarity.

^{(6) (}a) Levason, W.; Reid, G. In Comprehensive Coordination Chemistry II, from Biology to Nanotechnology; Lever, A. B. P., Ed.; Elsevier Ltd.: Oxford, U.K., 2004; Vol. 1, pp 391–398. (b) Murray, S. G.; Hartley, F. R. Chem. Rev. **1981**, 81, 365–414.



Figure 2. ORTEP drawing of 2 with thermal ellipsoids (50% probability).

The ¹H and ¹³C NMR spectra of **1** and **2** showed that the two terminal TbtS(o-phenylene) (for **1**) and PhS(o-phenylene) (for **2**) units were equivalent, respectively, suggesting flexible structures of **1** and **2** in solution.

Synthesis and Structure of Palladium(II) Complex 7 of Tetrathioether Ligand 1. When 1 was allowed to react with Na₂PdCl₄ in ethanol at reflux for 5 min, the starting red solution turned yellow with the formation of orange precipitates. Further refluxing for 30 min resulted in the almost quantitative formation of the corresponding dichloropalladium complex 7 as orange crystals, which are slightly soluble in ethanol (eq 1). The structure of 7 was determined by NMR spectrometry, ESI MS, elemental analysis, and single-crystal X-ray diffraction.



The X-ray structural analysis of **7** showed the C_2 symmetrical distorted octahedral structure, where the inner two sulfur atoms of ligand **1** and the two chlorine atoms form a square planar arrangement around the palladium(II) atom and the two terminal sulfur atoms of **1** weakly coordinate to the palladium center at the axial positions (Figure 3). A few examples for cationic, 6-coordinate complexes of palladium(II) have been synthesized using crown thioether ligands (e.g., 9S3,¹ 18S6,² and 10S3³), S- and N-donor macrocyclic ligands [e.g., 1,4,10,13-tetrathia-7,16-diaza-cyclooctadecane ([18]aneN₂S₄)¹² and 1,4-dithia-7-azacyclononane ([9]aneNS₂)¹³], and tris(2,4,6-trimethoxyphenyl)-

Table 1.	. Selected	Bond	Lengths	(Å),	Angles	(deg),	and	Torsion
Angles (deg)							

1		2	
I \$1-C1 \$1-C28 \$2-C33 \$2-C34 \$3-C39 \$3-C40 \$4-C45 \$54-C46	1.769(6) 1.765(6) 1.766(6) 1.776(6) 1.772(6) 1.771(7) 1.771(7) 1.767(6)	$\begin{array}{c} 2\\ \hline S1-C6\\ S1-C7\\ S2-C12\\ S2-C13\\ S3-C18\\ S3-C19\\ S4-C24\\ S4-C25\\ \hline $	1.7740(16) 1.7621(16) 1.7727(16) 1.7822(16) 1.7802(15) 1.7757(16) 1.7681(16) 1.7757(15)
C28-S1-C1 C33-S2-C34 C40-S3-C39 C46-S4-C45	101.8(3) 103.9(3) 101.1(3) 103.2(3)	C7-S1-C6 C12-S2-C13 C19-S3-C18 C24-S4-C25	103.53(7) $102.48(7)$ $102.64(7)$ $104.63(7)$
$\begin{array}{c} C28-S1-C1-C6\\ C1-S1-C28-C33\\ C34-S2-C33-C28\\ C33-S2-C34-C39\\ C40-S3-C39-C34\\ C39-S3-C40-C45\\ C46-S4-C45-C40\\ C45-S4-C46-C47\\ \end{array}$	$\begin{array}{c} 91.3(5)\\ 178.0(4)\\ 66.5(5)\\ -156.1(5)\\ 139.2(5)\\ 113.6(5)\\ -175.1(5)\\ 80.6(5) \end{array}$	$\begin{array}{c} C1-C6-S1-C7\\ C6-S1-C7-C12\\ C7-C12-S2-C13\\ C12-S2-C13-C18\\ C13-C18-S3-C19\\ C18-S3-C19-C24\\ C19-C24-S4-C25\\ C24-S4-C25-C30\\ \end{array}$	$\begin{array}{r} -123.94(13)\\ -157.99(13)\\ -77.49(14)\\ 164.94(12)\\ 165.82(12)\\ -78.62(14)\\ 167.38(12)\\ 132.39(12)\end{array}$

phosphine ligands.¹⁴ By contrast, to our knowledge, there has been no report on the X-ray structural analysis of a neutral, 6-coordinate palladium(II) complex except for 6-coordinate complexes having intermolecular or intra-molecular Pd····H interactions.¹⁵

The Pd1-Cl1 (2.3117(9) Å) bond distance is within the range of those in the reported tetracoordinate palladium(II) complexes (2.298–2.354 Å).¹⁶ The Pd1–S1 [2.2615(9) Å] bond distance is close to those of some 4- or 5-coordinate cis-dichloropalladium(II) complexes having macrocyclic thioether ligands $(2.246-2.269 \text{ Å})^{17}$ and is slightly shorter than those of cationic palladium(II)-crown thioether complexes $(2.31-2.33 \text{ Å})^{1,2}$ except for $[Pd(10S3)_2](PF_6)_2$ [2.2584(15) and 2.2789(14) Å].^{3a} The distances between the palladium and the two terminal sulfur atoms $[Pd1 \cdots S2 =$ 3.1755(8) Å] are shorter than the sum of the van der Waals radii of palladium and sulfur atoms (3.43 Å).¹⁸ This suggests weak interactions between the palladium (Pd1) and the terminal sulfur atoms (S2 and S2*) in the crystalline state. It is noteworthy that the torsion angles around the terminal sulfur atoms in palladium complex 7 [C9-S2-C10-C11 $= 94.5(3)^{\circ}$, C4-C9-S2-C10 $= 178.6(3)^{\circ}$] are almost similar to those of the free ligand 1. This result suggests that the extremely bulky Tbt groups put restrictions on the conformation of ligand 1 in the structure of complex 7 as

- (14) Dunbar, K. R.; Sun, J.-S. J. Chem. Soc., Chem. Commun. 1994, 2387–2388.
- (15) (a) Bailey, N. A.; Mason, R. J. Chem. Soc. A 1968, 2594–2605. (b) Debaerdemaeker, T.; Kutoglu, A.; Schmid, G.; Weber, L. Acta Crystallogr., Sect. B 1973, 29, 1283–1288.
- (16) Orpen, A. G.; Brammer, L.; Allen, F. H.; Kennard, O.; Watson, D. G.; Taylor, R. J. Chem. Soc., Dalton Trans. 1989, S1–S83.
- (17) (a) Arca, M.; Blake, A. J.; Lippolis, V.; Montesu, D. R.; McMaster, J.; Tei, L.; Schröder, M. Eur. J. Inorg. Chem. 2003, 1232–1241. (b) Lucas, C. R.; Liang, W.; Miller, D. O.; Bridson, J. N. Inorg. Chem. 1997, 36, 4508–4513. (c) Lee, G.-H.; Tzeng, B.-C. Acta Crystallogr. 1996, C52, 879–882. (d) Blake, A. J.; Freeman, G.; Schröder, M.; Yellowlees, L. J. Acta Crystallogr. 1993, C49, 167–168. (e) Blake, A. J.; Holder, A. J.; Roberts, Y. V.; Schröder, M. Acta Crystallogr. 1988, C44, 360–361.
- (18) Bondi, A. J. Phys. Chem. 1964, 68, 441-451.

⁽¹¹⁾ Allen, F. H.; Kennard, O.; Watson, D. G.; Brammer, L.; Orpen, A. G.; Taylor, R. J. Chem. Soc., Perkin Trans. 2 1987, S1–S19.

 ^{(12) (}a) Reid, G.; Blake, A. J.; Hyde, T. I.; Schröder, M. J. Chem. Soc., Chem. Commun. 1988, 1397–1399. (b) Blake, A. J.; Reid, G.; Schröder, M. J. Chem. Soc., Dalton Trans. 1990, 3363–3373.

⁽¹³⁾ Blake, A. J.; Crofts, R. D.; de Groot, B.; Schröder, M. J. Chem. Soc., Dalton Trans. 1993, 485–486.



Figure 3. ORTEP drawing of **7** with thermal ellipsoids (50% probability). Hydrogen atoms and solvent molecules were omitted for clarity. Selected bond distances (Å), angles (deg), and torsion angles (deg): Pd1-S1 = 2.2615(9), Pd1-C11 = 2.3117(9), $Pd1\cdots S2 = 3.1755(8)$; S1-Pd1-S1* = 89.45(5), S1-Pd1-C11 = 88.15(3), C11-Pd1-C11* = 95.37(5), S1-Pd1-S2 = 71.19(3), S1*-Pd1-S2 = 82.08(3), C11-Pd1-S2 = 89.37(3), C11*-Pd1-S2 = 116.55(3); C4-C9-S2-C10 = 178.6(3), C9-S2-C10-C11 = 94.5(3).

well as in that of 1; that is, the ligand 1 preferably has the conformation in which the C-S(terminal)-C planes are almost perpendicular to the benzene rings of the Tbt groups and almost parallel to the neighboring *o*-phenylene rings.

As for the cationic 6-coordinate complexes of palladium(II) with crown thioether ligands or S- and N-donor macrocyclic ligands, it has been known that complexes with relatively strong interactions with the sulfur atom at the axial positions have a blue or green color, while complexes with relatively weak or no interactions have a yellow or brown color. For example, [Pd(9S3)₂][PF₆]₂ [ax Pd-S 2.952(4) Å],¹ $[Pd(10S3)_2][BPh_4]_2$ [ax Pd-S 3.11 Å],^{3a} $[Pd(10S3)_2][PF_6]_2$ [ax Pd-S 3.034(1) Å]^{3b} and $[Pd([18]aneN_2S_4)][BPh_4]_2$ [ax Pd-S 3.034(1) Å]Pd-S 2.954(4) and 3.000(3) Å]¹² have an absorption maximum in the visible region at 615 (blue), 602 (bluegreen), 598 (green), and 514 nm (green) respectively, while [Pd(18S6)](PF₆)₂ [ax Pd-S 3.2730(17) Å] has yellow-brown color.² These absorptions around 500-600 nm have been assigned to d-d transitions by the small values of the extinction coefficient. Complex 7, the X-ray crystallographic analysis of which suggests the weak interactions between the palladium (Pd1) and the terminal sulfur atoms (S2 and S2*) in the crystalline state, has an orange color with absorption maxima at 398 (ϵ 6.5 × 10³), 305 (sh, 7.6 × 10³), and 294 (sh. 5.3 \times 10³) nm in hexane. The diffuse UV/vis reflectance spectrum of 7 in the solid state showed the



Figure 4. ORTEP drawing of **8** with thermal ellipsoids (50% probability). Selected bond distances (Å) and angles (deg): Pd1-S1 = 2.2448(7), Pd1-S2 = 2.2642(8), Pd1-Cl1 = 2.3115(8), Pd1-Cl2 = 2.3077(8), $Pd1\cdots S3 = 3.4305(8)$; S1-Pd1-S2 = 90.22(3), S1-Pd1-Cl1 = 87.92(3), S2-Pd1-Cl2 = 88.76(3), Cl1-Pd1-Cl2 = 93.35(3).

reflectance decrease in the region similar to the absorption region of **7** in hexane. This region is similar to the absorption region of square planar *cis*-dichloropalladium(II) complexes with dithioether ligands ($\lambda_{max} = 380-450$ nm).¹⁹ If the trend of the UV/vis spectra for the cationic 6-coordinate complexes of palladium(II) with these macrocyclic ligands can be applied to the neutral dichloropalladium(II) complex **7**, this result may suggest relatively weak or no interaction between the palladium and the terminal sulfur atoms both in the solid state and in solution.

Synthesis and Structure of Palladium(II) Complex 8 of Tetrathioether Ligand 2. To investigate the effects of bulky substituents, we examined the synthesis of a dichloropalladium complex coordinated with tetrathioether ligand 2. When a mixture of **2** and Na₂PdCl₄ was refluxed in ethanol for 1 h, the corresponding dichloropalladium complex 8 was obtained as orange crystals (eq 2). The X-ray structural analysis of 8 showed that the palladium metal is coordinated by the one terminal and the one inner sulfur atoms of ligand 2 (Figure 4) in contrast to the distorted octahedral coordination structure of Tbt-analogue 7. Complex 8 has a square planar structure with normal Pd-Cl bond lengths. The Pd1-S1 and Pd1-S2 bond lengths are similar to the Pd1-S1 bond lengths of 7, and the Pd1–S3 distance [3.4305(8) Å] is close to the sum of the van der Waals radii of palladium and sulfur atoms (3.43 Å).¹⁸ This result suggests that there is little interaction between Pd1 and S3 atoms in complex 8.

$$2 \xrightarrow{\text{Na}_2\text{PdCl}_4 (1.0 \text{ eq})}_{\text{EtOH, reflux, 1 h}} [\text{PdCl}_2(2)]$$
(2)

To clarify the reason for the preference of the **8b**-type structure over the **8a**-type structure in the crystalline state (Chart 2), the energies of these two linkage isomers, **8a**,**b**, were calculated with B3LYP/LANL2DZ(ECP) for the Pd and 6-31G(d) for C, S, Cl, and H (Figure 6).²⁰ Contrary to

^{(19) (}a) Hartley, F. R.; Murray, S. G.; Levason, W.; Soutter, F. E.; McAuliffe, C. A. *Inorg. Chim. Acta* **1979**, *35*, 265–277. (b) Takeda, N.; Shimizu, D.; Sasamori, T.; Tokitoh, N. *Acta Crystallogr., Sect. E* **2005**, *61*, m1408–m1410.

 ^{(20) (}a) Becke, A. D. J. Chem. Phys. 1993, 98, 5648-5652. (b) Becke, A. D. Phys. Rev. 1988, A38, 3098-3100. (c) Lee, C.; Yang, W.; Parr, R. G. Phys. Rev. 1988, B37, 785-789.

Chart 2



the result of the X-ray structural analysis, the theoretical calculations revealed that **8a** is more stable than **8b** by 4.3 kcal mol⁻¹. Furthermore, the conformation of the optimized structure for **8b** was different from that of the X-ray structure (Figures 4 and 6). Taking these results into consideration, the preference of the **8b**-type structure in the crystalline state may be interpreted in terms of the effects of the intermolecular interactions, which are observed in the X-ray structural analysis (Figure 5).

On the other hand, the calculated structure for **8a** showed the longer distances between the central palladium atom and the terminal sulfur atoms (Pd1...S1 = 3.575, Pd1...S4 = 3.589 Å) compared with those in complex **7** (Figures 3 and 6). This difference may be explained in terms of the effects of the bulky substituents; i.e., the steric hindrance of the Tbt groups in **7** leads to the conformation suitable for the



Figure 5. Intermolecular interaction in the crystalline state of **8**. Selected intermolecular C–C distances (Å): C14…C14* 3.461(3), C29…C2* 3.599(5), C14…C21* 3.582(4), C15…C21* 3.477(4), C15…C22* 3.460(4).



Figure 6. (a) Theoretically optimized structure of **8a**. Selected bond distances (Å) and angles (deg): Pd1-S2 = 2.377, Pd1-S3 = 2.377, Pd1-Cl1 = 2.373, Pd1-Cl2 = 2.373, Pd1-rs1 = 3.575, Pd1-rs1 = 3.575, Pd1-rs1 = 3.589; S2-Pd1-S3 = 88.323, S2-Pd1-Cl1 = 89.054, Cl1-Pd1-Cl2 = 94.085, S3-Pd1-Cl2 = 89.026, S1-Pd1-S2 = 65.8, S1-Pd1-S3 = 73.0, Cl1-Pd1-S1 = 101.0, Cl2-Pd1-S1 = 118.5, S4-Pd1-S2 = 72.2, S4-Pd1-S3 = 65.6, Cl1-Pd1-S4 = 118.5, Cl2-Pd1-S4 = 101.7. (b) Theoretically optimized structure of **8b**. Selected bond distances (Å) and angles (deg): Pd1-S1 = 2.383, Pd1-S2 = 2.369, Pd1-Cl1 = 2.375, Pd1-Cl2 = 2.366, Pd1-rs1 = 2.377, S1-Pd1-S2 = 88.331; S1-Pd1-Cl2 = 88.757, Cl1-Pd1-Cl2 = 94.742.

Table 2. Redox Potentials (V vs Fc/Fc^+) of 7–9

type	E_{pa}	$E_{\rm pc}$	$E_{1/2}$
oxidn ^a	-0.24	-0.33	-0.29
oxidn ^b	-0.04	-0.13	-0.09
redn ^b		-1.84	
oxidn ^a	+0.97		
redn ^b		-1.18	
oxidn ^a	-0.17	-0.26	-0.22
oxidn ^b	-0.02	-0.10	-0.06
redn ^b		-1.28	
	type oxidn ^a oxidn ^b redn ^b oxidn ^a oxidn ^a oxidn ^b redn ^b	$\begin{array}{c c} {\rm type} & E_{\rm pa} \\ \hline {\rm oxidn}^a & -0.24 \\ {\rm oxidn}^b & -0.04 \\ {\rm redn}^b \\ {\rm oxidn}^a & +0.97 \\ {\rm redn}^b \\ {\rm oxidn}^a & -0.17 \\ {\rm oxidn}^b & -0.02 \\ {\rm redn}^b \end{array}$	$\begin{array}{c cccc} type & E_{pa} & E_{pc} \\ \hline type & -0.24 & -0.33 \\ oxidn^{b} & -0.04 & -0.13 \\ redn^{b} & -1.84 \\ oxidn^{a} & +0.97 \\ redn^{b} & -1.18 \\ oxidn^{a} & -0.17 & -0.26 \\ oxidn^{b} & -0.02 & -0.10 \\ redn^{b} & -1.28 \\ \hline \end{array}$

^a 0.1 M (n-Bu)₄NPF₆ in CH₂Cl₂. ^b 0.1 M (n-Bu)₄NPF₆ in THF.

coordination of the two terminal sulfur atoms to the central palladium atom.

Interestingly, the ¹³C NMR spectrum of complex **8** in CDCl₃ at room temperature indicated that complex **8** has C_2 symmetry. This result suggests that **8** has the **8a**-type structure in solution, which is consistent with not the X-ray structure but the results of theoretical calculations. However, we cannot exclude a possibility of the existence of the rapid equilibrium among isomers such as **8a**,**b** in solution within the time scale of ¹³C NMR spectrometry at room temperature.²¹

Electrochemistry of Palladium(II) Complexes 7 and 8. As described in the Introduction, the cyclic polythioether ligands have been known to stabilize mononuclear palladium(III) species,⁴ which are generally known to be unstable. We investigated the electrochemical properties of complexes 7 and 8, comparing with those of [PdCl₂{PhS(o-phenylene)-SPh}] (9).¹⁹ The redox potentials of 7-9 were summarized in Table 2. The cyclic voltammogram of 7 showed a chemically reversible one-electron oxidation $[E_{1/2} = -0.29]$ V (0.1 M (*n*-Bu)₄NPF₆ in CH₂Cl₂) or $E_{1/2} = -0.09$ V (0.1 M $(n-Bu)_4$ NPF₆ in THF) vs Fc/Fc⁺ (ferrocene/ferrocenium)] and an irreversible reduction $[E_{pc} = -1.84 \text{ V} (0.1 \text{ M} (n-1.05 \text{ V}))]$ Bu)₄NPF₆ in THF) vs Fc/Fc⁺]. Since the $E_{1/2}$ values in the oxidation of 7 are close to those in the reversible one-electron oxidation of 9 [$E_{1/2} = -0.22$ V (0.1 M (*n*-Bu)₄NPF₆ in CH_2Cl_2) or $E_{1/2} = -0.06 V (0.1 M (n-Bu)_4 NPF_6 in THF)$ vs Fc/Fc^+], both these oxidations of 7 and 9 may be assigned to Pd(II)/Pd(III) couples as well as the case of the Pd complexes with crown thioethers.^{1b,3} The differential pulse voltammograms of 7 and 9 suggested that the irreversible reduction waves of 7 and 9 are one- and two-electron processes, respectively, indicating that these reductions may be assigned to different processes. On the other hand, cyclic voltammograms of 8 showed irreversible oxidation and reduction at $E_{pa} = +0.97 \text{ V} (0.1 \text{ M} (n-\text{Bu})_4\text{NPF}_6 \text{ in CH}_2\text{Cl}_2)$ and $E_{pc} = -1.18 \text{ V} (0.1 \text{ M} (n-\text{Bu})_4\text{NPF}_6 \text{ in THF}) \text{ vs Fc/Fc}^+$, respectively. It is interesting that the redox process of 8 is considered to be completely different from those of 7.

⁽²¹⁾ The ¹³C DEPT 90 NMR spectra of 8 at low temperatures (0, -20, -40, and -60 °C) showed the upper or lower field shifts of some signals and the appearance of some new peaks. However, these measurements did not lead to the observation of splitting of the signals, which indicates the existence of the rapid equilibrium among isomers such as 8a,b. These spectroscopic behaviors could not be analyzed in detail, though suggesting the existence of some equilibrium.

Conclusion

A new tetrathioether ligand **1** tethered with two bulky substituents, Tbt groups, was synthesized by taking advantage of the coupling reaction of thiols with iodobenzenes using Cu_2O in 2,4,6-trimethylpyridine. Complexation of **1** with Na₂PdCl₄ gave the corresponding dichloropalladium complex **7**, the X-ray structural analysis of which indicated its distorted octahedral structure with weak intramolecular interactions between the two terminal sulfur atoms and the central palladium metal.

On the other hand, a phenyl-substituted tetrathioether ligand **2** reacted with Na₂PdCl₄ to give the corresponding dichloropalladium complex **8**. X-ray crystallography revealed the structure of complex **8** in the crystalline state, where the dichloropalladium part is coordinated by the one terminal sulfur atom and the neighboring sulfur atom of **2**. By contrast, the ¹³C NMR spectrum of **8** in CDCl₃ at room temperature suggested a type of structure in solution different from that in the crystalline state, i.e., the **8a**-type structure, where the two inner sulfur atoms of **2** coordinate to the central metal. However, a possibility of the existence of rapid equilibriums among isomers such as **8a,b** in solution within the time scale of ¹³C NMR spectrometry could not be excluded.

We wish to propose a possible explanation for the distorted octahedral structure of **7** in the crystalline state as follows. Ligand **1** preferably has a conformation, in which the C–S (terminal)–C planes are almost perpendicular to the benzene rings of the Tbt groups and almost parallel to the neighboring *o*-phenylene rings, because of the steric effects of the bulky Tbt groups. It is considered that ligand **1** with this conformation can effectively stabilize the distorted octahedral structure.

Further studies on the properties and applications of the dichloropalladium complexes 7 and 8 are very interesting. In addition, the ligands 1 and 2 are expected to be useful for the synthesis of unusual complexes.

Experimental Section

General Procedures. All reactions were carried out under an argon atmosphere, unless otherwise noted. THF was purified by distillation from sodium diphenylketyl before use. Preparative gel permeation liquid chromatography (GPLC) was performed on an LC-908 or LC-918 instrument with JAI gel 1H+2H columns (Japan Analytical Industry) using chloroform or toluene as an eluent. Wet column chromatography (WCC) and short column chromatography were performed with Nacalai Tesque silica gel 60. The ¹H NMR (300 MHz) and ¹³C NMR (75 MHz) spectra were measured in CDCl₃ with a JEOL AL-300 spectrometer using CHCl₃ (7.25 ppm) as internal standards for ¹H NMR spectroscopy and CDCl₃ (77.0 ppm) as those for ¹³C NMR spectroscopy. High-resolution mass spectral data were obtained on a JEOL JMS-700 spectrometer. The electronic spectra were recorded on a JASCO V-570 UV/vis spectrometer, and diffuse UV/vis reflectance spectra were recorded on a JASCO V-570 UV/vis spectrometer equipped with an integrating sphere accessory. All melting points were determined on a Yanaco micro melting point apparatus and are uncorrected. Elemental analyses were performed by the Microanalytical Laboratory of the Institute for Chemical Research, Kyoto University. The electrochemical experiments were carried out with an ALS 602A potentiostat/galvanostat using a glassy carbon disk working electrode, a Pt wire counter electrode, and Ag/0.01 M AgNO₃ in 0.1 M *n*-Bu₄NPF₆/CH₃CN reference electrode. The measurements were carried out in CH₂Cl₂ and THF solution containing 0.1 M *n*-Bu₄NPF₆ as a supporting electrolyte with scan rates of 50-300 mV s⁻¹ at ambient temperature.

Synthesis of 2,4,6-Tris[bis(trimethylsilyl)methyl]benzenethiol (3). To a THF solution (120 mL) of TbtBr²² (7.54 g, 11.9 mmol) was gradually added t-BuLi (2.2 M pentane solution, 11.9 mL, 26.2 mmol) at -78 °C. After the yellow solution was stirred for 1 h at this temperature, S_8 (1.14 g, 4.46 mmol) was added. The reaction mixture was stirred at -78 °C for 1 h and then at room temperature for 3 h. The resulting dark brown solution was poured into ice water. The mixture was extracted with hexane (200 mL), and the organic layer was dried with MgSO₄. After removal of the solvents under reduced pressure, reprecipitation from CH₂Cl₂/CH₃CN gave thiol **3** (6.27 g, 10.7 mmol, 90%) as colorless crystals. **3**: colorless solid; mp 167.8–169.8 °C; ¹H NMR (300 MHz, CDCl₃) δ –0.01 (s, 36H), 0.00 (s, 18H), 1.27 (s, 1H), 2.39 (s, 1H), 2.70 (s, 1H), 2.82 (s, 1H), 6.32 (s, 1H), 6.44 (s, 1H); 13 C NMR (75 MHz, CDCl₃) δ 0.4 (q), 0.6 (q), 28.0 (d \times 2), 30.0 (d), 120.8 (s), 121.8 (d), 126.6 (d), 141.6 (s), 146.8 (s \times 2); HRMS (EI; m/z) found 584.3018 $([M]^+)$, calcd for C₂₇H₆₀SSi₆ 584.3031. Anal. Calcd for C₂₇H₆₀-SSi₆: C, 55.40; H, 10.33. Found: C, 55.26; H, 10.27.

Synthesis of 2-{2,4,6-Tris[bis(trimethylsilyl)methyl]phenylthio}iodobenzene (5). To a mixture of thiol 3 (6.0 g, 10.3 mmol) and Cu₂O (1.54 g, 10.8 mmol) in 2,4,6-trimethylpyridine (35 mL) was added 1,2-diiodobenzene (1.34 mL, 10.3 mmol), and the mixture was refluxed for 3 h. The reaction mixture was passed through a short column (SiO₂, hexane) to remove inorganic salts. All fractions were collected and washed with a 0.2 M aqueous solution of HCl three times (100 mL \times 3). The organic layer was dried with MgSO₄ and evaporated. The addition of EtOH to the resulting orange oil led to the formation of colorless precipitates, and the filtration of the mixture gave sulfide 5 (4.18 g, 5.31 mmol, 52%) as colorless crystals. 5: colorless solid; mp 188.1-190.1 °C; ¹H NMR (300 MHz, CDCl₃) δ -0.03 (s, 36H), 0.06 (s, 18H), 1.37 (s, 1H), 2.60 (s, 2H), 6.41 (s, 1H), 6.54 (s, 1H), 6.65 (dd, *J* = 1.5, 7.8 Hz, 1H), 6.71–6.78 (m, 1H), 7.01–7.08 (m, 1H), 7.74 (dd, *J* = 1.3, 7.9 Hz, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 0.7 (q), 0.8 (q), 26.7 (d × 2), 30.7 (d), 96.4 (s), 122.5 (d), 126.1 (s), 126.2 (d), 127.3 (d), 128.1 (d), 128.4 (d), 139.1 (d), 143.7 (s), 144.8 (s), 150.1 (s × 2); HRMS (EI; m/z) found 786.2306 ([M]⁺), calcd for C₃₃H₆₃ISSi₆ 786.2311. Anal. Calcd for C₃₃H₆₃ISSi₆: C, 50.34; H, 8.07. Found: C, 50.47; H. 8.10.

Synthesis of 1,2-Bis(2-{2,4,6-tris[bis(trimethylsilyl)methyl]phenylthio}phenylthio)benzene (1). A red-purple suspension of sulfide 5 (1.0 g, 1.27 mmol), Cu₂O (186 g, 1.30 mmol), and 1,2benzenedithiol (90 mg, 0.635 mmol) in 2,4,6-trimethylpyridine (13 mL) was refluxed for 3 h. The reaction mixture was subjected to short column chromatography (SiO₂, benzene) to remove inorganic salts. Hexane (100 mL) was added to the eluate, and then the mixture was washed with a 0.2 M aqueous solution of HCl three times (100 mL \times 3). After drying with MgSO₄, the solvents were removed under reduced pressure to give an orange oil. The addition of EtOH to the residue resulted in the formation of colorless precipitates, and the filtration of the mixture gave pure tetrathioether 1 (743 mg, 0.51 mmol, 80%) as colorless crystals. 1: colorless blocks; mp: 211.0–212.0 °C; ¹H NMR (300 MHz, CDCl₃) δ 0.03 (s, 72H), 0.08 (s, 36H), 1.38 (s, 2H), 2.66 (s, 4H), 6.43 (s, 2H), 6.58 (s, 2H), 6.61-6.66 (m, 2H), 6.93-7.04 (m, 4H), 7.05-7.16

⁽²²⁾ Okazaki, R.; Tokitoh, N.; Matsumoto, T. In Synthetic Methods of Organometallic and Inorganic Chemistry; Auner, N., Klingebiel, U., Eds.; Thieme: New York, 1996; Vol. 2, pp 260–269.

A Distorted Octahedral Pd(II)-Tetrathioether Complex

(m, 4H), 7.20–7.25 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 0.8 (q), 0.9(q), 26.3 (d × 2), 30.6 (d), 122.7 (d), 123.9 (s), 125.2 (d), 127.3 (d), 127.5 (d), 127.6 (d), 127.7 (d), 130.3 (s), 130.6 (d), 132.8 (d), 137.0 (s), 142.1 (s), 144.4 (s), 150.4 (s × 2); HRMS (FAB; *m/z*) found *m/z* 1458.6298 ([M]⁺), calcd for C₇₂H₁₃₀S₄Si₁₂ 1458.6287. Anal. Calcd for C₇₂H₁₃₀S₄Si₁₂•CHCl₃: C, 55.48; H, 8.35. Found: C, 55.27; H, 8.34.

Synthesis of 2-Iodophenyl Phenyl Sulfide (6). A red-purple suspension of benzenethiol (4) (3.51 mL, 34.2 mmol), Cu₂O (4.80 g, 33.6 mmol), and 1,2-diiodobenzene (11.1 g, 33.6 mmol) in 2,4,6-trimethylpyridine (170 mL) was refluxed for 3 h. After addition of 150 mL of hexane, the mixture was washed with 0.2 M aqueous solution of HCl four times (100 mL × 4). The organic layer was dried with MgSO₄ and evaporated. The residue was separated with WCC (SiO₂, hexane) to give sulfide **6** (3.7 g, 11.8 mmol, 35%) as colorless crystals. **6**²³ colorless solid; mp 52–53 °C; ¹H NMR (300 MHz, CDCl₃) δ 6.87 (dd, J = 7.8, ~8 Hz, 1H), 6.95 (d, J = 7.8 Hz, 1H), 7.19 (dd, J = 7.8, ~8 Hz, 1H), 7.34–7.47 (m, 5H), 7.83 (d, J = 7.8 Hz, 1H).

Synthesis of 1,2-Bis[2-(phenylthio)phenylthio]benzene (2). A red-purple suspension of sulfide 6 (3.69 g, 11.8 mmol), Cu₂O (1.69 g, 11.8 mmol), and 1,2-benzenedithiol (841 mg, 5.91 mmol) in 2,4,6-trimethylpyridine (20 mL) was refluxed for 3 h. The reaction mixture was subjected to short column chromatography (SiO₂, benzene) to remove inorganic salts. After addition of hexane (150 mL), the mixture was washed with a 0.2 M aqueous solution of HCl four times (100 mL \times 4). The organic layer was dried with MgSO₄ and evaporated under reduced pressure. The residue was subjected to WCC (SiO₂, hexane) and reprecipitated from CH₂Cl₂/ CH₃CN to give polythioether 2 (768 mg, 1.50 mmol, 25%) as colorless crystals. 2: colorless solid; mp 107.7-108.7 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.12–7.22 (m, 6H), 7.26–7.47 (m, 16H); ¹³C NMR (75 MHz, CDCl₃) δ 127.3 (d), 127.5 (d), 127.8 (d), 128.0 (d), 129.2 (d), 131.0 (d), 131.4 (d), 132.0 (d), 132.2 (d), 134.2 (s), 135.1 (s), 136.6 (s), 138.8 (s); HRMS (EI; m/z) found 510.0594 ([M]⁺), calcd for $C_{30}H_{22}S_4$ 510.0604. Anal. Calcd for $C_{30}H_{22}S_4$: C, 70.55; H, 4.34. Found: C, 70.63; H, 4.29.

Synthesis of [PdCl₂(1)] (7). Tetrathioether 1 (100 mg, 0.0684 mmol) was dissolved into EtOH (40 mL) by heating at reflux. To the solution was added Na₂PdCl₄ (40.3 mg, 0.137 mmol) using 3 mL of EtOH. The red solution was refluxed for 5 min, and the resulting orange suspension was further heated at reflux for 30 min. After evaporation of the solvents, CHCl₃ (20 mL) was added to the residue and the resulting orange suspension was filtered through Celite. The orange filtrate was evaporated to give pure palladium dichloride complex 7 (114.2 mg, quantitative). 7: orange blocks; mp 254.2–259.2 °C (dec); ¹H NMR (300 MHz, CDCl₃, 50 °C) δ -0.10 (s, 36H), 0.09 (s, 72H), 1.41 (s, 2H), 2.06 (s, 2H), 2.78 (s, 2H), 6.52 (s, 4H), 6.55 (dd, ${}^{3}J = 7.8$, ${}^{4}J = 1.5$ Hz, 2H), 7.13 (ddd, ${}^{3}J = 7.6, \sim 8$ Hz, ${}^{4}J = 1.5$ Hz, 2H), 7.20–7.26 (ddd, ${}^{3}J = 7.8, \sim 8$ Hz, ${}^{4}J = 0.8$ Hz, 2H), 7.39–7.43 (m, 4H), 7.83 (dd, ${}^{3}J = 7.6$ Hz, ${}^{4}J = 0.8$ Hz, 2H); ${}^{13}C$ NMR (75 MHz, CDCl₃, 50 °C) δ 0.5 (q), 0.8 (q), 1.4 (q), 26.5 (d), 27.2 (d), 30.9 (d), 120.4 (s), 123.2 (d), 124.3 (d), 127.8 (d), 128.1 (s), 128.3 (d), 130.6 (d), 131.3 (d), 132.1 (d), 135.9 (s), 136.3 (d), 142.8 (s), 145.8 (s), 149.9 (s), 151.8 (s); UV-vis (CHCl₃) λ_{max} 398 (ϵ 6.5 × 10³), 305 (sh, 7.6 × 10³), 294 $(sh, 5.3 \times 10^3)$; LRMS (ESI; m/z) found 1600 ($[M - Cl]^+$), calcd for $C_{72}H_{130}{}^{35}Cl^{106}PdS_4Si_{12}$ ([M - Cl]⁺) 1600. Anal. Calcd for C₇₂H₁₃₀ClPdS₄Si₁₂•CHCl₃: C, 49.88; H, 7.51. Found: C, 49.73; H, 7.42.

Synthesis of [PdCl₂(2)] (8). Tetrathioether 2 (25.0 mg, 0.0489 mmol) was dissolved into EtOH (10 mL) by heating at reflux. To the solution was added Na₂PdCl₄ (14.4 mg, 0.0489 mmol) using EtOH (3 mL). Heating of the orange solution at 60 °C for 5 min resulted in the formation of orange precipitates, and the resulting orange suspension was further refluxed for 30 min. After evaporation of the solvents, CH₂Cl₂ (100 mL) was added to the residue and the resulting orange suspension was filtered through Celite. The orange filtrate was evaporated to give pure palladium dichloride complex 8 (34.1 mg, quant) as orange crystals. 8: orange plates; mp 295.2-305.2 °C (dec); ¹H NMR (300 MHz, CDCl₃) δ 6.97-7.67 (m, 22H); ¹³C NMR (75 MHz, CDCl₃) δ 127.48 (d), 127.51 (d), 127.9 (d), 128.0 (d), 129.3 (d), 131.4 (d), 131.8 (d), 132.1 (d), 132.4 (d), 134.7 (s), 135.8 (s), 137.1 (s), 139.1 (s); UV-vis (CHCl₃) λ_{max} 417 (ϵ 3.6 × 10³), 308 (sh, 1.4 × 10³), 289 (sh, 9.8 × 10²); HRMS (FAB; m/z) found 650.9342 ([M - Cl]⁺), calcd for $C_{30}H_{22}{}^{35}Cl^{106}PdS_4$ ([M - Cl]⁺) 650.9328.

X-ray Structural Analysis of 1·CHCl₃, 2, 7·2(CH₂Cl₂), and 8. Single crystals of 1·CHCl₃, 2, 7·2(CH₂Cl₂), and 8 suitable for X-ray structural analysis were obtained by slow recrystallization from CHCl₃/CH₃CN, benzene, CH₂Cl₂/CH₃CN, and CH₂Cl₂/EtOH, respectively. The crystals were mounted on a glass fiber. The intensity data were collected on Rigaku/MSC Mercury CCD diffractometer with graphite-monochromated Mo K α radiation (λ = 0.710 69 Å for 1·CHCl₃ and λ = 0.710 70 Å for 2, 7·2(CH₂Cl₂), and 8). The structures were solved by direct methods $(SIR-97)^{24}$ (for 1·CHCl₃, 2, and 8) and Patterson methods (DIRDIF)²⁵ (for 7.2CH₂Cl₂) and refined by full-matrix least-squares procedures on F^2 for all reflections (SHELXL-97).²⁶ All the non-hydrogen atoms were refined anisotropically. All hydrogens for 1. CHCl3 and 7.2(CH₂Cl₂) were placed using AFIX instructions, and those for 2 and 8 were refined isotropically. The structural data are shown in Table 3. In the analysis of 7.2CH₂Cl₂, there are the highest peak $(1.559 \text{ e} \text{ Å}^{-3})$ and the deepest hole $(-1.634 \text{ e} \text{ Å}^{-3})$ near the CH₂Cl₂ solvent molecule due to disorders that could not be completely solved.

Theoretical Calculations. All theoretical calculations were carried out using the Gaussian 98 program²⁷ with density functional theory at the B3LYP level.²⁰ The LANL2DZ basis sets for Pd were used with effective core potential, and the 6-31G(d) basis sets were used for C, H, S, and Cl atoms. The LANL2DZ(ECP) basis sets were obtained from the Extensible Computational Chemistry Environment Basis Set Database (http://www.emsl.pnl.gov/forms/ basisform.html), version 02/25/04, as developed and distributed by

- (25) Beurskens, P. T.; Beurskens, G.; de Gelder, R.; Garcia-Granda, S.; Gould, R. O.; Israel, R.; Smits, J. M. M. *The DIRDIF-99 program system*; Crystallography Laboratory, University of Nijmegen: Nijmegen, The Netherlands, 1999.
- (26) Sheldrick, G. M. SHELX-97, Program for the Refinement of Crystal Structures; University of Göttingen: Göttingen, Germany, 1997.
- (27) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Zakrzewski, V. G.; Montgomery, J. A., Jr.; Stratmann, R. E.; Burant, J. C.; Dapprich, S.; Millam, J. M.; Daniels, A. D.; Kudin, K. N.; Strain, M. C.; Farkas, O.; Tomasi, J.; Barone, V.; Cossi, M.; Cammi, R.; Mennucci, B.; Pomelli, C.; Adamo, C.; Clifford, S.; Ochterski, J.; Petersson, G. A.; Ayala, P. Y.; Cui, Q.; Morokuma, K.; Salvador, P.; Dannenberg, J. J.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Cioslowski, J.; Ortiz, J. V.; Baboul, A. G.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Andres, J. L.; Gonzalez, C.; Head-Gordon, M.; Replogle, E. S.; Pople, J. A. *Gaussian 98*; Gaussian, Inc.: Pittsuburgh, PA, 1998.

⁽²³⁾ Beringer, F. M.; Kravetz, L.; Topliss, G. B. J. Org. Chem. 1965, 30, 1141–1148.

⁽²⁴⁾ Altomare, A.; Burla, M. C.; Camalli, M.; Cascarano, G. L.; Giacovazzo, C.; Guagliardi, A.; Moliterni, A. G. G.; Polidori, G.; Spagna, R. J. Appl. Crystallogr. 1999, 32, 115–119.

Table 3. Crystal Data and Refinement Details for 1. CHCl₃, 2, 7. 2CH₂Cl₂, and 8

param	1-CHCl ₃	2	7 •2(CH ₂ Cl ₂)	8		
empirical formula	C72H130S4Si12•CHCl3	$C_{30}H_{22}S_4$	C72H130Cl2PdS4Si12+2CH2Cl2	C ₃₀ H ₂₂ Cl ₂ PdS ₄		
fw	1580.45	510.72	1808.23	688.02		
temp (K)	103(2)	103(2)	103(2)	103(2)		
cryst system	monoclinic	triclinic	monoclinic	triclinic		
space group	<i>P</i> 2 ₁ (No. 4)	<i>P</i> 1 (No. 2)	<i>C</i> 2/ <i>c</i> (No. 15)	<i>P</i> 1 (No. 2)		
$a(\text{\AA})$	14.0803(6)	7.5404(10)	41.273(3)	9.4221(18)		
<i>b</i> (Å)	16.5775(7)	10.4513(16)	12.8906(6)	10.779(2)		
<i>c</i> (Å)	19.9887(9)	16.759(3)	19.3198(11)	14.826(3)		
α (deg)	90	88.513(6)	90	80.988(7)		
β (deg)	90.613(3)	78.037(5)	105.752(2)	76.200(7)		
γ (deg)	90	71.784(4)	90	73.758(5)		
$V(Å^3)$	4665.4(3)	1226.2(3)	9892.9(10)	1397.5(5)		
Ζ	2	2	4	2		
D_{calc} (Mg m ⁻³)	1.125	1.383	1.214	1.635		
abs coeff (mm ^{-1})	0.377	0.406	0.615	1.174		
cryst size (mm)	$0.40 \times 0.25 \times 0.03$	$0.40 \times 0.20 \times 0.10$	$0.35 \times 0.20 \times 0.10$	$0.20 \times 0.10 \times 0.05$		
θ range (deg)	1.78-25.50	2.36-26.00	1.91-25.00	2.63-25.00		
no. of reflcns measd	27 097	8636	39 657	9167		
no. of indep reflens	14 849	4661	8658	4828		
R _{int}	0.0834	0.0126	0.0219	0.0191		
completeness (%)	98.90	96.70	99.20	97.80		
data/restraints/params	14 849/1/865	4661/0/395	8658/0/456	4828/0/422		
goodness-of-fit on F^2	1.005	1.051	1.079	1.092		
final R indices $[I > 2\sigma(I)]^a$	$R_1 = 0.0636$	$R_1 = 0.0306$	$R_1 = 0.0549$	$R_1 = 0.0290$		
	$wR_2 = 0.1540$	$wR_2 = 0.0738$	$wR_2 = 0.1393$	$wR_2 = 0.0571$		
R indices (all data) ^a	$R_1 = 0.0930$	$R_1 = 0.0332$	$R_1 = 0.0576$	$R_1 = 0.0349$		
	$wR_2 = 0.1757$	$wR_2 = 0.0754$	$wR_2 = 0.1414$	$wR_2 = 0.0594$		
absolute struct param	-0.10(9)					
largest diff peak and hole (e $Å^{-3}$)	0.576 and -0.661	0.537 and -0.269	1.559 and -1.634	0.600 and -0.298		
${}^{a}\mathbf{R}_{1} = \Sigma F_{0} - F_{c} / \Sigma F_{0} , \mathbf{W}\mathbf{R}_{2} = [(\Sigma w (F_{0}{}^{2} - F_{c}{}^{2})^{2} / \Sigma w (F_{0}{}^{2})^{2}]^{1/2}.$						

the Molecular Science Computing Facility, Environmental and Molecular Sciences Laboratory, which is part of the Pacific Northwest Laboratory, P.O. Box 999, Richland, WA 99352, and funded by the U.S. Department of Energy. The Pacific Northwest Laboratory is a multiprogram laboratory operated by Battelle Memorial Institute for the U.S. Department of Energy under Contract DE-AC06-76RLO 1830. Contact Karen Schuchardt for further information. Computation time was provided by the Supercomputer Laboratory, Institute for Chemical Research, Kyoto University.

Acknowledgment. This work was partially supported by Grants-in-Aid for COE Research on Elements Science (No. 12CE2005), Creative Scientific Research (17GS0207), Scientific Research on Priority Area (No. 14078213), Young Scientist (B) (No. 15750031), and 21st Century COE Program on Kyoto University Alliance for Chemistry, from the Ministry of Education, Culture, Sports, Science, and Technology of Japan. We are grateful to Assistant Professor Takahiro Sasamori, and Dr. Noriyoshi Nagahora, Institute for Chemical Research, Kyoto University, for the valuable discussions.

Supporting Information Available: Crystallographic information files (CIF) for the compounds 1·CHCl₃, 2, 7·2(CH₂Cl₂), and 8, atomic coordinates for the calculated molecules 8a,b, VT-¹³C DEPT 90 NMR spectra of 8 (at 50, 25, 0, -20, -40, and -60 °C), and cyclic voltammograms of 7–9. This material is available free of charge via the Internet at http://pubs.acs.org.

IC050944V