CS_2 insertion into a gold–carbon bond. First syntheses and characterization of 2,2-diacetylethylene-1,1-dithiolato complexes. Crystal structure of $[N(PPh_3)_2][Au\{\eta^2-S_2C=C(COMe)_2\}_2]^{\dagger}$

José Vicente,*a⁺ María Teresa Chicote,*a Pablo González-Herrero^a and Peter G. Jones*b^b

^a Grupo de Química Organometálica,¶ Departamento de Química Inorgánica, Facultad de Química, Universidad de Murcia, Aptdo. 4021, E-30071 Murcia, Spain

^b Institut für Anorganische und Analytische Chemie, Technische Universität Braunschweig, Postfach 3329, 38023 Braunschweig, Germany

The reaction of $[N(PPh_3)_2][Au(acac)_2]$ (acac = acetylacetonate) with CS₂ gives $[N(PPh_3)_2]_2[Au_2\{\mu^2: \eta^2: S_2C=C(COMe)_2\}_2]$, which reacts with PhICl₂ to give $[N(PPh_3)_2][Au\{\eta^2:S_2C=C(COMe)_2\}_2]$, for which the crystal structure is determined; these are the first characterized 2,2-diacetylethylene-1,1-dithiolato complexes of any element.

Interest in gold complexes containing S-bonded ligands stems from their potential application in the glass and ceramic industries¹ and, more importantly, in medicine (chrysotherapy).² Thiolatogold(I) complexes such as the commercial Myocrisin, Allochrysine, Solganol, or Auranofin, are among the most efficient antiarthritic drugs.² In addition, Solganol and Auranofin show *in vitro* inhibitory effects on Human Immunodeficiency Virus 1,³ high cytotoxicity to tumor cells⁴ and activity against i.p. P388 leukaemia.⁵

We have found that CS₂ reacts with $[N(PPh_3)_2][Au(acac)_2]^6$ to give $[N(PPh_3)_2]_2[Au_2\{\mu^2:\eta^2-S_2C=C(COMe)_2\}_2]$ 1 (Scheme 1). As far as we are aware, this is the first fully characterized 2,2-diacetylethylene-1,1-dithiolato complex of any element.

In the absence of any mechanistic data, we assume that the CS_2 inserts into one of the gold–carbon bonds to produce a dithiocarboxylato complex **a** (see Scheme 1), likely to be in tautomeric equilibrium with the ethylene-1,1-dithiolato form **b**, the remaining acac ligand being responsible for intermolecular deprotonation and subsequent dimerization to give **1** with concomitant formation of acetylacetone.

Insertion reactions of CS₂ into transition-metal–H, –C, –N, –P, –S, or –Cl bonds are known to produce dithioformiato, dithiocarboxylato, dithiocarboxylato, phosphinodithiocarboxylato, trithio- or perthio-carbonato, or chlorodithioformiato

complexes, respectively.7 However, in the case of gold these reactions have scarcely been studied and thus only two other CS_2 insertion reactions have been described so far, namely those into the Au–C bond in $[Au{\eta^1-C_5Me_5}(PR_3)]$ complexes give dithiocarboxylato complexes $[Au{\eta^1}$ to $S_2CC_5Me_5)(PR_3)_n$ (R = Ph, n = 2; R = Prⁱ, n = 3)⁸ or into an Au-Cl bond of [Au2Cl6] to give the chlorodithioformate complex [AuCl₂(η^2 -S₂CCl)].⁹ We have ourselves described the insertion of CS₂ into an Au-S (or S-H) bond in [N(PPh₃)₂][Au(SH)₂] to produce the first trithiocarbonatogold complex.¹⁰ CS_2 with the reaction of In Me)CS₂ $_2$ [Au^I(C₆F₅)₂], resulting from a carbon–carbon coupling reaction, is obtained.11

The reaction of methylene active compounds with CS₂ and LiNR₂, NaH,¹² or K₂CO₃,¹³ led to the synthesis of the corresponding 2,2-disubstituted ethylene-1,1-dithiolate salts, which were treated in situ with MeI to give ketene dithioacetals. In this context acetylacetone has been used to prepare 2-acetyl-2-oxopropylidene-S, S-acetal [(MeS)₂C=Cthe $\{C(O)Me\}_2$.^{13–15} Our synthetic method is, therefore, unprecedented. Ketene dithioacetals are of current interest because the double carbon-carbon bond they present is amenable to both nucleophilic and electrophilic attack.14 Indeed, they are very versatile starting materials for the synthesis of a wide variety of fused heterocycles,13,16 polysubstituted pyridines,17 pyrazoles,18 etc. On the other hand, the interest in ketene dithioacetals with electron-withdrawing substituents stems from some of their physical properties as push-pull polarized ethylenes. Thus, the effect of chiral substituents or hydrogen bonding on second-order non-linearity,19 the free energy of activation for rotation around the carbon-carbon double



Scheme 1 Q = $[N(PPh_3)_2]$, aca = acetylacetonate. *Reagents and conditions*: i, CS₂ (excess) in acetone, 1 h, room temp; ii, -2 Hacac; iii, + Cl₂IPh - PhI - Q[AuCl₂], CH₂Cl₂, 45 min, room temp.



Fig. 1 Structure of one of the two independent anions of 2. Selected bond lengths (Å) and angles (°): Au(1)–S(1) 2.3234(13), 2.3375(13), Au(1)–S(2) 2.3479(12), 2.3217(12), C(1)–S(1) 1.758(4), 1.748(4), C(1)–S(2) 1.744(4), 1.755(4), C(1)-C(4) 1.353(6), 1.365(6); S(1)-Au(1)-S(2) 74.16(4), 74.46(4), S(2)-Au(1)-S(1A) 105.84(4), 105.54(4), S(1)-C(1)-S(2) 107.0(2), 107.2(2), S(1)-C(1)-C(4) 125.8(3), 126.1(3), S(2)-C(1)-C(4) 127.2(3), 126.8(4),

bond,15,20 or the influence of the double bond twisting on the position of the IR $v_{C=C}$ stretching band²¹ or on the photoionization energy²² have been studied. Our synthesis opens a new access for such studies. The only 2,2-diacetylethylene-1,1-dithiolato complexes (of Mn^{II}, Co^{II}, Ni^{II}, Cu^{II} and Zn^{II}) mentioned in the literature were prepared from the corresponding sodium salt of the ligand and their composition was only established by elemental analysis, while amperometric and potentiometric titrations were unsuccessful.23

Many dinuclear gold(I) complexes are known to undergo oxidative addition reactions with halogens, pseudohalogens or alkyl halides to give gold(II) complexes containing a gold-gold bond.²⁴ However, when complex **1** is treated with an equimolar amount of Cl₂IPh in dichloromethane a disproportionation reaction takes place, giving [N(PPh₃)₂][AuCl₂] along with the gold(III) complex $[N(PPh_3)_2][Au\{\eta^2-S_2C=C(COMe)_2\}_2]$ 2. The same behaviour has previously been observed for other dithiolato,²⁵ dialkyldithiocarbamato,²⁶ or trithiocarbonato²⁷ gold(I) complexes.

The isolated complexes 1 and 2 have been fully characterized by C, H, N, S analyses, IR spectroscopy and ¹H and ¹³C NMR. The X-ray crystal structure of complex 2 (Fig. 1), \parallel shows it to crystallize with two independent half-anions and one $[N(PPh_3)_2]$ cation in the asymmetric unit. Both anions show almost identical inversion-symmetric $[Au{\eta^2} S_2C=C(COMe)_2_2$ E,Z-conformations. The gold atoms are in distorted square-planar environments, the angles around gold differing from the ideal value of 90° because of the small bite of the dithiolate ligand $[S(1)-Au(1)-S(2) 74.16(4), 74.46(4)^{\circ}; S(1)-Au(1)-S(2) 105.84(4), 105.54(4)^{\circ}, respectively].$ The Au(1), S(1), S(2), C(1) and C(2) atoms are coplanar (mean deviation 0.054, 0.044 Å), with both acetyl groups twisted out of this plane, one of them considerably (torsion angle 55, 48°), the other one slightly (torsion angle 9, 13°, respectively). The C(1)-C(4) bond distances [1.353(6), 1.365(4) Å] lie in the range expected for $C(sp^2)=C(sp^2)$ double bonds $(1.294-1.392)^{28}$ and are similar to that found in the closely related ketene S,Sdithioacetal (MeS)₂C=C{C(O)Me}₂ (1.351 Å) which shows a twisted (torsion angle 14°) E,E-conformation.²⁹

Apart from bands due to the cation, the solid-state IR spectrum of 1 shows two strong v(C=O) bands at 1698 and 1678 cm^{-1} and a strong v(C=C) band at 1580 cm^{-1} , while the v(AuS)band(s) cannot be assigned unequivocally because of the presence of bands from the $[N(PPh_3)_2]$ cation in the 400–300 cm^{-1} region. The position of the v(C=O) band suggest that conjugation plays at best a marginal role.²⁹ The IR spectrum of **2** shows two strong v(C=O) bands at 1664 and 1636 cm⁻¹ and a medium v(AuS) band at 370 cm⁻¹. The frequency of the v(C=C) band is expected to decrease because of conjugation with one of the acetyl groups and cannot be unequivocally assigned.

We thank DGICYT (PB92-0982C) and the Fonds der Chemischen Industrie for financial support. P. G.-H. is grateful to Ministerio de Educación y Cultura for a Grant.

Footnotes and References

† Dedicated to Professor Pascual Royo on the occasion of his 60th birthday

- ‡ E-mail: jvs@fcu.um.es
- § E-mail: p.jones@tu-bs.de
- WWW: http://www.scc.um.es/gi/gqo/

 $\|$ Crystal data for complex 2: C₄₈H₄₂ÅuNO₄P₂S₄, triclinic, space group $P\overline{1}$, T = 143 K, a = 10.729(3), b = 11.853(3), c = 18.076(4) Å, $\alpha = 83.13(2)$, $β = 81.62(2), γ = 79.75(2)^\circ, U = 2227.6(10) Å³, Z = 2, λ(Mo-Kα) = 0.71073 Å, μ = 3.61 mm⁻¹, D_c = 1.616 Mg m⁻³. Data collection:$ Yellow prism $0.55 \times 0.40 \times 0.30$ mm, Stoe STADI-4 diffractometer, 9522 intensities of which 7867 independent, $2\theta_{max}$ 50°, absorption corrections based on ψ -scans with transmissions 0.776–0.933. Structure solution and refinement: heavy-atom method, refined anisotropically on F2 (program SHELXL-93, G.M. Sheldrick, University of Göttingen). H atoms with riding model or as rigid methyl groups. In the C(=O)Me groups of the second anion, the slightly short C–C and long C=O bonds and the difficulty in locating methyl H may indicate some disorder. Final wR(F2) 0.0742, R(F) 0.0296, 547 parameters, 464 restraints, S = 1.09, max $\Delta \rho 1.4$ e Å⁻³. CCDC 182/591.

- 1 A. N. Papazian, Gold Bull., 1982, 15, 81.
- 2 A. Lorber and T. M. Simon, Gold Bull., 1979, 12, 149; K. C. Dash and H. Schmidbaur, Metal Ions in Biological Systems, ed. H. Sigel, Marcel Dekker, New York, 1982; P. J. Sadler, Adv. Inorg. Chem., 1991, 36, 1.
- 3 T. Okada, B. K. Patterson, S.-Q. Ye and M. E. Gurney, Virology, 1993, 192. 631.
- 4 T. M. Simon, D. H. Kunishima, D. H. Vibert and A. Lorber, Cancer Res., 1981, 41, 94.
- 5 C. K. Mirabelli, R. K. Johnson, C. M. Sung, L. Faucette, K. Muirhead and S. T. Crooke, Cancer Res., 1985, 45, 32.
- 6 J. Vicente, M. T. Chicote, I. Saura-Llamas and M. C. Lagunas, J. Chem. Soc. Chem. Commun., 1992, 915.
- 7 K. K. Pandey, Coord. Chem. Rev., 1995, 140, 37; V. Cadierno, M. P. Gamasa, J. Gimeno and E. Lastra, J. Organomet. Chem., 1996, 510, 207.
- H. Otto and H. Werner, Chem. Ber., 1987, 120, 97.
- D. Jentsch, P. G. Jones, C. Thöne and E. Schwarzmann, J. Chem. Soc. Chem. Commun., 1989, 1495.
- 10 J. Vicente, M. T. Chicote, P. González-Herrero and P. G. Jones, J. Chem. Soc. Chem. Commun., 1995, 745.
- 11 M. Bardají, M. C. Gimeno, P. G. Jones, A. Laguna, M. Laguna, F. Merchán and I. Romeo, Organometallics, 1997, 16, 1083.
- 12 Z.-T. Huang and X. Shi, Synthesis, 1990, 162.
- 13 A. K. El-Shafei, A. M. M. El-Saghier and E. A. Ahmed, Synthesis, 1994, 152.
- 14 D. Villemin and A. B. Alloum, Synthesis, 1991, 301.
- 15 J. Sandström and I. Wennerbeck, Acta Chem. Scand., 1970, 24, 1191. 16 L.-B. Wang and Z.-T. Huang, Synth. Commun., 1996, 26, 459;
- Z.-T. Huang and X. Shi, Chem. Ber., 1990, 123, 541. 17 F. A. Abu-Shanab, M. H. Elnagdi, F. M. Ali and B. J. Wakefield,
- J. Chem. Soc., Perkin Trans. 1, 1994, 1449.
- 18 E. C. Taylor and P. W. Ronald, Heterocycles, 1977, 8, 243.
- 19 K. Mohanalingam, M. Nethaji and P. K. Das, J. Mol. Struct., 1996, 378, 177.
- 20 J. Sandström and U. Sjöstrand, Tetrahedron, 1978, 34, 3305.
- 21 D. Smith and P. J. Taylor, J. Chem. Soc., Perkin Trans. 2, 1979, 1376.
- 22 F. P. Colonna, G. DiStefano, J. Sandström and U. Sjöstrand, J. Chem. Soc. Perkin Trans 2, 1978, 279.
- 23 W. U. Malik, R. Bembi and V. K. Bharddwaj, J. Indian Chem. Soc., 1984, 65, 379.
- 24 J. P. Fackler Jr., Polyhedron, 1997, 16, 1.
- 25 M. N. I. Khan, S. Wang and J. P. Fackler Jr., Inorg. Chem., 1989, 28, 3579.
- 26 D. C. Calabro, B. A. Harrison, G. T. Palmer, M. K. Moguel, R. L. Rebbert and J. L. Burmeister, Inorg. Chem., 1981, 20, 4311.
- 27 J. Vicente, M. T. Chicote, P. González-Herrero and P. G. Jones, unpublished work.
- 28 F. H. Allen, O. Kennard, D. G. Watson, L. Brammer, A. G. Orpen and R. Taylor, J. Chem. Soc., Perkin Trans. 2, 1987, S1.
- 29 A. J. Jarvis and P. J. Taylor, J. Chem. Soc., Perkin Trans. 2, 1978, 972.

Received in Cambridge, UK, 15th July 1997; 7/05052F

2048 Chem. Commun., 1997