

Ruthenium–Acetylide-Mediated Catalytic Dimerization of $\text{RC}\equiv\text{CH}$ ($\text{R} = \text{Ph}$, CO_2Me) and the Formation of the New Ruthenium η^3 -Butadienyl Complex $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)[\eta^3\text{-PhCHCHC}=\text{C}(\text{Ph})\text{C}\equiv\text{CPh}]$

Chae S. Yi* and Nianhong Liu

Department of Chemistry, Marquette University, Milwaukee, Wisconsin 53201-1881

Arnold L. Rheingold and Louise M. Liable-Sands

Department of Chemistry, The University of Delaware, Newark, Delaware 19716

Received May 1, 1997[®]

The *in-situ*-generated ruthenium–acetylide species $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)\text{C}\equiv\text{CPh}$ (**1**) was found to catalyze the dimerization of $\text{RC}\equiv\text{CH}$ to give predominantly the head-to-head dimers *trans*- $\text{RCH}=\text{CHC}\equiv\text{CR}$ ($\text{R} = \text{Ph}$ (**3**), $\text{TON} = 607 \text{ h}^{-1}$; $\text{R} = \text{CO}_2\text{Me}$ (**4**), $\text{TON} = 975 \text{ h}^{-1}$). A new ruthenium η^3 -butadienyl complex $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)(\eta^3\text{-PhCHCHC}=\text{C}(\text{Ph})\text{C}\equiv\text{CPh})$ (**5**) was isolated at the end of the catalytic dimerization of $\text{PhC}\equiv\text{CH}$ (81% based on Ru). The structure of the 1.5 benzene-solvated complex **5**·1.5 C_6H_6 was established by X-ray crystallography. Complex **5** was independently prepared from the stoichiometric reaction of $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)(\text{Cl})=\text{C}=\text{CHPh}$ (**2**) with the dimer **3** in the presence of NEt_3 . The formation of an intermediate η^2 -alkyne complex $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)(\eta^2\text{-PhCH}=\text{CHC}\equiv\text{CPh})(\text{C}\equiv\text{CPh})$ (**6**) was observed during the reaction.

Introduction

The transition-metal-mediated dimerization of terminal alkynes is of considerable current interest because it can lead to a wide variety of organic enyne and oligoacetylene products that are useful synthetic precursors for organic conducting polymers and other carbon-rich allotropes.^{1–4} For example, the copper–acetylide catalysts have been employed in an industrial-scale dimerization of $\text{HC}\equiv\text{CH}$ to produce $\text{CH}_2=\text{CH}-\text{C}\equiv\text{CH}$, a key precursor for neoprene rubber.^{1a} Recently, transition-metal-based catalysts have been shown to selectively produce both the 1,3-disubstituted enynes² and the 1,4-disubstituted enynes³ from the dimerization

of terminal alkynes. The stereoselective formation of *cis*-1,4-disubstituted butatrienes has also been achieved from the dimerization of alkyl-substituted alkynes.⁴ We recently reported the ruthenium-catalyzed dimerization of terminal alkynes to produce both 1,3- and 1,4-disubstituted enynes and butatrienes and proposed a mechanism involving a coordinatively unsaturated ruthenium–acetylide species.⁵ We also developed an effective route to the coordinatively unsaturated acetylide species $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)\text{C}\equiv\text{CPh}$ (**1**) from the reaction of the vinylidene complex $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)(\text{Cl})=\text{C}=\text{CHPh}$ (**2**) with a base and found that the ruthenium acetylide **1** was reactive toward a variety of small molecules such as CO , H_2 , and CO_2 .⁶ Herein, we report the ruthenium–acetylide-mediated catalytic dimerization of terminal alkynes and the formation of a new (η^3 -butadienyl)-ruthenium complex.

[®] Abstract published in *Advance ACS Abstracts*, August 1, 1997.
 (1) For recent reviews and monographs, see: (a) Parshall, G. W.; Ittel, S. D. *Homogeneous Catalysis*, 2nd ed.; John Wiley & Sons: New York, 1992. (b) Henkelmann, J. In *Applied Homogeneous Catalysis with Organometallic Compounds*; Cornils, B., Herrmann, W. A., Eds.; VCH: New York, 1996. (c) *Modern Acetylene Chemistry*; Stang, P. J., Diederich, F., Eds.; VCH: New York, 1995. (d) Farina, V. In *Comprehensive Organometallic Chemistry II*; Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Pergamon Press: New York, 1994; Vol. 12. (e) Sonogashira, K. In *Comprehensive Organic Synthesis*; Trost, B. M., Ed.; Pergamon Press: New York, 1990; Vol. 2. (f) Kanis, D. R.; Ratner, M. A.; Marks, T. J. *Chem. Rev.* **1994**, *94*, 195. (g) *Materials for Nonlinear Optics: Chemical Perspectives*; Stucky, G. D., Marder, S. R., Sohn, J., Eds.; ACS Symposium Series 455; American Chemical Society: Washington, DC, 1991. (h) Lang, H. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 547.
 (2) For leading examples of 1,3-enynes, see: (a) Thompson, M. E.; Baxter, S. M.; Bulls, A. R.; Burger, B. J.; Nolan, M. C.; Santarsiero, B. D.; Schaefer, W. P.; Bercaw, J. E. *J. Am. Chem. Soc.* **1987**, *109*, 203. (b) Trost, B. M.; Chan, C.; Ruhter, G. *J. Am. Chem. Soc.* **1987**, *109*, 3486. (c) Ohshita, J.; Furumori, K.; Matsuguchi, A.; Ishikawa, M. *J. Org. Chem.* **1990**, *55*, 3277 and references therein. (d) Boese, W. T.; Goldman, A. S. *Organometallics* **1991**, *10*, 782. (e) Horton, A. D. *J. Chem. Soc., Chem. Commun.* **1992**, 185. (f) Barluenga, J.; González, J. M.; Llorente, I.; Campos, P. J. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 893. (g) Straub, T.; Haskel, A.; Eisen, M. S. *J. Am. Chem. Soc.* **1995**, *117*, 6364. (h) Trost, B. M.; Sorum, M. T.; Chan, C.; Harms, A. E.; Ruhter, G. *J. Am. Chem. Soc.* **1997**, *119*, 698 and references cited therein.

(3) For leading examples of 1,4-enynes, see: (a) Ohshita, J.; Furumori, K.; Matsuguchi, A.; Ishikawa, M. *J. Org. Chem.* **1990**, *55*, 3277. (b) Bianchini, C.; Peruzzini, M.; Zanobini, F.; Frediani, P.; Albinati, A. *J. Am. Chem. Soc.* **1991**, *113*, 5453. (c) Jun, C.-H.; Lu, Z.; Crabtree, R. H. *Tetrahedron Lett.* **1992**, *33*, 7119. (d) Duchateau, R.; van Wee, C. T.; Meetsma, A.; Teuben, J. H. *J. Am. Chem. Soc.* **1993**, *115*, 4931. (e) Evans, W. J.; Keyer, R. A.; Ziller, J. W. *Organometallics* **1993**, *12*, 2618. (f) Bianchini, C.; Frediani, P.; Masi, D.; Peruzzini, M.; Zanobini, F. *Organometallics* **1994**, *13*, 4616. (g) Matsuzaka, H.; Takagi, Y.; Ishii, Y.; Nishio, M.; Hidai, M. *Organometallics* **1995**, *14*, 2153. (h) Bianchini, C.; Innocenti, P.; Peruzzini, M.; Romerosa, A.; Zanobini, F. *Organometallics* **1996**, *15*, 272. (i) Slugovc, C.; Mereiter, K.; Zobetz, E.; Schmid, R.; Kirchner, K. *Organometallics* **1996**, *15*, 5275.
 (4) For leading examples of 1,4-disubstituted butatrienes, see: (a) Wakatsuki, Y.; Yamazaki, H.; Kumegawa, N.; Satoh, T.; Satoh, J. Y. *J. Am. Chem. Soc.* **1991**, *113*, 9604. (b) Schäfer, M.; Mahr, N.; Wolf, J.; Werner, H. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 1315. (c) Heeres, H. J.; Nijhoff, J.; Teuben, J. H.; Rogers, R. D. *Organometallics* **1993**, *12*, 2609. (d) Wakatsuki, Y.; Koga, N.; Yamazaki, H.; Morokuma, K. *J. Am. Chem. Soc.* **1994**, *116*, 8105.
 (5) Yi, C. S.; Liu, N. *Organometallics* **1996**, *15*, 3968.
 (6) Yi, C. S.; Liu, N.; Rheingold, A. L.; Liable-Sands, L. M.; Guzei, I. A. *Organometallics* **1997**, *16*, 3729.





Figure 2. Stoichiometric reaction profile of **2** (10 mg, 0.016 mmol) with $\text{PhC}\equiv\text{CH}$ (3 μL , 0.032 mmol, 2 equiv) and Et_3N (4 μL , 0.032 mmol, 2 equiv) in CD_2Cl_2 , as monitored by ^1H NMR at 20 $^\circ\text{C}$. For clarity, only complexes **2** (\square), **5** (\blacksquare), **6** (\blacktriangle), and the dimer **3a** (\bullet) are shown.

Scheme 1



expense of both complexes **6** and **3a**. Interestingly, no notable change in concentration of the *cis* dimer **3b** was observed during the entire reaction period.

The fact that **5** was generated even after most of $\text{PhC}\equiv\text{CH}$ was converted to dimer **3** and that **5** was formed at the expense of both **6** and dimer **3** suggested that **5** was obtained from the coupling reaction of **1** with **2** (Scheme 1). In support of this hypothesis, reaction of **2** with <1 equiv of $\text{PhC}\equiv\text{CH}$ produced a relatively long-lived complex **6**, as in this case, it could not be converted to **5** in the absence of **3**. Also, complex **5** was formed independently from a stoichiometric reaction of **2** (500 mg, 0.79 mmol) with $\text{PhC}\equiv\text{CH}$ (0.86 mL, 7.9 mmol) and Et_3N (1.1 mL, 7.9 mmol) in 81% isolated yield. The *syn* geometry between the $-\text{C}\equiv\text{CPh}$ group and the PPh_3 ligand of **5** was also consistent with the intramolecular migration of the acetylide to the coordinated enyne **3** via the alkyne intermediate **6**.

The formation of 1,4-disubstituted enynes **3** can be rationalized readily from a migratory insertion of the coordinated $\text{RC}\equiv\text{CH}$ ($\text{R} = \text{Ph}, \text{CO}_2\text{Me}$) to the ruthenium-acetylide and the subsequent reductive coupling with another alkyne substrate.^{1b,3} While the formation of the 1,4-enynes has been previously explained by invoking an acetylene-to-vinylidene rearrangement and the sub-

sequent acetylide migration,^{3,5} the direct insertion pathway seems to be preferred for alkynes with electron-withdrawing groups at relatively low temperature, where the acetylene-to-vinylidene tautomerization is relatively slow.^{1c,2} The formation of η^3 -butadienyl complex **5** can be rationalized similarly by invoking the intramolecular migratory insertion of the enyne **3** via the alkyne-coordinated **6** in which the acetylide ligand is selectively added to the phenyl-substituted acetylenic carbon. Since the acetylide migration has been commonly known to occur at the most electrophilic alkyne substrate,^{1c,d} a relatively more electron-withdrawing vinyl group of **3** should facilitate the acetylide migration to a relatively more electrophilic phenyl-substituted alkyne carbon.

In summary, the ruthenium-acetylide **1** was shown to be an effective catalyst for the dimerization of $\text{RC}\equiv\text{CH}$ ($\text{R} = \text{Ph}, \text{CO}_2\text{Me}$). A new ruthenium η^3 -butadienyl species **5** was formed from the reaction of **1** with the dimeric product **3** via an intermediate **6**. Further studies on the ruthenium-acetylide-mediated coupling reactions of alkenes and alkynes are currently in progress.

Experimental Section

General Information. All reactions were carried out in an inert-atmosphere glovebox or by using standard high-vacuum and Schlenk-line techniques unless otherwise mentioned. Tetrahydrofuran, benzene, and Et_2O were distilled from purple solutions of sodium and benzophenone immediately prior to use. The NMR solvents, C_6D_6 and CD_2Cl_2 , were dried from activated molecular sieves (4 Å). The ruthenium complexes, $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)_2\text{Cl}$ ¹⁰ and $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)(\text{Cl})=\text{C}=\text{CHPh}$ (**2**),^{6,11} were prepared according to literature procedures. Organic alkynes $\text{RC}\equiv\text{CH}$ ($\text{R} = \text{Ph}, \text{CO}_2\text{Me}$) were received from a commercial source (Aldrich Chemical Co.) and used without further purification. The ^1H and ^{13}C NMR spectra were recorded on a GE GN-Omega 300 MHz FT-NMR spectrometer. Mass spectra were recorded on a Hewlett-Packard HP 5970 GC/MS spectrometer. Elemental analyses were performed at the Midwest Microlab, Indianapolis, IN.

Formation of $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)(\eta^3\text{-PhCHCHC}=\text{C-PhC}=\text{CPh})$ (5**).** The ruthenium complex $\text{C}_5\text{Me}_5\text{Ru}(\text{PPh}_3)(\text{Cl})=\text{C}=\text{CHPh}$ (**2**) (500 mg, 0.79 mmol), Et_3N (1.10 mL, 10 equiv), and $\text{PhC}\equiv\text{CH}$ (0.86 mL, 10 equiv) were charged with 30 mL of CH_2Cl_2 in a 100 mL round-bottom flask. The reaction mixture was stirred at room temperature for 24 h. After the solvent was evaporated under high vacuum, the residue was dissolved in C_6H_6 and the solution filtered through a frit. The filtrate was evaporated under vacuum, and the remaining solid was washed with small amounts of Et_2O (3×5 mL) and hexanes (3×5 mL) to afford **5** as a yellow solid in 81% yield (513 mg).

^1H NMR (CD_2Cl_2 , 300 MHz): δ 7.8–7.0 (m, Ph), 3.68 (dd, $J_{\text{HH}} = 8.8$ Hz, $J_{\text{PH}} = 3.7$ Hz, PhCHCH), 3.00 (dd, $J_{\text{PH}} = 14.0$ Hz, $J_{\text{HH}} = 8.8$ Hz, PhCHCH), 1.36 (s, C_5Me_5). $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_2Cl_2 , 75 MHz): δ 192.6 (d, $J_{\text{PC}} = 17.1$ Hz, $\text{Ru}-\text{C}=\text{C}$), 144.8, 143.3, 133.2, 130.0, 129.3, 129.0, 128.8, 127.8, 127.0, 126.0 (Ph carbons), 125.3 ($=\text{C}(\text{Ph})\text{C}=\text{C}$), 114.8 (d, $J_{\text{PC}} = 6.4$ Hz, CHPh), 96.4 ($\text{CHPh}=\text{}$), 96.3 ($\text{C}=\text{CPh}$), 93.4 (C_5Me_5), 88.0 ($\text{C}=\text{CPh}$),

(10) (a) Serron, S. A.; Luo, L.; Li, C.; Cucullu, M. E.; Stevens, E. D.; Nolan, S. P. *Organometallics* **1995**, *14*, 5290. (b) Luo, L.; Nolan, S. P.; Fagan, P. J. *Organometallics* **1993**, *12*, 4305. (c) Conroy-Lewis, F. M.; Simpson, S. J. *J. Organomet. Chem.* **1987**, *322*, 221.

(11) (a) Bianchini, C.; Casares, J. A.; Peruzzini, M.; Romerosa, A.; Zanolini, F. *J. Am. Chem. Soc.* **1996**, *118*, 4585. (b) Bianchini, C.; Innocenti, P.; Peruzzini, M.; Romerosa, A.; Zanolini, F. *Organometallics* **1996**, *15*, 272. (c) Slugovc, C.; Mereiter, K.; Zobetz, E.; Schmid, R.; Kirchner, K. *Organometallics* **1996**, *15*, 5275.

(9) Selected spectral data for **6**: ^1H NMR (CD_2Cl_2 , 300 MHz) δ 6.73 (d, $J = 15.9$ Hz, $\text{CH}=\text{CHPh}$), 6.50 (d, $J = 15.9$ Hz, $\text{CH}=\text{CHPh}$), 1.44 (s, C_5Me_5); $^{31}\text{P}\{^1\text{H}\}$ NMR (CD_2Cl_2 , 121.6 MHz) δ 53.7 (s, PPh_3).

10.8 (C₅Me₅). ³¹P{¹H} NMR (CD₂Cl₂, 121.6 MHz): δ 60.9 (s, PPh₃). Anal. Calcd for C₅₂H₄₇PRu: C, 77.68; H, 5.89. Found C, 77.65; H, 6.05.

General Procedure of the Ruthenium-Catalyzed Dimerization of RC≡CH (R = Ph, CO₂Me). In a 25 mL Schlenk tube equipped with a Teflon stopcock, the ruthenium catalyst **1** (3–5 mol %) was generated from the treatment of C₅Me₅Ru(PPh₃)(Cl)=C≡CHPh (10 mg, 0.016 mmol) with Et₃N (22 μL, 10 equiv) in 10 mL of CH₂Cl₂. Excess alkyne (0.60 mmol) was added to the solution, and the reaction mixture was stirred at room temperature. Small samples were periodically drawn out from the solution and analyzed by the GC-MS. After 10 h, the solution was evaporated under high vacuum. The residue was extracted with Et₂O, and the Et₂O solution was chromatographed on silica gel (hexane/Et₂O) in air. The rotary evaporation led to the dimeric products as a pale-yellow oil. The spectroscopic data for both **3a** and **3b** have been previously reported.⁵

For *trans*-PhCH=CHC≡CPh (**3a**): ¹H NMR (C₆D₆, 300 MHz) δ 8.10–6.80 (m, Ph), 7.04 (d, *J* = 16.2 Hz, =CHPh), 6.30 (d, *J* = 16.2 Hz, =CHC≡C); ¹³C{¹H} NMR (C₆D₆, 75 MHz) δ 142.3 (=CHPh), 137.3, 132.2, 129.7, 129.4, 129.2, 127.3, 124.4 (Ph carbons), 109.0 (=CHC≡C), 92.3 (=CHC≡C), 89.8 (=CHC≡CPh); GC-MS *m/z* 204 (M⁺).

For *cis*-PhCH=CHC≡CPh (**3b**): ¹H NMR (C₆D₆, 300 MHz) δ 8.10–6.80 (m, Ph), 6.40 (d, *J* = 11.8 Hz, =CHPh), 5.79 (d, *J* = 11.8 Hz, =CHC≡C); ¹³H{¹H} NMR (C₆D₆, 75 MHz) δ 139.1 (=CHPh), 137.1, 131.7, 129.2, 128.7, 128.5, 128.4, 124.0 (Ph carbons), 107.7 (=CHC≡C), 96.7 (=CHC≡CPh), 88.9 (=CHC≡CPh); GC-MS *m/z* 204 (M⁺).

For *trans*-MeO₂CCH=CHC≡CCO₂Me (**4**): ¹H NMR (C₆D₆, 300 MHz) δ 6.45 (d, *J* = 16.2 Hz, =CHCO₂Me), 6.03 (d, *J* = 16.2 Hz, =CHC≡C), 3.23, 3.21 (s, CO₂Me); ¹³C{¹H} NMR (C₆D₆, 75 MHz) δ 164.6 (=CHCO₂Me), 153.4 (C≡CCO₂Me), 135.1 (=CHCO₂Me), 121.3 (=CHC≡CCO₂Me), 87.3 (C≡CCO₂Me), 81.9 (C≡CCO₂Me), 52.6, 51.5 (CO₂Me); GC-MS *m/z* 168 (M⁺).

X-ray Crystallographic Determination of 5. Bright yellow single crystals of **5** were obtained from the layering of hexanes to a benzene solution at room temperature. A yellow crystal (0.30 × 0.30 × 0.20 mm) was mounted on a Siemens P4/CCD diffractometer equipped with a graphite monochromator. Empirical methods were used to correct for absorption. No symmetry greater than triclinic was evident from the diffraction data. The asymmetric unit consists of one Ru complex and 1.5 molecules of benzene. The half molecule is located on a crystallographic inversion center. All non-hydrogen atoms were treated as idealized contributions. All software used were included in the SMART, SAINT, and SHELXTL program libraries (Siemens XRD, Madison, WI). Crystal data for **5**: C₅₂H₄₇PRu·1.5C₆H₆; yellow block, triclinic, *a* = 10.2582(1) Å, *b* = 15.8273(2) Å, *c* = 16.9688(2) Å, α = 101.0450(1)°, β = 107.5713(2)°, γ = 108.5520(1)°, *V* = 2360.54(4) Å³, *Z* = 2, *T* = 223 K, *R*(*F*) = 0.0508, *R*(w*F*²) = 0.1046.

Acknowledgment. Financial support from Marquette University, Committee on Research, is gratefully acknowledged. The authors from The University of Delaware acknowledge the National Science Foundation for their support in purchasing the CCD-based X-ray diffractometer (Grant No. CHE-9628768).

Supporting Information Available: Tables of crystal data and structure refinement, atomic coordinates, bond lengths and angles, anisotropic displacement parameters, and hydrogen coordinates for **5** (9 pages). Ordering information is given on any current masthead page.

OM970366T