

Pd(II)-Catalyzed Carbonylation of C(sp³)-H Bonds: A New Entry to 1,4-Dicarbonyl Compounds

Eun Jeong Yoo, Masayuki Wasa, and Jin-Quan Yu*

Department of Chemistry, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037, United States

Received September 10, 2010; E-mail: yu200@scripps.edu

Abstract: Pd(II)-catalyzed β -C(sp³)-H carbonylation of *N*-arylamides under CO (1 atm) has been achieved. Following amide-directed C(sp³)-H cleavage and insertion of CO into the resulting [Pd(II)-C(sp³)] bond, intramolecular C-N reductive elimination gave the corresponding succinimides, which could be readily converted to 1,4-dicarbonyl compounds. This method was found to be effective with substrates containing α -hydrogen atoms and could be applied to effect methylene C(sp³)-H carbonylation of cyclopropanes.

The development of diverse transformations using Pd catalysis to functionalize unactivated β -C(sp³)-H bonds in aliphatic acids and their amide derivatives could offer a new strategy for synthesizing molecules with complex tertiary and quaternary carbon centers.¹⁻³ To broaden the scope of this approach, we have reported a variety of novel C-C and C-heteroatom bond-forming reactions.^{1,3b} We envisioned that a reaction to effect β -carbonylation of C(sp³)-H bonds would be a highly desirable addition to our collection of transformations because replacement of C-H bonds by a highly oxidized carbonyl group installs a versatile handle for further structural elaboration. For example, β -carbonylation of aliphatic acids would provide a novel route to synthetically important 1,4-dicarbonyl compounds^{4,5} that are widespread in biologically important natural products (Figure 1).⁶






Figure 1. Natural products that contain 1,4-dicarbonyl moiety.

Despite landmark developments in Pd(0)-catalyzed carbonylation of aryl halides, triflates, and tosylates⁷ and recent progress on Pd(II)-catalyzed carbonylation of aryl C(sp²)-H bonds (eq 1),⁸⁻¹¹ Pd(II)-catalyzed C(sp³)-H carbonylation reactions that use the substrate as a limiting reagent have yet to be reported. Herein, we disclose a protocol for Pd(II)-catalyzed β -C(sp³)-H carbonylation of aliphatic amides to give succinimides which are readily hydrolyzed to afford broadly useful 1,4-dicarbonyl compounds (eq 2). The succinimide intermediates are also well-established synthons for preparation of *syn*-disubstituted 1,4-dicarbonyl compounds through regioselective and diastereoselective enolate chemistry (eq 2).¹² TEMPO (2,2,6,6-tetramethylpiperidine-1-oxyl) was found to be a crucial co-oxidant for efficient reoxidation of Pd(0) to Pd(II) in the presence of CO.

Previous work (ref 10a):




This work:



Early reports by Fujiwara described a Pd-catalyzed C(sp³)-H carbonylation reaction of alkanes (used in excess) under CO (20 to 50 atm) using K₂S₂O₈ as the reoxidant in TFA at 80 °C, which gave a regioisomeric mixture of the corresponding carboxylic acid products.¹³ The use of excess substrate and the lack of regioselectivity limited the usefulness of this chemistry for synthetic applications. Although regioselective C(sp²)-H carbonylation using 1 equiv of substrate under CO (1 atm) has been accomplished⁸⁻¹¹ through the use of proximate directing groups, extending this approach to C(sp³)-H carbonylation represents a significant challenge. One major obstacle is that excess CO inhibits the activation of the inert C(sp³)-H bonds by competitively occupying coordination sites on the Pd(II) center, thereby preventing the requisite C-H agostic interaction. In addition, both the insertion of CO into a [Pd(II)-C(sp³)] cyclopalladated intermediate and reductive elimination from Pd involving a C(sp³)-CO moiety have a limited number of precedents.¹⁴

With these considerations in mind, we initiated our investigation of Pd-catalyzed C(sp³)-H carbonylation by first identifying a highly efficient directing group for facile C(sp³)-H activation. Recently, our group established that acidic amides are superior directing groups for promoting C-H activation reactions with both Pd(0)/PR₃ and Pd(II) catalysts.^{1c,d} We therefore focused our efforts on using acidic amide substrates for our exploratory studies (Table 1).

Gratifyingly, we observed that **1a** could be transformed into the desired carbonylation product **2a** in 30% yield (based on ¹H NMR), using Pd(OAc)₂ (10 mol %) as the catalyst, AgOAc (2 equiv) as the oxidant, and KH₂PO₄ as the base, under CO (1 atm) in *n*-hexane at 130 °C (Table 1, entry 4). Following 1,1-migratory insertion to forge the C(sp³)-CO bond, the Pd intermediate undergoes Pd-

Table 1. Optimization of Reaction Conditions for Pd-Catalyzed C(sp³)-H Carbonylation^a


entry	additive	solvent	yield (%) ^b	entry	additive	solvent	yield (%) ^b
1	none	DMF	<1	6	Cu(OAc) ₂	<i>n</i> -hexane	4
2	none	toluene	7	7	DMF	<i>n</i> -hexane	54
3	none	C ₆ F ₆	8	8	PivOH	<i>n</i> -hexane	50
4	none	<i>n</i> -hexane	30	9	TEMPO ^c	<i>n</i> -hexane	80
5	BQ	<i>n</i> -hexane	13	10	TEMPO	<i>n</i> -hexane	95

^a Reaction conditions: **1a** (0.1 mmol), Pd(OAc)₂ (10 mol %), AgOAc (2.0 equiv), KH₂PO₄ (2.0 equiv), additive (2.0 equiv), solvent (1 mL), CO (1 atm), 130 °C, 18 h. ^b The yield was determined by ¹H NMR analysis of the crude product using 1,1,2,2-tetrachloroethane as an internal standard. ^c TEMPO (0.2 equiv).


mediated C–N reductive elimination to give the corresponding succinimide products.

After surveying a wide array of organic solvents, we observed that *n*-hexane gave the best reactivity. This finding was unexpected, as *n*-hexane has rarely been an effective solvent in our previous C–H activation reactions.¹ Several additives that are known to positively influence Pd-catalyzed C–H functionalization reactions (e.g., 1,4-benzoquinone and Cu(OAc)₂) were tested; however, they proved to be incompatible with the reaction conditions (entries 5 and 6). Among the other additives tested, DMF (2 equiv) increased conversion to 54% (entry 7), and pivalic acid improved the yield to 50% (entry 8). Further screening of the additives revealed that the addition of a catalytic amount of TEMPO (0.2 equiv) as a co-oxidant with AgOAc dramatically improved the conversion to over 80% (entry 9). Although the precise role of TEMPO remains to be elucidated, one plausible explanation is that oxoammonium salt¹⁵ (the oxidized form of TEMPO) reoxidizes Pd(0) to Pd(II) more efficiently than solely AgOAc. A stoichiometric amount of both TEMPO and AgOAc is required to achieve full conversion; using 2.0 equiv of TEMPO and AgOAc increased the yield to 95% (entry 10), while using 2.0 equiv of TEMPO gave only 12% conversion (see Supporting Information (SI)).

Notably, other directing groups that have previously been utilized in our laboratory for C(sp³)-H activation, such as carboxylic acids, hydroxamic acids, oxazolines, and pyridines, were unreactive under these conditions. An analogous acidic directing group *N*-toluenesulfonyl amide (CONHTs),¹⁶ however, gives a corresponding product in lower yield (63%, see SI).

With the optimized conditions in hand, we converted a range of commercial aliphatic acids into the corresponding CONHAr amides to examine the scope of this carbonylation protocol (Table 2). Substrates with a quaternary α -carbon atom gave good to excellent yields of the succinimide products (**2a–h**). Products containing ether groups (**2d–f** and **2k**) could also be obtained in good yields. The benzyl moiety proved to be a better protecting group than the TIPS group for β -hydroxyl substrates (**1e** and **1f**), while TBS-protected substrates gave none of the desired product. Notably, this method was also effective for the carbonylation of methylene C–H bonds in cyclopropane substrates (**2g**, **2h**, and **2l**). Intriguingly, cyclopropyl C(sp³)-H bonds could be selectively carbonylated over a methyl C(sp³)-H bond (in **1g**) and an *ortho* aryl C(sp²)-H bond (in **1h**).

In our early efforts to develop C(sp³)-H functionalization reactions with aliphatic carboxylic acids and their derivatives, substrates containing α -hydrogen atoms were often unreactive,

Table 2. Substrate Scope for C(sp³)-H Carbonylation^{a,b}



2a , 91%	2b , 99%	2c , 96%	2d , 96%
2e , 92%	2f , 70%	2g , 86%	2h , 65% ^c
2i , 65%	2j , 60% ^d	2k , 52%	2l , 65%

^a Reaction conditions: amide substrate **1** (0.1 mmol), Pd(OAc)₂ (10 mol %), TEMPO (2.0 equiv), AgOAc (2.0 equiv), KH₂PO₄ (2.0 equiv), *n*-hexane (1 mL), CO (1 atm), 130 °C, 18 h. ^b Isolated yield. ^c Ar' = C₆F₅. ^d Pd(OAc)₂ (20 mol %).

restricting the substrate scope to those containing quaternary α -carbon atoms. We were pleased to find that, in the present study, carbonylated products **2i–l** could be obtained in acceptable yields from substrates containing α -hydrogen atoms. These types of products are highly valuable synthons for the preparation of 1,4-dicarbonyl compounds.^{4–6}

To demonstrate the synthetic utility of this reaction, succinimide product **2a** was subjected to two different ring-opening conditions to obtain either 1,4-dicarboxylic acid **3** or 1,4-dicarbonyl molecule **4** (Scheme 1). Upon treatment of **2a** with TFA/AcOH under reflux, hydrolysis occurs to give 2,2-dimethylsuccinic acid **3** in 93% isolated yield. Similarly, treatment of **2a** with NaOMe in methanol at room temperature gave 80% of the ester product **4** without concomitant hydrolysis of the amide moiety; thus this group could potentially be used to direct further elaboration of the *gem*-dimethyl unit *via* Pd-catalyzed C–H functionalization.

Scheme 1. Ring-Opening of Succinimides



In summary, we have developed a novel protocol to effect carbonylation of C(sp³)-H bonds under CO (1 atm). Studies to expand the scope of the reaction to simple carboxylic acid substrates and to develop an enantioselective variant for substrates containing *gem*-dimethyl or cyclopropyl groups are currently underway in our laboratory.

Acknowledgment. We gratefully acknowledge The Scripps Research Institute, the National Institutes of Health (NIGMS, 1 R01 GM084019-02), the NSF (NSF CHE-1011898), Amgen, and Eli Lilly for financial support. This work was supported by a National

Research Foundation of Korea Grant funded by the Korean Government (NRF-2009-352-C00077). We also thank Bristol Myers Squibb for a predoctoral fellowship (M.W.).

Supporting Information Available: Experimental procedures and spectral data for all new compounds. This material is available free of charge *via* the Internet at <http://pubs.acs.org>.

References

- (1) (a) Giri, R.; Mangel, N.; Li, J.-J.; Wang, D.-H.; Breazzano, S. P.; Saunders, L. B.; Yu, J.-Q. *J. Am. Chem. Soc.* **2007**, *129*, 3510. (b) Wang, D.-H.; Wasa, M.; Giri, R.; Yu, J.-Q. *J. Am. Chem. Soc.* **2008**, *130*, 7190. (c) Wasa, M.; Engle, K. M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2009**, *131*, 9886. (d) Wasa, M.; Engle, K. M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 3680.
- (2) Renaudat, A.; Jean-Gerard, L.; Jazsar, R.; Kefalidis, C. E.; Clot, E.; Baudoin, O. *Angew. Chem., Int. Ed.* **2010**, *49*, 7261.
- (3) For early studies on β -C(sp³)-H functionalization reactions of carboxylic acids using strongly coordinating auxiliaries, see: (a) Zaitsev, V. G.; Shabashov, D.; Daugulis, O. *J. Am. Chem. Soc.* **2005**, *127*, 13154. (b) Giri, R.; Chen, X.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2005**, *44*, 2112.
- (4) For examples of a classic disconnection involving conjugate addition of acyl anions, see: (a) Corey, E. J.; Hegedus, L. S. *J. Am. Chem. Soc.* **1969**, *91*, 4926. (b) Stetter, H.; Schreckenberger, M. *Angew. Chem., Int. Ed.* **1973**, *12*, 81. (c) Groebel, B.-T.; Seebach, D. *Synthesis* **1977**, 357. (d) Kubota, Y.; Nemoto, H.; Yamamoto, Y. *J. Org. Chem.* **1991**, *56*, 7195.
- (5) For recent reports describing the synthesis of 1,4-dicarbonyl compounds, see: (a) Jang, H.-Y.; Hong, J.-B.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2007**, *129*, 7004. (b) DeMartino, M. P.; Chen, K.; Baran, P. S. *J. Am. Chem. Soc.* **2008**, *130*, 11546. (c) Liu, Q.; Perreault, S.; Rovis, T. *J. Am. Chem. Soc.* **2008**, *130*, 14066.
- (6) (a) Omura, S.; Hirano, A.; Iwai, Y.; Masuma, R. *J. Antibiot.* **1979**, *32*, 786. (b) Furusaki, A.; Matsumoto, T.; Ogura, H.; Takayanagi, H.; Hirano, A.; Omura, S. *J. Chem. Soc., Chem. Commun.* **1980**, 698. (c) Li, S.-H.; Wang, J.; Niu, X.-M.; Shen, Y.-H.; Zhang, H.-J.; Sun, H.-D.; Li, M.-L.; Tian, Q.-E.; Lu, Y.; Cao, P.; Zheng, Q.-T. *Org. Lett.* **2004**, *6*, 4327.
- (7) (a) *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E., de Meijere, A., Eds.; Wiley: New York, 2002; Vol. 2, p 2309. (b) Munday, R. H.; Martinelli, J. R.; Buchwald, S. L. *J. Am. Chem. Soc.* **2008**, *130*, 2754. (c) Barnard, C. F. J. *Organometallics* **2008**, *27*, 5402. (d) Brennfürer, A.; Neumann, H.; Beller, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 4114.
- (8) For lactam formation *via* C-H carbonylation, see: Orito, K.; Horibata, A.; Nakamura, T.; Ushito, H.; Nagasaki, H.; Yuguchi, M.; Yamashita, S.; Tokuda, M. *J. Am. Chem. Soc.* **2004**, *126*, 14342.
- (9) For the use of diethyl azodicarboxylate for C-H carbonylation, see: Yu, W.-Y.; Sit, W. N.; Lai, K.-M.; Zhou, Z.; Chan, A. S. C. *J. Am. Chem. Soc.* **2008**, *130*, 3304.
- (10) For Pd-catalyzed C-H carbonylation with CO, see: (a) Giri, R.; Yu, J.-Q. *J. Am. Chem. Soc.* **2008**, *130*, 14082. (b) Houlden, C. E.; Hutchby, M.; Bailey, C. D.; Ford, J. G.; Tyler, S. N. G.; Gagne, M. R.; Lloyd-Jones, G. C.; Booker-Milburn, K. I. *Angew. Chem., Int. Ed.* **2009**, *48*, 1830. (c) Giri, R.; Lam, J. K.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 686. (d) Wu, X.-F.; Anbarasan, P.; Neumann, H.; Beller, M. *Angew. Chem., Int. Ed.* **2010**, *49*, 7316.
- (11) For carbonylation reactions of C(sp²)-H bonds using other metals, see: (a) Guan, Z.-H.; Ren, Z.-H.; Spinella, S. M.; Yu, S.; Liang, Y.-M.; Zhang, X. *J. Am. Chem. Soc.* **2009**, *131*, 729. (b) Inoue, S.; Shiota, H.; Fukumoto, Y.; Chatani, N. *J. Am. Chem. Soc.* **2009**, *131*, 6898.
- (12) (a) Bennett, D. J.; Pickering, P. L.; Simpkins, N. S. *Chem. Commun.* **2004**, 1392. (b) Pohlmann, J.; Lampe, T.; Shimada, N.; Nell, P. G.; Pernerstorfer, J.; Svenstrup, N.; Brunner, N. A.; Schiffer, G.; Freiberg, C. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 1189.
- (13) (a) Nishiguchi, T.; Nakata, K.; Takaki, K.; Fujiwara, Y. *Chem. Lett.* **1992**, *21*, 1141. (b) Miyata, T.; Nakata, K.; Yamaoka, Y.; Taniguchi, Y.; Takaki, K.; Fujiwara, Y. *Chem. Lett.* **1993**, *22*, 1005. (c) Nakata, K.; Yamaoka, Y.; Miyata, T.; Taniguchi, Y.; Takaki, K.; Fujiwara, Y. *J. Organomet. Chem.* **1994**, *473*, 329.
- (14) (a) Balavoine, G.; Clinet, J. C. *J. Organomet. Chem.* **1990**, *390*, C84. (b) Dangel, B. D.; Godula, K.; Youn, S. W.; Sezen, B.; Sames, D. *J. Am. Chem. Soc.* **2002**, *124*, 11856. (c) Bloome, K. S.; Alexanian, E. J. *J. Am. Chem. Soc.* **2010**, *132*, 12823.
- (15) (a) Semmelhack, M. F.; Schmid, C. R.; Cortés, D. A.; Chou, C. S. *J. Am. Chem. Soc.* **1984**, *106*, 3374. (b) Kochkar, H.; Lassalle, L.; Morawietz, M.; Hölderich, W. F. *J. Catal.* **2000**, *194*, 343. (c) Kirchberg, S.; Vogler, T.; Studer, A. *Synlett* **2008**, 2841.
- (16) Inamoto, K.; Saito, T.; Hiroya, K.; Doi, T. *J. Org. Chem.* **2010**, *75*, 3900.

JA108754F