

Facile synthesis and biological assays of novel 2,4-disubstituted hydrazinyl-thiazoles analogs

Fateme Ghanbari Pirbasti¹ · Nosrat O. Mahmoodi¹

Received: 23 July 2015 / Accepted: 29 December 2015
© Springer International Publishing Switzerland 2016

Abstract A convenient, one-pot multi-component synthesis of new 2,4-disubstituted hydrazinyl-thiazoles was accomplished using different aldehydes/ketones, thiosemicarbazide, and 4-methoxy phenacyl bromide in the presence of a catalytic amount of AcOH in EtOH. Products were obtained in reasonable yields and high purity. The in vitro antioxidant activity of hydrazinyl-thiazoles was evaluated by DPPH radical scavenging activity in comparison to ascorbic acid. Synthesized thiazoles **14c** and **14g** possessed the lowest IC₅₀ values. Also, hydrazinyl-thiazoles were screened for their in vitro antibacterial activity against six strains of bacteria including *S. aureus*, *M. luteus*, *E. coli*, *Ps. aeruginosa*, *B. subtilis*, and *A. hydrophila* where some products showed good antibacterial activity. Moreover, compound **14a** showed anticancer activity against melanoma cancerous cell lines A375 with LC₅₀ = 0.55 mg/mL, slightly selective versus normal cell lines (Hu-2) with LC₅₀ = 1.19 mg/mL.

Keywords One-pot three-component reaction · Disubstituted hydrazinyl-thiazoles · Antioxidant · DPPH · Antibacterial · Anticancer · MCR

Introduction

Thiazoles have attracted a great deal of attention because of their diverse biological activities. Thiamine (vitamin B₁, **1**)

Electronic supplementary material The online version of this article (doi:10.1007/s11030-015-9654-7) contains supplementary material, which is available to authorized users.

✉ Nosrat O. Mahmoodi
nosmahmoodi@gmail.com; mahmoodi@guilan.ac.ir

¹ Department of Chemistry, Faculty of Science,
University of Guilan, P.O. Box 41335-1914, Rasht, Iran

is one of the B-complex vitamins that contains a thiazole ring, which is vitally important to keep a living organism operating properly. Commercial drugs, such as sulfathiazole **2** (antimicrobial drug), ritonavir **3** (antiretroviral drug), and abafungin **4** (antifungal drug) contain a thiazole moiety (Fig. 1).

Synthesized compounds containing a thiazole moiety have been used for the treatment of allergies [1], hypertension [2], inflammation [3], human immunodeficiency virus (HIV) infections [4], and schizophrenia [5]. Thiazoles also exhibit other biological activities, such as antibacterial [6,7], antitumor [8], analgesic [9], anticonvulsant [10], antimelanogenesis [11], antipsychotic [12], and antioxidant [13]. Also, thiazoles can act as cyclin-dependent kinase (CDK) inhibitors [14] and β -glucuronidase inhibitors [15].

Analogs such as 2-substituted-6-fluorobenzo[d]thiazoles [16], (thiazol-2-yl)hydrazine derivatives [17], 3-Allyl-2-(substituted imino)-4-phenyl-3*H*-thiazoles, and 2,2'-(1,3-phenylene)bis(3-substituted-2-imino-4-phenyl-3*H*-thiazole) derivatives [18] showed anti-*candida*, antibacterial, and acetylcholinesterase activities. Moreover, thiazoles have been used as semiconducting materials [19], thiazole-iridium complexes as phosphorescent organic light-emitting diodes [20], and naphthol-substituted thiazoles as ratiometric fluorescent chemosensors [21].

Recently, our research group reported the one-pot synthesis of new thiazolyl-pyrazoline derivatives **5–6** [22], bis-thiazoles **7–8** [23], and thiazolylpyridazinones **9** [24] exhibiting antibacterial activity, and new 1,4-dihydropyridines **10** bearing a thiazole moiety exhibiting high antioxidant activity [25] (Fig. 2). In continuation of our previous investigation [26], we report herein the one-pot MCR synthesis of new hydrazinyl-thiazole analogs and their biological activities.




Fig. 1 Vitamin B₁ and drugs containing a thiazole moiety




Fig. 2 Thiazoles accessed via one-pot synthesis

Results and discussion

Chemistry

New 2,4-disubstituted hydrazinyl-thiazoles **14a–g** were synthesized via the one-pot reaction of different aldehydes/ketones **11a–g**, thiosemicarbazide **12**, and 4-methoxy phenacyl bromide **13** in the presence of AcOH in EtOH under reflux conditions (Scheme 1). A literature survey indicated that hydrazone-thiazoles are generally synthesized in a two-step procedure. To the best of our knowledge, this is the first one-pot synthesis report for **14** analogs [27–30].

At the onset of our research, we explored the one-pot, multi-component preparation of **14a** in EtOH at room temperature in the absence of a catalyst (entry 1) which after 6h afforded the desired product in 56 % yield. In the presence of AcOH, the yield of **14a** increased from 56 to 60 % (entry 2). Then, the scope of the reaction was investigated under reflux conditions. As shown in Table 1, **14a** was obtained after 2h in EtOH or DMF in high yield (entries 4 & 8). Since DMF

is a toxic solvent, EtOH was chosen as it is an eco-friendly and green solvent.


Our results show that the products are obtained in high yields and are easily purified using non-chromatographic methods. The advantages of this method are simple set-up and product isolation, reproducibility, using cheap and eco-friendly solvent.

In most cases, upon addition of **13** to a solution of **12** in EtOH after 2 h under reflux condition with aldehydes/ketones **11a–g** in the presence of few drop of AcOH, the desired product precipitated. This is a good sign for the start of the reaction. Hydrazinyl-thiazoles **14a**, **14b**, and **14c** were obtained as pure solids without the need for further purification.

The structures of hydrazinyl-thiazoles **14a–g** were determined by IR, ¹H NMR, and ¹³C NMR spectra. In the IR spectra, the disappearance of the carbonyl and thiosemicarbazone N–H stretching vibration bands indicates the participation of thioamide moiety in the cyclization reaction formation of desired products. In ¹H NMR spectra, the presence of a singlet at 7.09–6.06 ppm confirms the thiazole ring closure. Also, the OCH₃ signal appeared as a

Table 1 Optimization of **14a**

Entry	Catalyst	Solvent	Temp	Time (h)	Yield % ^a
1	—	EtOH	r.t.	6	56
2	AcOH (20 mol%)	EtOH	r.t.	4	60
3	AcOH (20 mol%)	EtOH	Reflux (60 °C)	2	75
4	AcOH (20 mol%)	EtOH	Reflux (80 °C)	2	80
5	AcOH (20 mol%)	H ₂ O	Reflux	4	65
6	AcOH (20 mol%)	EtOH:H ₂ O (1:1)	Reflux	3	70
7	AcOH (20 mol%)	MeOH	Reflux	3	70
8	AcOH (20 mol%)	DMF	Reflux	2	80

^a After purification**Scheme 1** Synthesis of new thiazoles **14a–g**

singlet at 3.87–3.67 ppm. Other aromatic protons appeared in the expected region around 8.94–6.68 ppm. The N–H signal appeared as a broad singlet at 14.50–11.45 ppm. In addition, the N–H signal can be endo- or exocyclic. However, according to literature [31], the signal at 14.50–11.45 ppm is related to an exocyclic N–H. The ¹³C NMR spectra of **14a–g** provided the expected number and types of carbons. The C–H of the thiazole ring appeared at 100.4–98.7 ppm. The signal at 55.9–55.3 is attributed to OCH₃ and other aromatic carbons appeared at 112.6–171.5 ppm.

In order to apply this procedure to carbonyl compounds with steric hindrance, we investigated the 3MCRs of camphor and 2-adamantanone **15** with thiosemicarbazide **12**, and 4-methoxy phenacyl bromide **13**. The reaction of camphor was unsuccessful and produced several by-products. However, **15** led to mixture of products **16A–16D** in moderate yield (Scheme 2).

The ¹H NMR spectra did not show a single product; rather, it showed a mixture of two sets of stereoisomer **16A/B** and **C/D**. The protons of the adamantane moiety appeared at 2.20–1.25 ppm, OCH₃ of isomers **16C/D** appeared at expected region at 3.96–3.72 ppm, and the proton of the thiazole moiety as well as other aromatic protons appeared at 7.80–6.71 ppm. Several signals at 3.25–3.01 and 2.74–2.58 ppm corresponded to CH–N=N isomers **C** and **D**.

Biology

Antioxidant assay

Using DPPH (diphenylpicrylhydrazyl) is one of the simplest methods to evaluate antioxidant activity. DPPH is a stable, free radical of violet color. When a compound donates a radical hydrogen atom to a DPPH molecule, it reduces

Scheme 2 Synthesis of thiazoles **16A–D** from adamantanone and proposed structures





Fig. 3 DPPH radical scavenging activity of hydrazinyl-thiazoles **14a–g**




DPPH and so the absorbance of DPPH decreases. In brief, the higher the antioxidant activity, the lighter the violet hue. The antioxidant activity of hydrazinyl-thiazoles **14a–g** was evaluated by the DPPH radical scavenging activity method. In this method, a decrease in absorption band at 517 nm indicates that the test compound possesses antioxidant activity. The radical scavenging activity of **14a–g** was screened at a concentration of 4000–62.5 µg/mL and monitored at 517 nm. Also, IC_{50} values (the concentration of **14a–g** to scavenge 50 % of DPPH radical concentration) were calculated from line equation which can be obtained from Fig. 3. Ascorbic acid was used as standard. All hydrazinyl-thiazoles **14a–g** showed dose-dependent antioxidant activity (Fig. 3).

The IC_{50} values of **14a–g** are in the range 2.90–0.23 µM (Table 2). It is worth noting that hydrazinyl-thiazole **14c** has the lowest IC_{50} value (0.23 µM) while compound **14a** has the highest IC_{50} value (2.90 µM) compared to the IC_{50} value of ascorbic acid (antioxidant compound) (1.30 µM). The higher antioxidant activity is reflected in a lower IC_{50} . Compounds

14a–g were determined to exhibit potent radical scavenging activity in a DPPH assay with IC_{50} values in the following order **14c**, **14g**, **14b**, **14e**, ascorbic acid, **14f**, **14d**, and **14a** respectively.

According to the literature [32] and our recent report [25], the high antioxidant activity of **14a–g** is attributed to the presence of the thiazole ring. The N–H at the C₂ position of the thiazole ring can readily donate a hydrogen radical to DPPH radical and form a radical of the test compound. Therefore, the radical is highly resonance-stabilized through the thiazole ring and =C–N–NH–C moiety (Scheme 3).

In addition to the presence of the thiazole moiety, the high antioxidant activity of **14g** can be due to the presence of benzylic hydrogens. These benzylic hydrogens are easily converted to the stable benzyl radical by DPPH. Furthermore, the antioxidant activity of **14b** and **14c** can be due to the presence of aliphatic hydrogens which can form tertiary, secondary, and primary carbon radicals. As it is mentioned in Table 2, these compounds exhibit better activity than ascorbic acid. Therefore, the presence of benzylic and



Scheme 3 Proposed mechanism of radical scavenging activity

Table 2 IC₅₀ values for DPPH radical scavenging activity of hydrazinyl-thiazoles **14a–g**

Compound	DPPH assay IC ₅₀ (μM)
14a	2.90
14b	0.87
14c	0.23
14d	2.10
14e	0.88
14f	1.69
14g	0.28
Ascorbic acid	1.30

aliphatic radicals as well thiazole and =C–N–NH–C moieties possibly play an effective role in antioxidant activity (Scheme 3).

Antibacterial assay

Products **14a–g** were screened for their in vitro antibacterial activity against Gram-positive and Gram-negative bacteria strains including *Staphylococcus aureus* (*S. aureus*), *Micrococcus luteus* (*M. luteus*), *Escherichia coli* (*E. coli*), *Pseudomonas aeruginosa* (*Ps. aeruginosa*), *Bacillus subtilis* (*B. subtilis*), *Aeromonas hydrophila* (*A. hydrophila*) using a well-diffusion method. DMSO was used as negative control and showed no activity against the above bacterial strains. Penicillin G and Cefixime were used as positive controls (Table 3).

Hydrazinyl-thiazoles **14a–g** were evaluated for their antibacterial activity at a concentration of 4000 μg/mL in DMSO. The experiments were performed in triplicate. The results are presented as mean ± standard deviation in mil-

limeter. According to the results, only **14d** was active against all six bacterial strains, and **14f** had the highest antibacterial activity against *S. aureus* and *M. luteus*. Compound **14d** was more active against *E. coli* and also showed significant inhibition activity against *B. subtilis*. In addition, **14e** showed significant inhibition activity against *Ps. aeruginosa*. Only **14b** showed good antibacterial activity against *A. hydrophila*.


Some reports showed moderate antibacterial activity of hydrazinyl-thiazoles. For example, Bharti and co-workers [33] studied the antibacterial activity of several hydrazinyl-thiazole derivatives, finding that their thiazoles showed no antibacterial activity against *E. coli* and only a few were active against *Ps. aeruginosa*. However, in our study, **14c–e** had moderate antibacterial activity against *E. coli*, while **14e** good antibacterial activity against *Ps. aeruginosa*. Moreover, Lee et al. [34] studied the antibacterial activity of hydrazinyl-thiazoles on several bacterial strains. According to their report, none of their synthesized compounds had antibacterial activity against *Ps. aeruginosa*, and only a few of them had good activity against *S. aureus*, *B. subtilis*, and *E. coli*. Also, the hydrazinyl-thiazoles showed no antibacterial activity against *A. hydrophila*.

Anticancer assay

Melanoma is a cancer that develops in melanocytes (the pigment cells present in the skin). It is known that melanoma can spread to other parts of the body causing serious illness and death. Melanoma can be more serious than two other forms of skin cancers, e.g., basal-cell cancer (BCC) and squamous-cell cancer (SCC). The primary treatment for melanoma is surgery. There are a few drugs used in chemotherapy of melanoma cancer; however, none of them

Table 3 Antibacterial activity of hydrazinyl-thiazoles **14a–g** (4000 µg/mL) as zone of inhibition in millimeter

Antibacterial activity (mean ± SD) (mm)						
Compound	<i>S. aureus</i>	<i>M. luteus</i>	<i>E. coli</i>	<i>Ps. aeruginosa</i>	<i>B. subtilis</i>	<i>A. hydrophila</i>
14a	–	8.33 ± 0.57	–	7.00 ± 0.73	6.66 ± 0.57	–
14b	7.33 ± 1.15	–	9.66 ± 1.15	7.33 ± 1.52	7.66 ± 0.57	11.00 ± 1.0
14c	6.66 ± 0.57	7.66 ± 0.57	11.00 ± 1.0	9.66 ± 0.57	11.00 ± 1.0	–
14d	7.66 ± 0.57	8.66 ± 0.57	13.00 ± 1.0	9.33 ± 0.57	21.33 ± 0.57	7.66 ± 0.57
14e	8.66 ± 0.57	–	11.33 ± 0.57	17.33 ± 0.57	12.33 ± 1.15	–
14f	11.33 ± 1.52	10.33 ± 0.57	–	7.66 ± 0.57	–	–
14g	7.66 ± 1.15	–	–	–	7.66 ± 0.57	6.66 ± 0.57
DMSO ^a	–	–	–	–	–	–
Penicillin G ^b	23	55	45	24	33	48
Cefixime ^b	38	35	39	30	32	41

^a Negative control^b Positive control**Fig. 4** In vitro cytotoxic activity of **14a** on human melanoma cancer cell lines A375. Results are presented as the percentage of cell viability ± SD using the MTT protocol

contain hydrazinyl-thiazole moiety, e.g., dacarbazine and temozolomide contain imidazole and carboxamide moieties, respectively. An in vitro cytotoxicity study of **14a** was carried out on human melanoma cancer cell lines A375 and Hu-02 (normal human skin cell lines) obtained from the National Centre for Cell Science, Tehran, Iran. Cell viability of **14a** was evaluated using the colorimetric MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay protocol at concentrations of 0.5, 0.25, 0.125, 0.075, 0.037 mg/mL in DMSO. DMSO was used as negative control. The MTT method is based on the reduction of soluble MTT by mitochondrial reductase of the viable cells to insoluble purple crystals of formazan ((*E, Z*)-5-(4,5-dimethylthiazol-2-yl)-1,3-diphenylformazan). The time of exposure was 24 h and 48 h (Fig. 4). Results are reported as the percent of cell viability ± standard deviation. Statistical analysis demonstrated that **14a** had cytotoxic activity on Hu-02 (69.4 %) and A375 (58.8 %) cell lines at 24 h time of exposure and on Hu-02 (64.3 %) and A375 (51.2 %)

Table 4 The LC₅₀ values of cytotoxic effects of **14a** on melanoma cancer cell lines (A375) (mean ± SD)

Entry	LC ₅₀ (mg/mL)			
	Hu-02 (24 h)	A375 (24 h)	Hu-02 (48 h)	A375 (48 h)
14a	1.19 ± 0.12	0.55 ± 0.01	0.81 ± 0.06	0.47 ± 0.02

at 48 h time of exposure at a concentration of 0.5 mg/mL ($P \leq 0.05$).

The LC₅₀ values of **14a** are shown in Table 4. The LC₅₀ values are between 1.19 and 0.55 mg/mL. As it is shown, the LC₅₀ values on melanoma cancer cells (A375) are lower than on skin cells (Hu-02). However, increasing the time of exposure from 24 h to 48 h did not have a significant effect on the LC₅₀ values.

Conclusions

Several new hydrazinyl-thiazoles derivatives **14a–g** were synthesized via one-pot reaction of aldehydes/ketones **11a–g**, thiosemicarbazide **12**, and 4-methoxy phenacyl bromide **13** through a reliable procedure for high yields and high purity. Hydrazinyl-thiazoles **14a**, **14b**, and **14c** were obtained as pure solids without further purification. All products showed high antioxidant activity in comparison to ascorbic acid. Compounds **14b** and **14e** had the highest antioxidant activity; their IC₅₀ was less than that of ascorbic acid. Compounds **14a–g** showed low to moderate antibacterial activity. Compound **14d** possessed the highest antibacterial activity against *B. subtilis*. Compound **14a** had cytotoxic activity at a concentration of 0.5 mg/mL upon 24 h and 48 h time of exposure on melanoma cancer cell lines.

Experimental section

Materials and instruments

Starting materials, reagents, biological cultures, and solvents were obtained from commercial suppliers Fluka, Merck, Sd-fine, Quelab and were used without further purification. All reactions were monitored by silica gel-coated TLC plates (60 F₂₅₄ Merck). IR spectra were recorded on a Shimadzu IR-470 spectrophotometer in anhydrous KBr. ¹H NMR and ¹³C NMR spectra were recorded on a 400 MHz and 500 MHz Bruker spectrometers using DMSO-*d*₆ and CDCl₃ as solvents. Chemical shifts are expressed relative to TMS as singlet (s), doublet (d), triplet (t), multiple (m), doublet of doublet (dd), doublet of triplet (dt), triplet of triplet (tt), doublet of doublet of doublet (ddd), and quintet (quin). Coupling constants are expressed in Hertz (Hz). Elemental analyses were carried out on a Carlo-Erba EA1110 CNNO-S analyzer. Melting points were determined with a Mettler Fp5 apparatus, and were uncorrected. Absorbance in the antioxidant assay was recorded on an Unico 2100 spectrophotometer.

General procedure for the synthesis of hydrazinyl-thiazoles 14a–g

To a solution of thiosemicarbazide **12** (1 mmol) in EtOH, aldehydes/ketones **11a–g** (1 mmol) and a few drops of AcOH were added and this mixture was refluxed and stirred. After a few minutes (1–15 min), 4-methoxy phenacyl bromide **13** (1 mmol) was added and the reaction was refluxed and stirred until completion (1.5–2 h). Thin layer chromatography (TLC) was used to monitor the progress of the reaction (EtOAc:*n*-hexane 3:6). After completion of the reaction, the reaction mixture was cooled down to room temperature. The resulting solid was filtered and washed with or recrystallized in EtOH.

2-(2-(6-Methoxy-3,4-dihydronaphthalen-1(2H)-ylidene)hydrazinyl)-4-(4-methoxyphenyl)thiazole (**14a**):

Cream solid (0.3 g, 80 %); mp 233–236 °C (from EtOH); IR (KBr) ν_{\max} 3210 (stretch N–H), 3100 (stretch C–H aromatic), 2920, 2830 (stretch C–H aliphatic), 1610 (stretch C=N), 1570, 1500 (stretch C=C), 1250, 1240, 1045 (stretch C–O), 825, 760, 700 (OOP. C–H) cm^{−1}; *R*_f (83 %, hexane: EtOAc 6:3); ¹H NMR (400 MHz, CDCl₃) δ 12.46 (s, 1H, NH), 8.02 (*d*, *J* = 8.8 Hz, 1H), 7.67 (*d*, *J* = 8.0 Hz, 2H), 6.99 (*d*, *J* = 8.8 Hz, 2H), 6.82 (dd, *J* = 8.8, 2.8 Hz, 1H), 6.68 (*d*, *J* = 2.4 Hz, 1H), 6.06 (s, 1H), 3.84 (s, 3H), 3.83 (s, 3H), 2.90 (*t*, *J* = 6.4 Hz, 2H), 2.79 (*t*, *J* = 6.0 Hz, 2H), 2.02 (quin, *J* = 6.3 Hz, 2H) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 169.3, 161.2, 160.7, 154.0, 142.5, 127.1, 126.9, 123.7, 121.5, 114.7, 113.5, 112.6, 99.1, 55.4, 55.3, 29.7, 27.2, 21.6 ppm;

Anal. calcd. for C₂₁H₂₁N₃O₂S: C, 66.49, H, 5.56, N, 11.10. Found: C, 66.51; H, 5.59; N, 11.06 %.

4-(4-Methoxyphenyl)-2-(2-(4-phenylcyclohexylidene)hydrazinyl)thiazole (**14b**)

Orange solid (0.3 g, 85 %); mp 186–189 °C (from EtOH); IR (KBr) ν_{\max} 3210 (stretch N–H), 3020 (stretch C–H aromatic), 2920 (stretch C–H aliphatic), 1610 (stretch C=N), 1560, 1540, 1510 (stretch C=C), 1250, 1040 (stretch C–O), 820, 740, 690 (OOP. C–H) cm^{−1}; *R*_f (80 %, hexane: EtOAc 6:3); ¹H NMR (400 MHz, CDCl₃) δ 12.45 (s, 1H, NH), 7.67 (*d*, *J* = 8.8 Hz, 2H), 7.33 (*t*, *J* = 7.2 Hz, 2H), 7.26–7.22 (*m*, 3H), 7.01 (*d*, *J* = 8.8 Hz, 2H), 6.56 (s, 1H), 3.86 (s, 3H), 3.32 (*d*, *J* = 14.8 Hz, 1H), 2.88 (tt, *J* = 12, 3.3 Hz, 1H), 2.72 (*d*, *J* = 14.4 Hz, 1H), 2.47 (dt, *J* = 14, 4.9 Hz, 1H), 2.32 (dt, *J* = 14.1, 5.3 Hz, 1H), 2.27–2.17 (*m*, 2H), 1.79 (ddd, *J* = 25.5, 13.1, 4.3 Hz, 2H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ 170.1, 164.2, 161.5, 145.1, 140.8, 129.0, 127.5, 127.1, 127.0, 120.3, 115.4, 98.8, 55.8, 43.5, 35.2, 34.6, 33.3, 29.4 ppm; Anal. calcd. for C₂₂H₂₃N₃OS: C, 70.03; H, 6.11; N, 11.09. Found: C, 69.98; H, 6.15; N, 11.12 %.

2-(2-(Heptan-4-ylidene)hydrazinyl)-4-(4-methoxyphenyl)thiazole (**14c**)

Orange solid (0.26 g, 83 %); mp 127–131 °C (from EtOH); IR (KBr) ν_{\max} 3210 (stretch N–H), 3100 (stretch C–H aromatic), 2950, 2850 (stretch C–H aliphatic), 1610 (stretch C=N), 1560, 1540, 1505 (stretch C=C), 1250, 1020 (stretch C–O), 830 (OOP. C–H) cm^{−1}; *R*_f (82 %, hexane: EtOAc 6:3); ¹H NMR (400 MHz, CDCl₃) δ 12.39 (s, 1H, NH), 7.66 (*d*, *J* = 8.2 Hz, 2H), 6.99 (*d*, *J* = 7.8 Hz, 2H), 6.53 (s, 1H), 3.85 (s, 3H), 2.49 (*t*, *J* = 9 Hz, *J* = 8 Hz, 2H), 2.34 (*t*, *J* = 7.6 Hz, 2H), 1.70–1.61 (*m*, 4H), 1.11 (*t*, *J* = 7.2 Hz, 3H), 0.98 (*t*, *J* = 7.4 Hz, 3H) ppm; ¹³C NMR (125 MHz, DMSO-*d*₆) δ 170.2, 166.1, 161.5, 140.7, 127.5, 120.4, 115.3, 98.7, 55.8, 38.9, 33.8, 19.8, 19.7, 14.6, 14.1 ppm; Anal. calcd. for C₁₇H₂₃N₃OS: C, 64.35; H, 7.28; N, 13.21. Found: C, 64.30; H, 7.31; N, 13.17 %.

2-(2-((4-Chlorophenyl)(phenyl)methylene)hydrazinyl)-4-(4-methoxyphenyl)thiazole (**14d**)

Orange crystal (0.32 g, 78 %); mp 239–242 °C (from EtOH); IR (KBr) ν_{\max} 3120 (stretch C–H aromatic), 1610 (stretch C=N), 1580, 1560, 1540 (stretch C=C), 1250, 1030 (stretch C–O), 830, 740, 690 (OOP. C–H) cm^{−1}; *R*_f (73 %, hexane: EtOAc 6:3); ¹H NMR (400 MHz, CDCl₃) δ 12.07 (s, 1H, NH), 7.69 (*d*, *J* = 8.8 Hz, 2H), 7.66 (*d*, *J* = 8.4 Hz, 2H), 7.60 (dd, *J* = 7.8, 1.3 Hz, 2H), 7.49 (tt, *J* = 6.6, 1.8 Hz, 1H), 7.42 (tt, *J* = 8.8, 1.2, 2H), 7.35 (*d*, *J* = 8.8 Hz, 2H), 6.85 (*d*, *J* = 8.8 Hz, 2H), 6.63 (s, 1H), 3.85 (s, 3H) ppm; ¹³C

NMR (125 MHz, $\text{CDCl}_3/\text{DMSO}-d_6$) δ 170.0, 161.3, 137.2, 135.7, 131.3, 130.7, 130.6, 130.4, 129.7, 129.6, 128.9, 128.5, 128.3, 127.6, 115.1, 114.6, 100.4, 55.7 ppm; Anal. calcd. for $\text{C}_{23}\text{H}_{18}\text{ClN}_3\text{OS}$: C, 65.81; H, 4.29; N, 9.98. Found: C, 65.85; H, 4.33; N, 10.03 %.

4-(4-Methoxyphenyl)-2-(phenyl(pyridin-2-yl)methylene)hydrazinylthiazole (14e)

Red solid (0.27 g, 71 %); mp 215–220 °C (from EtOH); IR (KBr) ν_{max} 1605 (stretch C=N), 1550, 1510, 1480 (stretch C=C), 1240, 1050 (stretch C–O), 830, 760, 730, 700 (OOP. C–H) cm^{-1} ; R_f (74 %, hexane: EtOAc 6:3); ^1H NMR (400 MHz, CDCl_3) δ 14.51 (*s*, 1H, NH), 8.93 (*d*, $J = 3.6$ Hz, 1H), 7.83–7.77 (*m*, 3H), 7.64–7.60 (*m*, 2H), 7.52–7.39 (*m*, 5H), 6.97 (*d*, $J = 8.8$ Hz, 2H), 6.78 (*s*, 1H), 3.87 (*s*, 3H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ 171.5, 160.6, 150.2, 148.6, 138.7, 136.0, 129.9, 129.6, 129.3, 128.7, 128.6, 127.6, 127.3, 126.0, 124.7, 114.6, 100.3, 55.3 ppm; Anal. calcd. for $\text{C}_{22}\text{H}_{18}\text{N}_4\text{OS}$: C, 68.41; H, 4.72; N, 14.53. Found: C, 68.36; H, 4.67; N, 14.48 %.

2-Methoxy-4-((2-(4-(4-methoxyphenyl)thiazol-2-yl)hydrazono)methyl)-6-nitrophenol (14f)

Red solid (0.3 g, 75 %); mp 207–209 °C (from EtOH); IR (KBr) ν_{max} 3250 (stretch N–H), 3100 (stretch C–H aromatic), 2980, 2830 (stretch C–H aliphatic), 1610 (stretch C=N), 1570, 1480 (stretch C=C), 1540, 1340 (stretch NO_2), 1250, 1050 (stretch C–O), 840, 760, 740, 690 (OOP. C–H) cm^{-1} ; R_f (60 %, hexane: EtOAc 6:3); ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ~ 11.45 (*s*, 1H, NH), 7.95 (*s*, 1H), 7.74 (*d*, $J = 8.76$ Hz, 2H), 7.67 (*dd*, $J = 1.52$ Hz, 1H), 7.49 (*dd*, $J = 1.48$ Hz, 1H), 7.09 (*s*, 1H), 6.92 (*d*, $J = 8.7$ Hz, 2H), 3.90 (*s*, 3H), 3.74 (*s*, 3H) ppm; ^{13}C NMR (125 MHz, $\text{DMSO}-d_6$) δ 168.8, 159.6, 151.1, 150.6, 144.2, 140.2, 138.1, 128.3, 127.6, 126.2, 115.1, 114.8, 113.1, 102.3, 57.4, 55.9 ppm; Anal. calcd. for $\text{C}_{18}\text{H}_{16}\text{N}_4\text{O}_5\text{S}$: C, 54.03; H, 4.06; N, 13.97. Found: C, 53.97; H, 4.02; N, 14.04 %.

4-(4-Methoxyphenyl)-2-(2-(2,4,6-trimethylbenzylidene)hydrazinyl)thiazole (14g)

Cream solid (0.26 g, 75 %); mp 229–233 °C (from EtOH); IR (KBr) ν_{max} 3400 (stretch N–H), 3100 (stretch C–H aromatic), 2950, 2820 (stretch C–H aliphatic), 1610 (stretch C=N), 1510 (stretch C=C), 1250, 1020 (stretch C–O), 850, 750, 740, (OOP. C–H) cm^{-1} ; R_f (75 %, hexane: EtOAc 6:3); ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ ~ 13.00 (*s*, 1H, NH), 8.65 (*s*, 1H), 7.52 (*d*, $J = 8.6$ Hz, 2H), 6.81 (*d*, $J = 6.9$ Hz, 2H), 6.75 (*s*, 2H), 6.56 (*s*, 1H), 3.67 (*s*, 3H), 2.31 (*s*, 6H), 2.13 (*s*, 3H) ppm; ^{13}C NMR (125 MHz, $\text{DMSO}-d_6$) δ ~ 170, 161.3, 151.2, 141.1, 140.9, 139.1, 130.4, 127.9,

126.8, 120.6, 115.0, 100.2, 55.7, 22.2, 21.5 ppm. Anal. calcd. for $\text{C}_{20}\text{H}_{21}\text{N}_3\text{OS}$: C, 68.37; H, 6.05; N, 11.94. Found: C, 68.32; H, 5.98; N, 11.98 %.

2-(Adamantan-2-ylidenehydrazono)-4-(4-methoxyphenyl)-2,3-dihydrothiazole (16)

Orange solid (0.17g, 50 %); mp 140–148 °C (from EtOH); IR (KBr) ν_{max} 2917, 2849 (stretch C–H aliphatic), 1605 (stretch C=N), 1549, 1503 (stretch C=C) cm^{-1} ; R_f (35 %, hexane: EtOAc 6:3); ^1H NMR (500 MHz, CDCl_3) δ 7.80–6.71 (*m*, Ar, NH, H-thiazole), 3.96–3.72 (OCH_3), 3.01–2.55 (CH–C=N of C and D), 2.20–1.25 (CH and CH_2 adamantan) ppm.

Biological methods

DPPH radical scavenging assay

The DPPH radical scavenging activity of **14a–d** was evaluated according to the literature [25]. A 2,2-diphenyl-2-picrylhydrazyl (DPPH) solution was prepared by dissolving an appropriate amount of DPPH in MeOH to give a concentration of 6.25×10^{-5} M. Compounds **14a–g** and DPPH with different concentrations (4000, 2000, 1000, 500, 250, 125, 62.5 $\mu\text{g/mL}$) in MeOH were prepared. Then, 0.1 mL of each hydrazinyl-thiazole solution was added to 3.9 mL of DPPH solution and was shaken vigorously. Samples were kept in darkness for 30 min and then their absorbance was measured at 517 nm. MeOH was used as blank. Radical scavenging activity was calculated as follows:

$$\text{Radical scavenging activity \%} = \left(\frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \right) \times 100$$

where A_{control} is the absorbance of the negative control (containing all reagents except test compounds), A_{sample} the absorbance of the test compounds. IC_{50} values of the test compounds were determined by plotting the radical scavenging activity percentage against concentration of the test compound.

Antibacterial assay

The antibacterial activity of hydrazinyl-thiazoles was evaluated biologically using a well-diffusion method. First, nutrient agar and nutrient broth cultures were prepared according to manufacturer's instructions and were incubated at 37 °C. After incubation for the appropriate time, a suspension of 30 μL of each bacterium was added to the nutrient agar plates. Cups (5 mm in diameter) were cut in the agar using sterilized glass tube. Each well received 30 μL of the test compounds at concentration of 4000 $\mu\text{g/mL}$ in DMSO. Then, plates were incubated at 37 °C for 24 h, after this

time the zone of inhibition was measured. The values are expressed in millimeters (mm). The experiments were performed in triplicate. The results are reported as mean \pm SD of zone of inhibition in millimeter. Antibacterial activity of each hydrazinyl-thiazole was compared with Penicillin G and Cefixime as standards. DMSO was used as negative control.

In vitro anticancer assay

The anticancer activity of the hydrazinyl-thiazole **14a** was determined against melanoma cancerous cell lines Hu-02 and A375 using the MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) method. The melanoma cancerous cell lines were provided by the Iranian Biological Resource Center, Tehran. The MTT method is based on the reductive cleavage of the tetrazolium ring of MTT into insoluble purple formazan. 100 μ L of the cell suspension were added to each well of 96-well microplate and then incubated for 12–24 h at 37 °C in a CO₂ incubator. Compound **14a** was diluted in concentrations of 0.037–0.5 mg/mL in DMSO by two-fold serial dilution. DMSO was used as negative control. Different concentrations of a test compound were added to each well and then incubated for 24 and 48 h at 37 °C in a CO₂ incubator. Then, 10 μ L of MTT (5 mg/mL) stock solution in PBS were added to each well and incubated for 4 h, the upper solution was removed, and 100 μ L of DMSO were added to the media. The plates were shaken slowly until the insoluble crystals of the produced formazan dissolved. The plates were kept for 2–4 h in the dark. After that, the plates were read at 360 nm on an ELISA reader. The experiments were performed in triplicate.

The percentage of cell viability was calculated using the following formula:

$$\% \text{ cell viability} = \frac{\text{Mean Absorbance of sample}}{\text{Mean Absorbance of control}} \times 100$$

The LC₅₀ values (lethal concentration of the test compound required to kill 50 % of sample population) were determined by plotting cell viability percentage against concentration of the sample.

Acknowledgments The authors are greatly thankful to Dr. Zohreh Ramezanpour and Miss Somaie Rasouli Dogaheh for their helpful assistance in running biological assays. The authors appreciate the valuable and helpful comments of Professor Dr. Guillermo A. Morales, Editor-In-Chief of Molecular Diversity.

References

- Hargrave KD, Hess FK, Oliver JT (1983) *N*-(4-substituted-thiazolyl)oxamic acid derivatives, a new series of potent, orally active antiallergy agents. *J Med Chem* 26:1158–1163. doi:10.1021/jm00362a014
- Patt WC, Hamilton HW, Taylor MD, Ryan MJ, Taylor DG Jr, Connolly CJC, Doherty AM, Klutchko SR, Sircar I, Steinbaugh BA, Batley BL, Painchaud CA, Rapundalo ST, Michniewicz BM, Olson SCJ (1992) Structure-activity relationships of a series of 2-amino-4-thiazole-containing renin inhibitors. *J Med Chem* 35:2562–2572. doi:10.1021/jm00092a006
- Aggarwal R, Kumar S, Kaushik P, Dhirender K, Girish Kumar G (2013) Synthesis and pharmacological evaluation of some novel 2-(5-hydroxy-5-trifluoromethyl-4,5-dihydropyrazol-1-yl)-4-(coumarin-3-yl)thiazoles. *Eur J Med Chem* 62:508–514. doi:10.1016/j.ejmech.2012.11.046
- Cantrell AS, Engelhardt P, Högberg M, Jaskunas SR, Gunnar Johansson N, Jordan CL, Kangasmetsä J, Kinnick MD, Lind P, Morin JM, Muesing MA Jr, Noreén R, Öberg B, Pranc P, Sahlberg C, Ternansky RJ, Vasileff RT, Vrang L, West SJ, Zhang H (1996) Phenethylthiazolethiourea (PETT) compounds; a new class of HIV-1 reverse transcriptase inhibitors. Synthesis and basic structure activity relationship studies of PETT analogs. *J Med Chem* 39:4261–4274. doi:10.1021/jm950639r
- Jaen JC, Wise LD, Caprathe BW, Tecle H, Bergmeier S, Humblet CC, Heffner TG, Meltzner LT, Pugsley TA (1990) 4-1,2,5,6-Tetrahydro-1-alkyl-3-pyridinyl)-2-thiazolamines: A novel class of compounds with central dopamine agonist properties. *J Med Chem* 33:311–317. doi:10.1021/jm00163a051
- Tsuji K, Ishikawa H (1994) Synthesis and anti-pseudomonal activity of new 2-isocephems with a dihydroxypyridone moiety at C-7. *Bioorg Med Chem Lett* 4:1601–1606. doi:10.1016/S0960-894X(01)80574-6
- Jukić M, Dorđević A, Lazarević J, Gobec M, Šmelcerović A, Anderluh A (2013) Antimicrobial activity and cytotoxicity of some 2-amino-5-alkylidene-thiazol-4-ones. *Mol Divers* 17:773–780. doi:10.1007/s11030-013-9474-6
- Li M, Sim Y, Wook Ham S (2010) Discovery of 2-aminothiazole derivatives as antitumor agents. *Bull Korean Chem Soc* 31:1463–1464. doi:10.5012/bkcs.2010.31.6.1463
- Regnier G, Canevar L, Le RJ, Douarec JC, Halstop S, Daussy J (1972) Triphenylpropylpiperazine derivatives as new potent analgetic substances. *J Med Chem* 15:295–301. doi:10.1021/jm00273a600
- Siddiqui N, Waqar A (2010) Triazole incorporated thiazoles as a new class of anticonvulsants: Design, synthesis and in vivo screening. *Eur J Med Chem* 45:1536–1543. doi:10.1016/j.ejmech.2009.12.062
- Ha YM, Uehara Y, Park D, Jeong HO, Park JY, Park YJ, Lee JY, Lee HJ, Song YM, Moon HR, Chung HY (2012) Synthesis and preliminary in vitro biological evaluation of 5-chloro-2-(substituted Phenyl)benzo[d]thiazole derivatives designed as novel antimelanogenesis agents. *Appl Biochem Biotechnol* 168:1416–1433. doi:10.1007/s12010-012-9867-5
- Chandra Sekhar KVG, Rao VS, Deuther-Conrad W, Sridhar D, Nagesh HN, Kumar VS, Brust P, Kumar MMK (2013) Design, synthesis, and preliminary in vitro and in vivo pharmacological evaluation of 4-4-[2-(4-(2-substitutedquinoxalin-3-yl)piperazin-1-yl)ethyl]phenylthiazoles as atypical antipsychotic agents. *Med Chem Res* 22:1660–1673. doi:10.1007/s00044-012-0164-1
- Anbazhagan R, Sankaran KR (2013) Syntheses, spectral characterization, single crystal X-ray diffraction and DFT computational studies of novel thiazole derivatives. *J Mol Struct* 1050:73–80. doi:10.1016/j.molstruc.2013.07.019
- McIntyre NA, McInnes C, Griffiths G, Barnett AL, Kontopidis G, Slawin AM, Jackson W, Thomas M, Zheleva DI, Wang S, Blake DG, Westwood NJ, Fischer PM (2010) Design, synthesis, and evaluation of 2-methyl- and 2-amino-*N*-aryl-4,5-dihydrothiazolo[4,5-*h*]quinazolin-8-amines as ring-constrained 2-anilino-4-(thiazol-5-yl)pyrimidine cyclin-dependent kinase inhibitors. *J Med Chem* 53:2136–2145. doi:10.1021/jm901660c

15. Mohammed Khan K, Karim A, Saied S, Ambreen N, Rustamova X, Naureen S, Mansoor S, Ali M, Perveen S, Choudhary MI, Antonio Morales G (2014) Evaluation of the thiazole Schiff bases as β -glucuronidase inhibitors and their in silico studies. *Mol Divers* 18:295–306. doi:[10.1007/s11030-013-9500-8](https://doi.org/10.1007/s11030-013-9500-8)
16. Imramovsky A, Pejchal V, Štěpánková Š, Vorčáková K, Jampílek J, Vancod J, Šimunek P, Královce K, Brucková L, Mandíková J, Trejtnar F (2013) Synthesis and in vitro evaluation of new derivatives of 2-substituted-6-fluorobenzo[d]thiazoles as cholinesterase inhibitors. *Bioorg Med Chem* 21:1735–1748. doi:[10.1016/j.bmc.2013.01.052](https://doi.org/10.1016/j.bmc.2013.01.052)
17. Carradori S, Secci D, Bolasco A, Rivanera D, Mari E, Zicari A, Lotti LV, Bizzarri B (2013) Synthesis and cytotoxicity of novel (thiazol-2-yl)hydrazine derivatives as promising anti-candida agents. *Eur J Med Chem* 65:102–111. doi:[10.1016/j.ejmech.2013.04.042](https://doi.org/10.1016/j.ejmech.2013.04.042)
18. Abbasi Shiran J, Yahyazadeh A, Mamaghani M, Rassa M (2013) Regioselective synthesis of novel 3-allyl-2-(substituted imino)-4-phenyl-3H-thiazole and 2, 2'-(1,3-phenylene)bis(3-substituted-2-imino-4-phenyl-3H-thiazole) derivatives as antibacterial agents. *J Mol Struct* 1039:113–118. doi:[10.1016/j.molstruc.2013.02.003](https://doi.org/10.1016/j.molstruc.2013.02.003)
19. Kudrjasova J, Herckens R, Penxten H, Adriaenssens P, Lutsen L, Vanderzandea D, Maes W (2014) Direct arylation as a versatile tool towards thiazolo[5,4-d]thiazole-based semiconducting materials. *Org Biomol Chem* 12:4663–4672. doi:[10.1039/C4OB00360H](https://doi.org/10.1039/C4OB00360H)
20. Giridhar T, Cho W, Park J, Park J-S, Gal Y-S, Kang S, Leec JY, Jin S-H (2013) Facile synthesis and characterization of iridium(III) complexes containing an N-ethylcarbazole-thiazole main ligand using a tandem reaction for solution processed phosphorescent organic light-emitting diodes. *J Mater Chem C* 1:2368–2378. doi:[10.1039/C3TC00323J](https://doi.org/10.1039/C3TC00323J)
21. Hela A, Harun M, Choi C-H, Kima H-S (2012) New regioisomeric naphthol-substituted thiazole based ratiometric fluorescence sensor for Zn^{2+} with a remarkable red shift in emission spectra. *Tetrahedron* 68:647–653. doi:[10.1016/j.tet.2011.10.106](https://doi.org/10.1016/j.tet.2011.10.106)
22. Sharifzadeh B, Mahmoodi NO, Mamaghani M, Tabatabaiean K, Salimi Chirani A, Nikokar I (2013) Facile regioselective synthesis of novel bioactive thiazolyl-pyrazoline derivatives via a three-component reaction and their antimicrobial activity. *Bioorg Med Chem Lett* 23:548–551. doi:[10.1016/j.bmcl.2012.11.024](https://doi.org/10.1016/j.bmcl.2012.11.024)
23. Mahmoodi NO, Parvizi J, Sharifzadeh B, Rassa M (2013) Facile regioselective synthesis of novel bis-thiazole derivatives and their antimicrobial activity. *Arch Pharm Chem Life Sci* 346:860–864. doi:[10.1002/ardp.201300187](https://doi.org/10.1002/ardp.201300187)
24. Mahmoodi NO, Safari N, Sharifzadeh B (2014) One-Pot synthesis of novel 2-(thiazol-2-yl)-4,5-dihydropyridazin-3(2H)-one derivatives catalyzed by activated KSF. *Synth Commun* 44:245–250. doi:[10.1080/00397911.2013.801077](https://doi.org/10.1080/00397911.2013.801077)
25. Mahmoodi NO, Ramzanpour S, Ghanbari Pirbasti F (2015) One-pot multi-component synthesis of 1,4-dihydropyridines using Zn^{2+} @KSF and evaluation their antibacterial and antioxidant activities. *Arch der Pharm* 348:275–282. doi:[10.1002/ardp.201400414](https://doi.org/10.1002/ardp.201400414)
26. Mahmoodi NO, Mamaghani M, Behzadi T (2012) Synthesis and structure-behavior relationships of tetra-substituted imidazole derivatives of 1, 3-diazabicyclo [3, 1, 0] hex-3-ene. *Mol Divers* 16:737–747. doi:[10.1007/s11030-012-9409-7](https://doi.org/10.1007/s11030-012-9409-7)
27. Caputto ME, Ciccarelli A, Frank F, Moglioni AG, Moltrasio GY, Vega D, Lombardo E, Finkelsztejn LM (2012) Synthesis and biological evaluation of some novel 1-indanone thiazolylhydrazones derivatives as anti-trypanosoma cruzi agents. *Eur J Med Chem* 55:155–163. doi:[10.1016/j.ejmech.2012.07.013](https://doi.org/10.1016/j.ejmech.2012.07.013)
28. Secci D, Bolasco A, Carradori S, D'Ascenzio M, Nescatelli R, Yáñez M (2012) Recent advances in the development of selective human MAO-B inhibitors: (Hetero)arylidene-(4-substituted-thiazol-2-yl)hydrazines. *Eur J Med Chem* 58:405–417. doi:[10.1016/j.ejmech.2012.10.032](https://doi.org/10.1016/j.ejmech.2012.10.032)
29. Gaikwad ND, Patil SV, Bobade VD (2012) Hybrids of ravuconazole: synthesis and biological evaluation. *Eur J Med Chem* 54:295–302. doi:[10.1016/j.ejmech.2012.05.010](https://doi.org/10.1016/j.ejmech.2012.05.010)
30. Arshad A, Osman H, Bagley MC, Kit Lam C, Mohamad S, Safirah A, Zahariluddin M (2011) Synthesis and antimicrobial properties of some new thiazolyl coumarin derivatives. *Eur J Med Chem* 46:3788–3794. doi:[10.1016/j.ejmech.2011.05.044](https://doi.org/10.1016/j.ejmech.2011.05.044)
31. Hassan AA, Ibrahim YR, El-Sheref EM, Abdel-Aziz M, Bräse S, Nieger M (2013) Synthesis and antibacterial activity of 4-aryl-2-(1-substituted ethylidene)thiazoles. *Arch Pharm Chem Life Sci* 346:562–570. doi:[10.1002/ardp.201300099](https://doi.org/10.1002/ardp.201300099)
32. Osman H, Arshad A, Kit Lam C, Bagley MC (2012) Microwave-assisted synthesis and antioxidant properties of hydrazinyl thiazolyl coumarin derivatives. *Chem Cent J* 6:32. doi:[10.1186/1752-153X-6-32](https://doi.org/10.1186/1752-153X-6-32)
33. Bharti SK, Nath G, Tilak R, Singh SK (2010) Synthesis, antibacterial and anti-fungal activities of some novel Schiff bases containing 2,4-disubstituted thiazole ring. *J Med Chem* 45:651–660. doi:[10.1016/j.ejmech.2009.11.008](https://doi.org/10.1016/j.ejmech.2009.11.008)
34. Alam MS, Liu L, Lee YE, Lee DU (2011) Cytotoxicity of new 5-phenyl-4,5-dihydro-1,3,4-thiadiazole analogues. *Chem Pharm Bull* 59:568–573. doi:[10.1002/chin.201215131](https://doi.org/10.1002/chin.201215131)