This article was downloaded by: [Mount Allison University OLibraries] On: 05 June 2013, At: 11:27 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Coordination Chemistry

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gcoo20

Synthesis and characterization of two acetato-bridged dinuclear copper(II) complexes with 4-bromo-2-((4 or 6methylpyridin-2-ylimino)methyl)phenol as ligand

Farshideh Hasanvand $^{\rm a}$, Ali Hoseinzadeh $^{\rm a}$, Javad Zolgharnein $^{\rm a}$ & Saeid Amani $^{\rm a}$

^a Department of Chemistry, Arak University, Dr Beheshti Ave., Arak 38156, Iran Published online: 12 Oct 2009.

To cite this article: Farshideh Hasanvand , Ali Hoseinzadeh , Javad Zolgharnein & Saeid Amani (2010): Synthesis and characterization of two acetato-bridged dinuclear copper(II) complexes with 4-bromo-2-((4 or 6-methylpyridin-2-ylimino)methyl)phenol as ligand, Journal of Coordination Chemistry, 63:2, 346-352

To link to this article: http://dx.doi.org/10.1080/00958970903330294

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <u>http://www.tandfonline.com/page/terms-and-</u> conditions

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



Synthesis and characterization of two acetato-bridged dinuclear copper(II) complexes with 4-bromo-2-((4 or 6-methylpyridin-2-ylimino)methyl)phenol as ligand

FARSHIDEH HASANVAND, ALI HOSEINZADEH, JAVAD ZOLGHARNEIN and SAEID AMANI*

Department of Chemistry, Arak University, Dr Beheshti Ave., Arak 38156, Iran

(Received 26 July 2009; in final form 27 July 2009)

The synthesis and characterization of two new acetato-bridged dinuclear copper(II) complexes are described. Both compounds have the general formula $[Cu(L)(\mu-O_2C-CH_3)]_2$, in which L = 4-bromo-2-((4-methylpyridin-2-ylimino)methyl)phenol or 4-bromo-2-((6-methylpyridin-2-ylimino)methyl)phenol. The title compounds consist of dinuclear units with bridging acetato groups and a ligand linked to each copper via the phenol oxygen and nitrogen. Both compounds were synthesized in a one-step reaction and characterized by elemental analysis, Fourier transform infrared (FTIR), electron spin resonance (ESR), and electronic spectra and by room temperature magnetic moments. The compounds exhibit antiferromagnetic interactions at room temperature. UV-Vis spectra show four absorptions attributed to d-d transitions of copper, ligand \rightarrow metal charge transfer and $\pi \rightarrow \pi^*$ or $n \rightarrow \pi^*$ transitions of ligand. The FTIR spectra indicate a $Cu_2O_4C_2$ ring vibration. Both complexes show room temperature magnetic moments of about 1.6 B.M. per copper. The X-band ESR studies indicate a weak half-field band, characteristic of the Cu(II)-Cu(II) dimer, observed at 1552 and 1558 G for the complexes, strongly suggesting that the hyperfine structure arises from a spin triplet species. The spectra of frozen samples in DMSO or DMF at liquid nitrogen temperature show a typical $\Delta m = 1$ transition.

Keywords: Dinuclear copper(II) complexes; Spin-spin interaction; Diacetato-bridged

1. Introduction

A fascinating field of research is the study of polynuclear complexes. Magnetic properties of a polynuclear complex of paramagnetic metal ions are, in general, different from the sum of the magnetic properties of each metal ion fragment separately [1]. Bleaney and Bowers [2] demonstrated that the magnetic behavior of copper(II) acetate monohydrate was due to the dinuclear nature and magnetic exchange interaction. Since then, exchange coupled polynuclear systems have seen exponential increase in publications [3–8]. Such interaction between metal centers lies at the center of widely separated areas, such as physics of magnetic materials and the role of polynuclear reaction sites in biological [9–12] and catalytic processes [13–16]. Dinuclear copper(II)

^{*}Corresponding author. Emails: s-amani@araku.ac.ir; amani1952@yahoo.com

complexes have relatively simple single-ion properties facilitating interpretation of magnetic data. The simplest bridging systems use monoatomic bridges, such as hydroxo [17], alkoxo [18], chloro [19], bromo [20], fluoro [21], and azido [22]. Despite a vast number of experimental and theoretical studies, our understanding of the magnitude of the copper(II)–copper(II) coupling is still imperfect. In the present study, two dinuclear copper(II) complexes are reported with 4-bromo-2-((4-methylpyridin-2-ylimino)–methyl)phenol or 4-bromo-2-(6-methylpyridin-2-ylimino)methyl)phenol (hereafter abbreviated L_1 and L_2 , respectively), $[Cu_2(L_1)_2(O_2C-CH_3)_2]$ or $[Cu_2(L_2)_2(O_2C-CH_3)_2]$.

2. Experimental

2.1. Chemicals

All chemicals were of reagent grade quality purchased from Merck Chemical Company and used as received.

2.2. Physical measurements

The C, H, and N determinations were made in the Research Institute of Petroleum Industry of Iran. Cu determination was carried out on a Perkin-Elmer 2380 Atomic Absorption spectrophotometer at 324.7 nm. Electronic spectra were recorded on a Perkin-Elmer Lambda 900 spectrophotometer. Fourier transform infrared (FTIR) spectra were made from 4000–250 cm⁻¹ using a Galaxy 5000 spectrophotometer. Spectra were calibrated using polystyrene bands at 3028, 1601, and 1208 cm⁻¹. X-band electron paramagnetic resonance spectra were recorded on powder and frozen solutions at room and at liquid nitrogen temperatures in DMF or DMSO on an IBM electron spin resonance (ESR) spectrometer using DPPH (g = 2.0036) as a standard. A Johnson Matthey Alfa products magnetic susceptibility balance was used to measure the room temperature magnetic moments.

2.3. Preparation of the ligands

The ligands were prepared according to the following procedures.

2.3.1. 4-Bromo-2-((4-methypyridin-2-ylimino)methyl)phenol (L₁). 10 mmol 2-amino-4-methylpyridine in 15 mL methanol was added to 10 mmol 4-bromo-2-hydroxybenzaldeyhyde in 25 mL CH₃OH. By the addition of five drops concentrated H₂SO₄ and 2 h reflux, yellow precipitate separated was washed with hot methanol and air dried. Yield $ca \approx 96\%$. Elemental analysis for C₁₃H₁₁N₂BrO: Found (%): C, 52.65; H, 3.74; N, 9.87. Calcd (%): C, 53.36; H, 3.81; N, 9.62. ¹H NMR (CDCl₃, ppm): δ 2.28(s, 3H_{met.}), 6.95– 8.40(m, 6H_{aro.}), 9.47(s, 1H_{imin.}), 13.05(s, 1H_{phen.}) (scheme 1).

2.3.2. 4-Bromo-2-((6-methypyridin-2-ylimino)methyl)phenol (L₂). This compound was obtained by similar method as described for L₁. Yield $ca \approx 95\%$. Elemental analysis for



Scheme 1. Structure of ligand L₁.



Scheme 2. Structure of ligand L₂.



Scheme 3. Proposed structure for 1.

 $C_{13}H_{11}N_2BrO$: Found (%): C, 54.58; H, 3.91; N, 9.50. Calcd (%): C, 53.36; H, 3.81; N, 9.62. ¹H NMR (CDCl₃, ppm): $\delta 2.66(s, 3H_{met.})$, 6.92–7.78(m, $6H_{aro.})$, 9.40(s, $1H_{imin.}$), 13.10(s, $1H_{phen.}$) (scheme 2).

2.4. Preparation of the complexes

Coordination compounds were prepared according to the following general procedure.

2.4.1. $[Cu_2(L_1)_2(\mu-O_2C-CH_3)_2]$ (1). 2 mmol ligand L₁ and 2 mmol triethylamine were dissolved in 25 mL methanol; 2 mmol Cu(O₂C-CH₃)₂ · H₂O was also dissolved in 50 mL CH₃OH. The Cu(II) salt solution was then added slowly to the ligand solution. After 4 h reflux, the brown precipitate separated and was washed with hot methanol and THF. Yield $ca \approx 55\%$. Elemental analysis for C₃₀H₂₆Br₂O₆N₄Cu₂: Found (%): C, 44.13; H, 3.44; N, 6.37; Cu, 15.73. Calcd (%): C, 43.65; H, 3.18; N, 6.79; Cu, 15.40 (scheme 3).



Scheme 4. Proposed structure for 2.

Table 1. Spectroscopic data for 1 and 2.

Complex	Absorption UV-Vis (nm)	IR (acetate) (cm ⁻¹)	$\begin{array}{c} IR(Cu–O) \\ (Cu–N) \\ (cm^{-1}) \end{array}$	ESR half-field	ESR powder	ESR solution in DMF	μ _{Cu} (R.T.) B.M.
1	667, 418, 295, 270	1556, 1415	549, 464, 349, 279	$g_{\rm h} = 4.36$	g=2.06	$g_{\parallel} = 2.06$ $A_{\parallel} \approx 165$ $A_{n\perp} \approx 15$	1.58
2	671, 413, 292, 277	1527, 1381	534, 439, 354, 283	$g_{\rm h} = 4.04$	g=2.06	$g_{\parallel} = 2.07$ $A_{\parallel} \approx 160$ $A_{N\perp} \approx 15$	1.63

2.4.2. $[Cu_2(L_2)_2(\mu-O_2C-CH_3)_2]$ (2). This complex was obtained by similar method as reported for 2. Yield $ca \approx 45\%$. Elemental analysis for $C_{30}H_{26}Br_2O_6N_4Cu_2$: Found (%): C, 44.28; H, 2.97; N, 6.61; Cu, 16.22. Calcd (%): C, 43.65; H, 3.18; N, 6.79; Cu, 15.40 (scheme 4).

3. Results and discussion

3.1. Electronic spectra

Spectroscopic data for all compounds are presented in table 1. Electronic spectra of both complexes were obtained from a solid sample using diffuse reflectance and are very similar, illustrating similar geometry. The compounds show a broad band at 667 nm for 1 and 671 for 2 due to ligand field transition for a CuN₂O₃ chromophore [23, 24]. An absorption band at 418 nm for 1 and 413 nm for 2 are assigned to charge transfer from oxygen to $d_{x^2-y^2}$ orbital of copper(II) [23, 25]. The last two absorptions at 295 and 275 nm for both 1 and 2 are associated with $\pi \to \pi^*$ or $n \to \pi^*$ transitions of the ligand [26].

3.2. Infrared spectra

The infrared spectra of acetate and $Cu_2N_2O_2$ are of particular interest [27–34]. All bands present in the free ligand are also observed in the spectra of the complexes.



Figure 1. Frozen solution (77 K) ESR X-band for complex 1 in DMF. *Denotes (diphenylpicrylhydrazine) DPPH marker (g = 2.0036).

Based on data from earlier reports [35, 36], we assign bands at 549 and 464 cm⁻¹ and at 534 and 439 cm⁻¹ (for 1 and 2, respectively) to Cu–O vibrations. The bands at 349 and 279 cm⁻¹ and at 354 and 283 cm⁻¹ (for 1 and 2, respectively) are assigned to Cu–N vibrations: however, these vibrations are likely coupled Cu–N and Cu–O vibrations. These data indicate L_1 is closer to copper(II) than L_2 due to the steric effect of methyl ortho to nitrogen in L_2 . The bridging acetate in 1 has higher Cu–O vibration energy in 2 consistent with shorter Cu–Cu distance in 1 than in 2.

3.3. Magnetic moment and ESR

Both complexes are dimeric with room temperature magnetic moments of 1.58 and 1.63 B.M. per copper for 1 and 2, respectively, low for a d⁹ configuration from spin–spin interaction through the bridging ligands [37]. The spin–spin interaction in 1 is higher than in 2, in agreement with data from IR spectra [8]. To provide further evidence for antiferromagnetic interaction, X-band powder and frozen solution ESR spectra were obtained at liquid nitrogen and at room temperatures. In the solid state, the spectra were similar and ESR silent with only a weak signal due to monomeric impurity (g = 2.06). Frozen solution ESR spectra of both compounds is typical for a dinuclear copper(II) complex. A weak $\Delta m = 2$ transition for both compounds confirms Cu ··· Cu magnetic exchange interaction at half-field (1552 G for 1 and 1558 G for 2), which is characteristic of a dimer [2, 38, 39].

The g_{\parallel} values of 2.06 and 2.07 (for **1** and **2**, respectively) with $A_{\parallel} \approx 162 \text{ G}$ is in agreement with a $d_{x^2-y^2}$ ground state and a square-pyramidal geometry [40–43]. The A_N value of $\approx 15 \text{ G}$ and the presence of seven peaks for nitrogen superhyperfine structure for the complexes agrees with two N donors per copper(II) (see figure 1 and Supplementary material).

4. Conclusion

Two new bis-(µ-alkoxo)-bridged dinuclear copper(II) complexes with 4-bromo-2-((4-methylpyridine-2-ylimino)methyl)phenol or 4-bromo-2-((6-methylpyridine-2-ylimino)-methyl)phenol as the ligands have been made in a one-step synthesis and

spectroscopically characterized. Both complexes show subnormal magnetic moments between 1.58 and 1.63 B.M. per copper at room temperature, suggesting an antiferromagnetic spin-exchange interaction within each molecule. Data from IR spectroscopy show that L_1 is closer to copper(II) than L_2 due to the steric effect of methyl; the IR for bridging acetate is in agreement with the magnetic moments. The X-band ESR spectra of 1 and 2 with $g_{\parallel} = 2.06$ and $A_{\parallel} \approx 163$ G are typical of dimeric square-pyramidal copper(II) complexes.

Acknowledgements

The authors would like to thank the Research Council of Arak University for financial support of this research. The authors also appreciate the Chemistry Institute and DAAD Organization in Chemnitz University in Germany for the services, facilities, and physical measurements.

References

- [1] O. Kahn. Angew. Chem. Int. Ed. Engl., 24, 834 (1985).
- [2] B. Bleaney, K.D. Bowers. Proc. R. Soc. London Ser. A, 214, 451 (1952).
- [3] C.J. Cains, D.H. Busch. Coord. Chem. Rev., 69, 1 (1986).
- [4] O. Kahn. Inorg. Chim. Acta, 62, 3 (1982).
- [5] O. Kahn. Struct. Bond., 68, 89 (1987).
- [6] E. Sinn. Coord. Chem. Rev., 3, 313 (1970).
- [7] P.W. Ball. Coord. Chem. Rev., 4, 361 (1969).
- [8] R.L. Martin. In New Pathway in Inorganic Chemistry, E.A.V. Absworth, A.G. Maddock, A.G. Sharp (Eds), p. 175, Cambridge University Press, London (1968).
- [9] P.A. Vigato, S. Tamburini. Coord. Chem. Rev., 106, 25 (1990).
- [10] J. Peisach, P. Aisen, W.E. Blumberg. The Biochemistry of Copper, Academic Press, New York (1966).
- [11] E.I. Solomon. In Copper Proteins, T.G. Spiro (Ed.), p. 2, Wiley-Interscience, New York (1981).
- [12] E.I. Solomon, D.E. Wilcox. In Magneto-Structural Correlations in Exchange Coupled Systems, NATO Advanced Study Institute Series, D. Gatteschi, O. Kahn, R.D. Willett (Eds), Vol. C140, p. 463, D. Reidel Dordrecht, Holland (1984).
- [13] G. Challa, W. Chen, J. Reedijk. Makromol. Chem. Macromol., Symp., 59 (1992).
- [14] M.M. Rogic, T.R. Demmin. J. Am. Chem. Soc., 100, 5472 (1978).
- [15] H.C. Meiders, F. Van Bolhuis, G. Challa. J. Mol. Catal., 5, 225 (1979).
- [16] J. Reedijk. Bioinorganic Catalysis, Marcel Dekker, New York (1993).
- [17] D.J. Hodgson. J. Mol. Catal., 23, 219 (1984).
- [18] R. MerGehenn, L. Merz, W. Hasse. J. Chem. Soc., Dalton Trans., 1703 (1980).
- [19] G.O. Bannon, R.D. Willett. Inorg. Chim. Acta, 52, L131 (1981).
- [20] C.P. Landee, R.E. Greeney. Inorg. Chem., 25, 3771 (1986).
- [21] F.J. Rietmeijer, R.A.G. De Graaff, J. Reedijk. Inorg. Chem., 23, 151 (1984).
- [22] S. Sikorav, I. Bkouche-Waksman, O. Kahn. Inorg. Chem., 23, 490 (1984).
- [23] A. Bencini, D. Gatesschi, C. Zanchini, W. Haase. Inorg. Chem., 24, 3485 (1985).
- [24] A.B.P. Lever. Inorganic Electronic Spectroscopy, 2nd Edn, Elsevier, Amsterdam (1984).
- [25] B.J. Hathaway, H. Elliott. Inorg. Chem., 5, 885 (1966).
- [26] Y. Nishida, S. Kida. Coord. Chem. Rev., 27, 275 (1979)
- [27] E. Gomashchi. Spectrosc. Lett., 27, 829 (1994).
- [28] S. Amani, N. Foroughifar, H. Hamidi, L.J. Theriot. J. Sci. I.R. Iran, 2, 143 (1998).
- [29] S. Amani, G.A. Van Albada, I. Mutikainen, U. Trupinen, J. Reedijk. Polyhedron, 14, 1991 (1999).
- [30] S. Amani, G.A. Van Albada, H. Kooijman, A.L. Spek, J. Reedijk. Inorg. Chim. Acta, 286, 24 (1999).
- [31] S. Amani, G.A. Van Albada, H. Kooijman, A.L. Spek, J. Reedijk. Inorg. Chim. Acta, 287, 226 (1999).
- [32] S. Amani, N. Sadeghy, Z. Talebi. J. Chem. Soc. Pak., 4, 271 (2000).
- [33] M. Dadpu, A.R. Mahjoob, S. Amani. Iran. J. Chem. Chem. Eng., 1, 37 (2003).

- [34] S. Amani, N. Sadeghi, J. Zolgharnein. J. Sci. Teach. Train. Univ., 3, 32 (2003).
- [35] N. Sadeghy, S. Amani. Polish J. Chem., 79, 1749 (2005).
- [36] G.A. Van Albada, M.T. Lakin, N. Veldman, A.L. Spek, J. Reedijk. Inorg. Chem., 34, 4910 (1995).
- [37] G.A. Van Albada, W.J.J. Smeets, A.L. Spek, J. Reedijk. Inorg. Chim. Acta, 260, 151 (1997).
- [38] P.W. Sekwood. Magnetochemistry, 2nd Edn, Interscience, New York (1956).
- [39] A.P. Ginsberg. Inorg. Chim. Acta Rev., 5, 45 (1971).
- [40] S.M. Nelson, C.V. Knox. J. Chem. Soc., Dalton Trans., 2525 (1981).
- [41] B.J. Hathaway, G. Wilkinson, R.D. Gill, J.A. McClevery (Eds). *Comprehensive Coordination Chemistry*, Vol. 5, Pergamon, Oxford (1987).
- [42] A.L. Abuhijleh, C. Woods. Inorg. Chim. Acta, 209, 187 (1993).
- [43] H. Zheng, L. Que Jr. Inorg. Chim. Acta, 263, 301 (1997).