

Table 2. Selected geometric parameters (Å, °)

Zr—O2	2.1126 (11)	C1—C7	1.485 (2)
Zr—O1	2.1436 (11)	C7—C8	1.386 (2)
Zr—Cl	2.5484 (9)	C8—C9	1.386 (3)
O1—C7	1.277 (2)	C9—C10	1.484 (2)
O2—C9	1.287 (2)		
O2—Zr—O2 ⁱ	115.81 (2)	O2—Zr—Cl	78.02 (3)
O2—Zr—O1 ⁱ	149.67 (5)	O1—Zr—Cl	132.30 (3)
O2—Zr—O1	76.55 (4)	C7—O1—Zr	134.57 (10)
O2 ⁱ —Zr—O1	77.80 (4)	C9—O2—Zr	136.28 (10)
O1 ⁱ —Zr—O1	79.66 (5)		

Symmetry code: (i) 1 - y, x - y, z.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989). Cell refinement: *CAD-4 Software*. Data reduction: *SDP-Plus* (Frenz, 1988). Program(s) used to solve structure: *SHELXS86* (Sheldrick, 1985). Program(s) used to refine structure: *SHELXL93* (Sheldrick, 1993). Molecular graphics: *ORTEP3* for Windows (Farrugia, 1997).

The authors thank the DFG, the Fonds der Chemischen Industrie, and the graduate college 'Synthetic, mechanistic and reaction engineering aspects of metal catalysts' at the TU, Berlin, for financial support.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: CF1194). Services for accessing these data are described at the back of the journal.

References

- Chun, H. K., Steffen, W. L. & Fay, R. C. (1979). *Inorg. Chem.* **18**, 2458–2465.
- Cox, E. G., Cruickshank, D. W. J. & Smith, J. A. S. (1958). *Proc. R. Soc. London Ser. A*, **247**, 14–21.
- Cox, M., Lewis, L. & Nyholm, R. S. (1964). *J. Chem. Soc.* pp. 6113–6120.
- Enraf–Nonius (1989). *CAD-4 Software*. Version 5.0. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1997). *ORTEP3* for Windows. Version 1.0.1β. University of Glasgow, Scotland.
- Frenz, B. A. (1988). *Enraf–Nonius SDP-Plus Structure Determination Package*. Version 5.0. Enraf–Nonius, Delft, The Netherlands.
- Goddard, R., Haenel, M. W., Herndon, W. C., Krüger, C. & Zander, M. (1995). *J. Am. Chem. Soc.* **117**, 30–41.
- Hunter, C. A. (1993). *Angew. Chem. Int. Ed. Engl.* **32**, 1584–1586.
- Janiak, C., Scharmann, T. G. & Lange, K. C. H. (1994). *Macromol. Rapid Commun.* **15**, 655–658.
- Longo, P., Proto, A. & Oliva, L. (1994). *Macromol. Rapid Commun.* **15**, 151–155.
- Muller, E. G., Day, V. W. & Fay, R. C. (1975). *J. Am. Chem. Soc.* **98**, 2165–2172.
- Orpen, A. G., Brammer, L., Allen, F. H., Kennard, O., Watson, D. G. & Taylor, R. (1989). *J. Chem. Soc. Dalton Trans.* pp. S1–83.
- Piermarini, G. J., Mighell, A. D., Weir, C. E. & Block, S. (1969). *Science*, **165**, 1250–1255.
- Sheldrick, G. M. (1985). *SHELXS86. Program for the Solution of Crystal Structures*. University of Göttingen, Germany.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Silverton, J. V. & Hoard, J. L. (1963). *Inorg. Chem.* **2**, 243–249.
- Stezowski, J. J. & Eick, H. A. (1965). *J. Am. Chem. Soc.* **91**, 2890–2894.
- Ueki, S., Furuhashi, H., Murata, M. & Toki, S. (1992). *Chem. Abstr.* **116**, 7090b; (1993). US Patent 5 223 465, Tonen Corporation, Japan.
- VonDreele, R. B., Stezowski, J. J. & Fay, R. C. (1971). *J. Am. Chem. Soc.* **93**, 2887–2893.
- Wolf, L. & Tröltzsch, C. (1962). *J. Prakt. Chem.* **17**, 78–91

Acta Cryst. (1998). **C54**, 212–214

(9,10-Dioxo-9,10-dihydro-1-anthrylaminato-N)(triphenylphosphine-P)gold

ANDREI V. CHURAKOV,^{a†} LYUDMILA G. KUZ'MINA^a AND JUDITH A. K. HOWARD^b

^aN. S. Kurnakov Institute of General and Inorganic Chemistry of Russian Academy of Science, 31 Leninskii prospect, Moscow 117907, Russia, and ^bDepartment of Chemistry, University of Durham, Science Laboratories, South Road, Durham DH1 3LE, England. E-mail: andrei.churakov@durham.ac.uk



(Received 29 April 1997; accepted 2 October 1997)

Abstract

In the title compound, [Au(C₁₄H₈NO₂)(C₁₈H₁₅P)], the AuPPh₃ group is located at the amino group. The Au—N and Au—P distances are 2.019 (3) and 2.2420 (8) Å, respectively. The amino group H atom is involved in an intramolecular hydrogen bond with a carbonyl O atom. The bond-length distribution in the aminoanthraquinone ligand is slightly different from that in the free ligand, manifesting itself as a more pronounced contribution from the quinoid structure.

Comment

The title compound, (1), is a (triphenylphosphine)gold derivative of 1-aminoanthracene-9,10-dione, (2). The initial photochromic organic compound (2) can exist in two tautomeric forms, amino and imino (see scheme below).



In the imino form, two types of imino group configuration, resulting in different kinds of hydrogen bonds, may be proposed. Based on X-ray structural investiga-

[†] Current address: Department of Chemistry, University of Durham, Science Laboratories, South Road, Durham DH1 3LE, England.

tions, only the amino form has been found to occur in the crystalline state of (2) (Almond *et al.*, 1983; Foitzik & Paulus, 1986), although examples of organic molecules existing in both tautomeric forms in the crystal are known, for example, 2-pyridone-6-chloro-2-hydroxypyridine (Almlöf *et al.*, 1971).

The title compound is the first representative of a series of mono- and poly-aurated amino derivatives of anthraquinone, which we regard as models for structures with chains of intermolecular $\text{Au} \cdots \text{Au}$ interactions.

The structure determination of the title compound shows that the amino tautomeric form exists in the crystal, revealing a change in the geometrical parameters of the aminoanthracene chromophore due to the introduction of the AuPPH_3 substituent at the amino group. In the amino form, either a hydrogen bond or an $\text{Au} \cdots \text{O}$ secondary bond stabilizes the structure of (1). The latter has been observed in three crystalline modifications of (8-hydroxyquinolino)(triphenylphosphine)gold (Kuz'mina, Dvortsova *et al.*, 1989) and in (8-quinolinato-*O,N*)bis(triphenylphosphine)digold(I) tetrafluoroborate (Kolb *et al.*, 1993).

The amino tautomeric form of a molecule of (1) is shown in Fig. 1. Competition between an intramolecular hydrogen bond and an intramolecular $\text{Au} \cdots \text{O}$ secondary bond is resolved in favour of the hydrogen bond. The $\text{H} \cdots \text{O1}$ and $\text{N} \cdots \text{O1}$ distances [1.95 (5) and 2.686 (4) Å, respectively] are comparable with those in the free molecule of (2). The $\text{Au}-\text{N}$ distance of 2.019 (3) Å is somewhat shorter than those in other AuPPH_3 derivatives of arylamines, *e.g.* 2.038 (6) (Kuz'mina, Burtseva *et al.*, 1989), 2.050 (6) (Perevalova *et al.*, 1988) and 2.046 Å (Kuz'mina *et al.*, 1990). We have shown previously that the $\text{Au}-\text{P}$ bond length is sensitive to a *trans* ligand in Au^{I} compounds (Kuz'mina, 1992). The $\text{Au}-\text{P}$ bond length of 2.2420 (8) Å is typical of Au^{I} compounds containing a $\text{P}-\text{Au}-\text{N}$ linear fragment.




Fig. 1. The molecular structure of (1) showing 50% probability displacement ellipsoids.

In the aminoanthraquinone fragment of (1), the bond-length distribution is slightly but significantly different from that observed in (2). In (1), the $\text{N}-\text{C1}$ bond [1.336 (4) Å] is shorter than in (2) [1.357 (4) Å found by Almond *et al.* (1983) and 1.358 Å reported by Foitzik & Paulus (1986)]. In addition, the amino-substituted ring shows some localization of double-bond character at $\text{C2}-\text{C3}$ [1.365 (5) Å] and $\text{C4}-\text{C12}$ [1.375 (5) Å]. Such a bond-length distribution, typical of the imino tautomeric form, is less pronounced in (2) [comparable bond lengths are 1.377 (4) and 1.384 (4) Å reported by Almond *et al.* (1983), and 1.371 and 1.385 Å reported by Foitzik & Paulus (1986)]. The observed reduction of the $\text{C}-\text{C}-\text{C}$ angle at atom C1 [116.7 (3)°] in (1) compared with those in (2) (118.4 and 118.5°) may be considered as an indication of the greater positive inductive effect of the AuPPH_3 substituent compared with hydrogen. A similar effect of some quinoid-like contribution to the general benzenoid structure is also observed in the above-mentioned tautomeric hydroxypyridine derivative. The remaining geometrical parameters of the ligand closely resemble those of (2).

The two $\text{C}-\text{O}$ bonds, 1.247 (4) and 1.229 (5) Å, differ by 0.018 (6) Å. The elongation of the $\text{O1}-\text{C9}$ bond is due to the involvement of O1 in the hydrogen bond.

In the crystal, the molecules are packed in stacks running along the *a* axis. In such a stack, any two adjacent molecules are related by a symmetry centre. None of the interatomic distances is less than the sum of the corresponding van der Waals radii. The mean interplanar distances between molecules related by two different symmetry centres (0,0,0 and $\frac{1}{2},0,0$) are 3.42 and 3.59 Å, respectively.

Experimental

The title compound was prepared by reaction of 1-aminoanthracene-9,10-dione with tris(triphenylphosphine)gold(I) oxonium tetrafluoroborate in the presence of aqueous potassium carbonate. Single crystals were grown from a chloroform-ether (1:7) solution.

Crystal data

$[\text{Au}(\text{C}_{14}\text{H}_8\text{NO}_2)(\text{C}_{18}\text{H}_{15}\text{P})]$

$M_r = 681.45$

Triclinic

$P\bar{1}$

$a = 8.8328$ (2) Å

$b = 9.4173$ (2) Å

$c = 16.1617$ (3) Å

$\alpha = 96.941$ (1)°

$\beta = 98.159$ (1)°

$\gamma = 101.396$ (1)°

$V = 1289.06$ (5) Å³

$Z = 2$

$D_x = 1.756$ Mg m⁻³

D_m not measured

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 9492

reflections

$\theta = 1.29-27.51^\circ$

$\mu = 5.799$ mm⁻¹

$T = 153$ (2) K

Block

$0.24 \times 0.18 \times 0.10$ mm

Red

Data collection

Siemens SMART diffractometer 5842 independent reflections
 5407 reflections with $I > 2\sigma(I)$
 ω scans $R_{\text{int}} = 0.028$
 Absorption correction: $\theta_{\text{max}} = 27.51^\circ$
 multi-scan (SHELXTL; $h = -11 \rightarrow 11$
 Sheldrick, 1994) $k = -8 \rightarrow 12$
 $T_{\text{min}} = 0.287$, $T_{\text{max}} = 0.560$ $l = -20 \rightarrow 20$
 9492 measured reflections

Refinement

Refinement on F^2 $\Delta\rho_{\text{max}} = 1.13 \text{ e } \text{\AA}^{-3}$
 $R[F^2 > 2\sigma(F^2)] = 0.025$ $\Delta\rho_{\text{min}} = -0.67 \text{ e } \text{\AA}^{-3}$
 $wR(F^2) = 0.064$ Extinction correction:
 $S = 1.120$ SHELXL93 (Sheldrick, 1993)
 5660 reflections Extinction coefficient:
 427 parameters 0.0000 (2)
 All H atoms refined Scattering factors from
 $w = 1/[\sigma^2(F_o^2) + (0.011P)^2$ International Tables for
 $+ 2.5925P]$ Crystallography (Vol. C)
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$

Table 1. Selected geometric parameters ($\text{\AA}$, $^\circ$)

Au—N	2.019 (3)	C3—C4	1.403 (6)
Au—P	2.2420 (8)	C4—C12	1.375 (5)
N—C1	1.336 (4)	C9—C11	1.453 (5)
N—H1	0.89 (5)	C9—C14	1.496 (5)
O1—C9	1.247 (4)	C10—C13	1.485 (6)
O2—C10	1.229 (5)	C10—C12	1.489 (5)
C1—C2	1.433 (5)	C11—C12	1.426 (5)
C1—C11	1.442 (5)	C13—C14	1.404 (5)
C2—C3	1.365 (6)		
N—Au—P	178.21 (9)	C11—C9—C14	119.0 (3)
C1—N—Au	128.0 (3)	O2—C10—C13	120.8 (4)
C1—N—H1	111 (3)	O2—C10—C12	121.3 (4)
Au—N—H1	122 (3)	C13—C10—C12	117.9 (3)
N—C1—C2	119.7 (3)	C12—C11—C1	119.2 (3)
N—C1—C11	123.6 (3)	C12—C11—C9	120.0 (3)
C2—C1—C11	116.7 (3)	C1—C11—C9	120.8 (3)
C3—C2—C1	122.1 (4)	C4—C12—C11	121.6 (3)
C2—C3—C4	121.0 (4)	C4—C12—C10	117.0 (3)
C12—C4—C3	119.4 (4)	C11—C12—C10	121.4 (3)
O1—C9—C11	123.1 (3)	C14—C13—C10	120.3 (3)
O1—C9—C14	117.9 (3)	C13—C14—C9	121.3 (3)

Crystal decay was checked during the data reduction and no decay was observed. The structure was solved by direct methods. H atoms were found from the difference Fourier synthesis and refined in the isotropic approximation. The largest peak in the difference electron-density map was found at a distance of 0.94 $\text{\AA}$ from the Au atom.

Data collection: SMART (Siemens, 1995a). Cell refinement: SAINT (Siemens, 1995b). Data reduction: SAINT. Program(s) used to solve structure: SHELXS86 (Sheldrick, 1990). Program(s) used to refine structure: SHELXL93 (Sheldrick, 1993). Molecular graphics: SHELXTL (Sheldrick, 1994). Software used to prepare material for publication: SHELXTL.

This work was supported by the Russian Foundation for Basic Research (project 95-03-08616a) and The Royal Society (UK).

Supplementary data for this paper are available from the IUCr electronic archives (Reference: BM1166). Services for accessing these data are described at the back of the journal.

References

- Almlöf, J., Kvik, Å. & Olovsson, I. (1971). *Acta Cryst.* **B27**, 1201–1208.
 Almond, P., Cutbush, S. D., Islam, S. A., Kuroda, R., Neidle, S., Gandeche, B. M. & Brown, J. R. (1983). *Acta Cryst.* **C39**, 627–630.
 Foitzik, J. K. & Paulus, H. (1986). *Mol. Cryst. Liq. Cryst. Lett. Sect.* **3**, 15–21.
 Kolb, A., Bissinger, P. & Schmidbaur, H. (1993). *Inorg. Chem.* **32**, 5132–5135.
 Kuz'mina, L. G. (1992). *Metalloorg. Khim.* **5**, 744–781.
 Kuz'mina, L. G., Burtseva, O. Yu., Dvortsova, N. V. & Porai-Koshits, M. A. (1989). *Metalloorg. Khim.* **2**, 1242–1247.
 Kuz'mina, L. G., Burtseva, O. Yu., Dvortsova, N. V., Porai-Koshits, M. A. & Smyslova, E. I. (1990). *Koord. Khim.* **16**, 70–74.
 Kuz'mina, L. G., Dvortsova, N. V., Porai-Koshits, M. A., Smyslova, E. I., Grandberg, K. I. & Perevalova, E. G. (1989). *Metalloorg. Khim.* **2**, 1344–1350.
 Perevalova, E. G., Struchkov, Yu. T., Kravtsov, D. N., Kuz'mina, L. G., Smyslova, E. I., Grandberg, K. I., Kalinina, O. N., Dyadchenko, V. P. & Voevodskaya, T. I. (1988). *Zh. Obshch. Khim.* **58**, 62–71.
 Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
 Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
 Sheldrick, G. M. (1994). *SHELXTL*. Release 5. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Siemens (1995a). *SMART*. Version 4.050. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Siemens (1995b). *SAINT*. Version 4.050. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Acta Cryst. (1998). **C54**, 214–217

3-(η^3 -Allyl)-3,3-dicarbonyl-4-(dimethyl sulfido)-3,1,2-molybdadicarba-closo-dodecaborane(12)

KJELD JOHANSEN, GEORGINA M. ROSAIR, ANDREW S. WELLER AND ALAN J. WELCH

Department of Chemistry, Heriot-Watt University, Riccarton, Edinburgh EH14 4AS, Scotland. E-mail: chegmr@bonaly.hw.ac.uk

(Received 22 May 1997; accepted 22 August 1997)

Abstract

The title charge-compensated allyl carbamolybdaborane 3-(η^3 -C₃H₅)-3,3-(CO)₂-4-SMe₂-3,1,2-closo-MoC₂B₉H₁₀, or [Mo(C₃H₅)(C₄H₁₆B₉S)(CO)₂], has been synthesized and its solid-state structure determined. In the solid state, the conformation of the {Mo(η^3 -C₃H₅)(CO)₂} fragment is such that the carbonyl ligands are approximately *trans* to the cage C atoms and the conformation of the allyl group is *exo*.