

tion-metal complexes with tertiary aminomethyl substituents³ may be attributed to the lack of an available β -hydride elimination pathway for the latter compounds.

Experimental Section

General Data. Pentane was dried over concentrated H_2SO_4 and fractionally distilled. Reagent grade dichloromethane was used as received. Reagent grade methanol and acetonitrile were dried over type 3A molecular sieves. Isopropylamine, CD_2Cl_2 , and CD_3CN were obtained from Aldrich and used directly. Et_4NBH_4 was prepared by a method described previously.⁸ The organometallic starting materials *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CHOCH}_3)^+\text{CF}_3\text{SO}_3^-$, *mer,trans*- $\text{Mn}(\text{CO})_3(\text{PPh}_3)_2(\text{CHNCH}_2\text{Ph})^+\text{CF}_3\text{SO}_3^-$, *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CHNHCH}_2\text{Ph})^+\text{CF}_3\text{SO}_3^-$, and *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CHNCH}_2\text{Ph})$ were prepared as previously reported.¹ Spectroscopic measurements were obtained on the following instruments: ^1H NMR, Varian XL-300 and EM-390; ^{13}C NMR, Varian XL-300; IR, Perkin-Elmer 599B and Nicolet SX170. ^1H and ^{13}C NMR chemical shifts were referenced to tetramethylsilane. Melting points were obtained on a Thomas-Hoover capillary melting point apparatus and are uncorrected. Elemental analyses were performed by Galbraith Laboratories, Knoxville, TN. All reactions were performed under an atmosphere of prepurified nitrogen.

***mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CH}_2\text{NHCH}_2\text{Ph})$ (2a).** (a) By Reduction of *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CHNHCH}_2\text{Ph})^+\text{CF}_3\text{SO}_3^-$ (1a). To a stirred solution of 1a (0.10 g, 0.094 mmol) in CH_3OH (10 mL) at room temperature was added Et_4NBH_4 (0.014 g, 0.094 mmol). After 5 min, a white precipitate began to form; after 30 min, the solvent was evaporated from the mixture. The residue was scraped from the sides of the flask and stirred with 30 mL of water for 5 min. The residue was then collected by filtration and dried, in vacuo, to yield 0.079 g (92%) of product, mp 159–160 °C. Efforts to recrystallize the product resulted in slight decomposition and lower yield; analytical data were obtained on the crude product. Anal. Calcd for $\text{C}_{47}\text{H}_{40}\text{NO}_3\text{P}_2\text{Re}$: C, 61.70; H, 4.41; N, 1.53. Found: C, 61.66; H, 4.46; N, 1.33. IR (CH_2Cl_2): ν_{CO} 2015 (vw), 1923 (s), 1884 (m) cm^{-1} . IR (neat, DRIFTS): ν_{NH} 3348 (w) cm^{-1} . ^1H NMR (CD_2Cl_2): δ 7.35 (m), 2.93 (s), 2.08 (t, $J_{\text{PH}} = 7.3$ Hz), 0.45 (s, br). $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_2Cl_2): δ 199.0 (t, $J_{\text{PC}} = 9.2$ Hz), 196.1 (t, $J_{\text{PC}} = 7.5$ Hz), 142.2 (s), 135.8 (s), $J_{\text{PC}} = 23.4$ Hz), 133.8 (t, $J_{\text{PC}} = 5.6$ Hz), 130.1 (s), 128.4 (s), 128.4 (t, $J_{\text{PC}} = 4.6$ Hz), 128.1 (s), 126.3 (s), 63.2 (s), 28.9 (t, $J_{\text{PC}} = 6.4$ Hz).

(b) By Reduction of *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CHNCH}_2\text{Ph})$. To a stirred solution of *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CHNCH}_2\text{Ph})$ (0.12 g, 0.13 mmol) in CH_2Cl_2 (10 mL) at 0 °C was added Et_4NBH_4 (0.019 g, 0.13 mmol). After 50 min, the volume of the solvent was reduced on a rotary evaporator, CH_3OH (20 mL) was added, and the solution was chilled to –20 °C to precipitate the product. White crystals were collected by filtration and dried for 2 h under vacuum to give 0.080 g (67%) of 2a. The spectral properties of the product were identical with those above.

***mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2[\text{CHNHCH}(\text{CH}_3)]^+\text{CF}_3\text{SO}_3^-$ (1b).** To 15 mL of CH_2Cl_2 containing *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CHOCH}_3)^+\text{CF}_3\text{SO}_3^-$ (0.40 g, 0.40 mmol) and maintained at 0 °C was added isopropylamine (34 μL , 0.40 mmol) dropwise from a syringe with stirring. The reaction was complete after 5 min at which time 10 mL of pentane was added, and the solution was chilled to –20 °C overnight. White crystals were collected by filtration, washed with 10 mL of pentane, and dried, in vacuo, to give 0.37 g (90%) of product, mp 213–214 °C dec. Anal. Calcd for $\text{C}_{44}\text{H}_{39}\text{F}_3\text{NO}_3\text{P}_2\text{ReS}$: C, 52.07; H, 3.87; N, 1.38. Found: C, 52.14; H, 3.86; N, 1.30. IR (CH_2Cl_2): ν_{CO} 2060 (w), 1960 (s), 1935 (sh) cm^{-1} . IR (neat, DRIFTS): ν_{NH} 3231 (m) cm^{-1} . ^1H NMR (CD_2Cl_2): δ 10.2 (d, br, $J_{\text{HH}} = 21.0$ Hz), 10.0 (d, $J_{\text{HH}} = 21.0$ Hz), 7.5 (m), 2.9 (hep, $J_{\text{HH}} = 6.6$ Hz), 0.55 (d, $J_{\text{HH}} = 6.6$ Hz). At this point, the broad doublet at lower field was assigned to the NH group while the sharper doublet at δ 10.0 was assigned to the isopropyl methine group. These assignments were confirmed by an NMR experiment using CD_3CN as the solvent. In CD_3CN , the doublet assigned to the methine proton becomes the lower

field resonance while the broad doublet assigned to the NH group moves slightly upfield. When a drop of D_2O was added to this NMR sample, the lower field doublet collapsed to a singlet and the other low-field doublet disappeared. $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_2Cl_2): δ 222.3 (t, $J_{\text{PC}} = 9.0$ Hz), 194.6 (t, $J_{\text{PC}} = 6.3$ Hz), 191.5 (t, $J_{\text{PC}} = 8.6$ Hz), 134.0 (t, $J_{\text{PC}} = 24.6$ Hz), 133.3 (t, $J_{\text{PC}} = 5.6$ Hz), 131.2 (s), 129.3 (t, $J_{\text{PC}} = 5.0$ Hz), 63.3 (s), 21.5 (s).

***mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2[\text{CH}_2\text{NHCH}(\text{CH}_3)_2]$ (2b).** To a stirred solution of 1b (0.30 g, 0.30 mmol) in CH_3CN (15 mL) at room temperature was added Et_4NBH_4 (0.043 g, 0.30 mmol). A white precipitate began to form immediately. After 5 min, the product was collected and dried in vacuo to yield 0.22 g (87%), mp 216 °C dec. Anal. Calcd for $\text{C}_{43}\text{H}_{40}\text{NO}_3\text{P}_2\text{Re}$: C, 59.57; H, 4.65; N, 1.62. Found: C, 58.89; H, 4.67; N, 1.82. IR (CH_2Cl_2): ν_{CO} 2015 (vw), 1920 (s), 1885 (m) cm^{-1} . IR (neat, DRIFTS): ν_{NH} 3290–3340 (vw, br) cm^{-1} . ^1H NMR (CD_2Cl_2 , –4 °C): δ 7.50 (m), 1.96 (t, $J_{\text{PH}} = 6.6$ Hz; the NH proton is thought to underlie this triplet as it is broadened; this assignment is supported by the integration), 1.51 (hep, $J_{\text{HH}} = 6.1$ Hz), 0.50 (d, $J_{\text{HH}} = 6.1$ Hz). $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_2Cl_2 , –4 °C): δ 198.8 (t, $J_{\text{PC}} = 9.4$ Hz), 195.4 (t, $J_{\text{PC}} = 6.8$ Hz), 135.4 (t, $J_{\text{PC}} = 23.2$ Hz), 133.6 (t, $J_{\text{PC}} = 5.4$ Hz), 129.9 (s), 128.3 (t, $J_{\text{PC}} = 4.6$ Hz), 56.6 (s), 24.7 (t, $J_{\text{PC}} = 5.9$ Hz), 22.0 (s).

Thermolysis of *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CH}_2\text{NHCH}_2\text{Ph})$ (2a). (a) A sample of 2a (0.025 g, 0.027 mmol) was placed in a Schlenk tube and stoppered. The sample was heated at 170 °C and then allowed to cool to 150 °C; after 2 h at 150 °C, the sample was cooled to room temperature. The flask was opened, and the residue was triturated with pentane. The major pentane-soluble product was identified as $\text{CH}_3\text{NHCH}_2\text{Ph}$ by comparison of the NMR spectrum with that of an authentic sample. The pentane-insoluble residue was chromatographed on a florisil column with CH_2Cl_2 /pentane (9:1). The first fractions off the column contained only *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2\text{H}$ (0.12 g, 57% yield) identified by comparison of its spectral properties with those of an authentic sample.⁹ Additional carbonyl-containing products were then eluted, in very small amounts; these products have not been identified.

(b) A second experiment was conducted by using equimolar amounts of 2a and *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2\text{H}$. Again, the major organic product was identified as $\text{CH}_3\text{NHCH}_2\text{Ph}$, and the major metal carbonyl product was *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2\text{H}$. No difference in product distributions were observed (after correction for the added hydride) in comparing this experiment with the one above.

Attempted Reductions of *mer,trans*- $\text{Mn}(\text{CO})_3(\text{PPh}_3)_2(\text{CHNHCH}_2\text{Ph})^+\text{CF}_3\text{SO}_3^-$ (1c). To 25 mL of CH_2Cl_2 containing 1c (0.50 g, 0.58 mmol) at room temperature was added Et_4NBH_4 (0.083 g, 0.58 mmol). After 5 min, the solvent was removed and the residue triturated with pentane. The pentane extracts smelled strongly of amine and showed no carbonyl stretching bands in the IR spectrum. A ^1H NMR study of the pentane-soluble portion in CD_2Cl_2 showed a complicated spectrum. However, the major product was determined to be $\text{CH}_3\text{NHCH}_2\text{Ph}$ by comparison of the NMR spectrum with that of the authentic sample. The pentane-insoluble product was dissolved in CH_2Cl_2 , the solution was mixed with decolorizing charcoal and filtered through Celite, and the volume of the solvent was reduced. Pentane was added until the solution became slightly cloudy, and this was chilled to –20 °C overnight. Pale yellow crystals of *mer,trans*- $\text{Mn}(\text{CO})_3(\text{PPh}_3)_2\text{H}$ were collected and dried, in vacuo, to yield 0.32 g, 82%. The hydride complex was identified by comparison of its spectral properties with those of an authentic sample.¹⁰ Doing this reaction at –78 °C gave similar results as did changing the solvent to CH_3OH or CH_3CN . When half of 1 molar equiv of Et_4NBH_4 was used, only half of the starting material reacted, but the same products were formed.

Thermolysis of *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CHNCH}_2\text{Ph})$. A sample of *mer,trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2(\text{CHNCH}_2\text{Ph})$ (0.073 g, 0.093 mmol) was placed in a Schlenk tube and stoppered. The sample was heated to 170 °C and then allowed to cool to 150 °C. This temperature was maintained for 2 h, during which time the

(8) Gibson, D. H.; Ahmed, F. U.; Phillips, K. R. *J. Organomet. Chem.* 1981, 218, 325.

(9) Flitcroft, N.; Leach, J. M.; Hopton, F. J. *J. Inorg. Nucl. Chem.* 1970, 32, 137.

(10) Hieber, W.; Duchatsch, H. *Chem. Ber.* 1965, 98, 2933.

white sample melted and became yellow. After 2 h, the sample was allowed to cool to room temperature. NMR and IR spectral data confirm that the major metal carbonyl product is *mer*-, *trans*- $\text{Re}(\text{CO})_3(\text{PPh}_3)_2\text{H}$ (by comparison of its spectral properties with those of an authentic sample).

Acknowledgment. Support of this work by the Na-

tional Science Foundation (Grant RII-8610671) and the Commonwealth of Kentucky (EPSCoR Program) is gratefully acknowledged. Partial support of this work by the Department of Energy, Division of Chemical Sciences (Office of Basic Energy Sciences), is also gratefully acknowledged.

Additions and Corrections

Jeffrey W. Freeman and Fred Basolo*: Kinetics and Mechanisms of Ligand Substitution Reactions of the Vanadium Triad Metals. Syntheses and Reactivities of $(\eta^5\text{-C}_5\text{H}_5)\text{M}(\text{CO})_3(\text{C}_4\text{H}_9\text{E})$ ($\text{M} = \text{Nb}$, $\text{E} = \text{S}$, Se , Te ; $\text{M} = \text{Ta}$, $\text{E} = \text{S}$). 1991, 10, 256–263.

On column 2, line 4 of page 256 the sentence beginning with "Equilibria studies..." should be replaced with "Studies of $\text{ReBr}(\text{CO})_3(\text{EMe}_2)_2$, $(\text{Cp-Me})\text{Mn}(\text{CO})_2\text{EMe}_2$, and $\text{CpV}(\text{CO})_3\text{EMe}_2$ showed¹¹ that the stability of these complexes increases in the order $\text{S} < \text{Se} < \text{Te}$." To ref 11 add the following: Belforte, A.; Calderazzo, F.; Vitali, D.; Zanazzi, P. F. *Gazz. Chim. Ital.* 1985, 115, 125.