

# Chemoenzymatic Cascades toward Methylated Tetrahydroprotoberberine and Protoberberine Alkaloids

Rebecca Roddan, Fabiana Subrizi, Joseph Broomfield, John M. Ward, Nicholas H. Keep, and Helen C. Hailes\*



Cite This: <https://doi.org/10.1021/acs.orglett.1c02110>



Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information



**ABSTRACT:** Tetrahydroprotoberberine and protoberberine alkaloids are a group of biologically active natural products with complex molecular scaffolds. Isolation from plants is challenging and stereoselective synthetic routes, particularly of methylated compounds are limited, reducing the potential use of these compounds. In this work, we describe chemoenzymatic cascades toward various 13-methyl-tetrahydroprotoberberine scaffolds using a stereoselective Pictet-Spenglerase, regioselective catechol O-methyltransferases and selective chemical Pictet-Spengler reactions. All reactions could be performed sequentially, without the workup or purification of any synthetic intermediates. Moreover, the naturally occurring alkaloids have the (+)-configuration and importantly here, a strategy to the (–)-isomers was developed. A methyl group at C-8 was also introduced with some stereocontrol, influenced by the stereochemistry at C-13. Furthermore, a single step reaction was found to convert tetrahydroprotoberberine alkaloids into the analogous protoberberine scaffold, avoiding the use of harsh oxidizing conditions or a selective oxidase. This work provides facile, selective routes toward novel analogues of bioactive alkaloids.

Natural products and related analogues are a major source of therapeutics, making up 42% of FDA-approved drugs from 1981 to 2019.<sup>1</sup> Many have a diverse range of molecular scaffolds, often with multiple, defined stereocenters. However, accessing natural products is challenging and different approaches are used with varying successes. For example, isolation from plant sources is hindered by low production and issues with separation from other structurally similar metabolites.<sup>2</sup> Total synthesis is commercially viable in some cases, although can be unattainable or not cost-effective for more complex products.<sup>3</sup> *In vivo* fermentation processes involve significant bioengineering efforts and achieving high enantiopurities can be challenging.<sup>4,5</sup> High regio- and stereoselective control under benign conditions, with minimal side-reactions can be achieved however by mimicking the biosynthesis *in vitro*, using recombinantly expressed enzymes.<sup>6</sup> Nevertheless, this approach can be limited by challenges such as enzyme stability issues. Cascade processes, using a combination of traditional synthetic methods and biocatalytic enzymes, can alternatively lead to the generation of complex products, in fewer steps with easier purification processes than solely *in vivo* or organic synthetic routes.<sup>7–9</sup>

Alkaloids are important nitrogen-containing natural products, many of which are biologically active.<sup>10</sup> Protoberberine (PB) and tetrahydroprotoberberine (THPB) alkaloids isolated from plants of the *Corydalis* genus are unique among other


isolated alkaloids of this type as they possess a methyl group at C-13.<sup>11,12</sup> Many 13-Me-PB and 13-Me-THPB alkaloids have been shown to have promising bioactivities,<sup>13</sup> including as inhibitors of reverse transcriptase activity<sup>14</sup> and enterovirus 71.<sup>15</sup> Others have been shown to be dopamine D-1 receptor agonists and to have antihepatitis B activities.<sup>16,17</sup>

Synthetic routes toward racemic 13-Me-PB and 13-Me-THPB alkaloids have been reported,<sup>18–20</sup> and between them only a few are stereoselective. The asymmetric synthesis of natural 13-Me-THPBs has been described for example by Zhou et al. (Scheme 1a).<sup>21,22</sup> After the four-step synthesis of an enantiopure PINAP ligand, the (13*R*,13*aS*)-13-Me-THPBs were synthesized in three-steps in yields of 47–65% and high selectivities (91–96% ee). Previous work in our group has shown that the Pictet-Spenglerase<sup>23</sup> norcoclaurine synthase (NCS) can accept  $\alpha$ -methyl-substituted aldehydes as substrates.<sup>24</sup> Notably, an active site variant of *Thalictrum flavum* NCS (*Tf*NCS) M97V gave (1*S*,1'*R*)-tetrahydroisoquinolines


Received: June 25, 2021

**Scheme 1. Routes towards 13-Me-THPB alkaloids (a) Previous route using a chiral catalyst;<sup>21</sup> (b) This work using a chemoenzymatic approach**

**a) Previous work:**




**b) This work:**



(THIQs) in high yields (up to 96%) and diastereomeric ratios (d.r. = 98:2) in a single-step. Herein, the application of this methodology in combination with other biocatalytic or chemical steps is described to generate a range of 13-Me-THPBs isolated from *Corydalis* plants. Importantly, the products generated have opposing stereochemistry to those isolated from plants, thus providing routes to useful natural product analogues (Scheme 1b).

To generate the desired THIQ scaffold (Scheme 2), NCS-mediated reactions between dopamine **1a** and the aldehyde

**Scheme 2. Stereoselective Synthesis of the THIQ Scaffold Using Norcoclaurine Synthase**



<sup>a</sup>d.r. values correspond to the ratio of the two diastereomers formed, (1*S*,1'*R*):(1*S*,1'*S*) and were determined by <sup>1</sup>H NMR spectroscopy and HPLC (method 2). <sup>b</sup>Conversions were determined by HPLC analysis, based upon calibration curves of the purified products (see SI). <sup>c</sup>Isolated yields after preparative HPLC purification.


**2**<sup>24,25</sup> were investigated. Variants of TfNCS, M97V, L76V, and M97F, which previously gave improved selectivities compared with the wild type for the acceptance of  $\alpha$ -methyl phenylacetaldehyde, were explored (Supporting Information (SI) Figure S6): when this racemic aldehyde had been used in NCS catalyzed reactions, the *R*-enantiomer was accepted preferentially over the *S*-isomer.<sup>24</sup> This had been determined by performing reactions with single enantiomer  $\alpha$ -methyl

aldehydes. Notably, it resulted in the generation of two defined chiral centers in the resulting THIQs, with the major product assigned as (1*S*,1'*R*).<sup>24</sup> Aldehyde **2** was prepared in three steps<sup>25</sup> and after the final step taken through without purification due to its oxidative sensitivity. TfNCS-M97V gave the THIQ products with **2** in quantitative yield by HPLC analysis, with the major diastereomer generated assigned as (1*S*,1'*R*)-**3a** (d.r. = 96:4).<sup>24</sup> The minor diastereomer observed was (1*S*,1'*S*)-**3a**, as no racemic NCS background reaction was observed and NCS has been shown to generate the *S*-stereochemistry at C-1.<sup>26,27</sup> Compound **3a** was isolated by preparative HPLC (with the reaction performed on a 0.10 mmol scale, with 20 mg **1a**) for characterization purposes but otherwise was taken through directly for the cascade process to telescope the synthetic approach. While these initial experiments used 2 equiv of **2**, similar results and slightly higher stereoselectivities were observed when using just 1 equiv (>99% conversion, 18% isolated, d.r. = 97:3) as **2** can racemize *in situ*.<sup>28</sup> The challenges of THIQ product isolation have previously been reported<sup>29</sup> which can lower isolated yields, highlighting the advantages of directly taking material through to the next step using cascaded reaction sequences which is described below. This reaction was also amenable to scale up with conversions and stereoselectivities retained on a 50 mL, 10 mM scale.

The *para*-hydroxyl group of dopamine **1a** is nonessential for a productive NCS reaction,<sup>30</sup> so the 7-OMe THIQ **3b** was also generated using a TfNCS-M97V reaction between **2** and the 7-OMe dopamine analogue, **1b**,<sup>26</sup> on a 0.10 mmol scale (with 20 mg of **1b**). This reduced the oxidative sensitivity of the THIQ scaffold. The product, (1*S*,1'*R*)-**3b** (Scheme 2), was generated in high yield (>99% HPLC conversion, 81% isolated) and in a reasonable d.r. (92:8).

To generate the analogous C6-OMe-THIQ (**3c**), use of regioselective *O*-methyltransferases (*O*-MTs) were explored. Previous work has shown that two promising *O*-MTs, RnCOMT (isolated from *Rattus norvegicus*) and MxSafC (isolated from *Myxococcus xanthus*), are capable of regioselectively methylating the C6-OH or C7-OH of various 1-benzylic and 1-aryl-THIQs.<sup>26,31–34</sup> Both are *S*-adenosyl-L-methionine (SAM) dependent enzymes, and although SAM is expensive and has issues of instability, recent developments in SAM supply systems have meant that such biocatalytic methylations are viable on preparative scales.<sup>35,36</sup> Here, we used a previously described *in situ* SAM generation system which utilizes ATP, L-methionine, and a methionine adenosyl transferase from *Escherichia coli* (EcMAT E.C.2.5.1.6). After the methylation reaction, *S*-adenosyl-L-homocysteine (SAH) is generated which can inhibit the *O*-MT, so another enzyme methylthioadenosine/SAH nucleosidase (EcMTAN E.C.3.2.2.9) is used to breakdown SAH.<sup>35,37</sup> The use of other SAM generation systems was not attempted, but other *in situ* SAM supply methods could be applied.<sup>38,39</sup>

The THIQ substrate **3a** formed (using TfNCS-M97V) was directly lyophilized, and since the *O*-MT enzymes are known to be highly selective toward the catechol moiety and no **1a** remained after the NCS reaction, there was no need for a purification step. Using previously reported conditions,<sup>26,31</sup> RnCOMT was used as clarified cell lysate and MxSafC was used as a purified enzyme. Both *O*-MTs interestingly exhibited high regioselectivity toward the 6-OH, generating the product **3c** (Scheme 3) with complete conversions by HPLC analysis and no observable methylation on the 7-OH.

Scheme 3. Regioselective Methylations, Chemical Pictet–Spengler Reactions and Generation of the 13-Me-PB Scaffold<sup>a,b,c,d</sup>

<sup>a</sup>Conversions were determined by HPLC analysis against product standards. <sup>b</sup>Conversions were determined by HPLC analysis, based upon starting material depletion (see calibration curves SI, Figures S7–S9). <sup>c</sup>Isolated yields after preparative HPLC purification. <sup>d</sup>Epimeric ratio determined based upon previous studies in which the stereochemistries at C-13 and C-13a were determined, and NOESY <sup>1</sup>H NMR analysis of the isolated product (see SI). Isolated yields were lower than HPLC yields/conversions due to issues of oxidative sensitivity of the compounds generated and minor impurities having similar retention times to the desired products by preparative HPLC. To obtain the compounds in high purity for characterization purposes, isolated yields are lower than those typically reported.

Reactions were performed on a 0.10 mmol scale (20 mg **1a**). Since both enzymes exhibited similar reactivities, yet *RnCOMT* could be used as clarified cell lysate, *RnCOMT* was used in subsequent reactions. Regioselectivities were established using Nuclear Overhauser Effect Spectroscopy (NOESY). This result is consistent with the reported regioselectivity of these enzymes toward THIQs with a phenyl or cyclohexane ring at C-1.<sup>26,31</sup> However, with substrates such as dopamine, *RnCOMT* methylates the *meta*-OH and *MxSafC* the *para*-OH.<sup>36</sup> Such variations in the regioselectivity highlights that bulkier groups attached to C-1 can hinder methylation at the C7-OH position in THIQs for steric reasons. Reactions were performed on a preparative scale (20 mL, 5 mM) using *RnCOMT* to give (1*S*,1'*R*)-**3c** in quantitative conversion by HPLC against product standards and a 12% isolated yield.

Biosynthetic routes to give the THPB scaffold involve the berberine bridge enzyme (BBE).<sup>40</sup> The enantioselective production of various (*S*)-THPBs from racemic, *N*-methylated, 1-benzylic THIQs using recombinant BBE has been achieved in high conversions (98%) and selectivities (>97% ee).<sup>41</sup> Although a highly productive route, there is the requirement to generate the starting material and *in situ* deracemization of the (*R*)-THIQ is needed, involving a selective monoamine oxidase and borane reduction. Here, to generate the tetracyclic THPB scaffold, chemical Pictet–Spengler (PS) reactions were explored (Scheme 3a).<sup>42</sup> Previous work has demonstrated that a phosphate-mediated PS reaction with formaldehyde was capable of converting 1-benzylic-THIQs into THPBs in high yields and good regioselectivities (7:1 10,11-OMe:9,10-OMe THPB).<sup>43</sup> Both reported reaction conditions were attempted here with **3c**, but no conversion was observed, presumably because the aromatic dimethoxy groups reduce the reactivity of the substrates compared to phenolic groups. Instead, formic acid catalyzed PS reactions were explored as described by Qian et al.<sup>44</sup> A telescoped synthesis was then performed, with crude, lyophilized **3a** or **3c** used. Complete conversion, by monitoring

the consumption of starting material by HPLC to the products (13*S*,13*aR*)-**4a** and (13*S*,13*aR*)-**4b**, was observed with 24% and 23% isolated yields, over two or three steps, respectively. Reactions were performed on a 5 mL, 10 mM scale. Complete regioselectivity was observed on the D-ring (see Scheme 1), with solely 10,11-dimethoxy-substituted products generated. Nuclear Overhauser Spectroscopy (NOESY) analysis of the products formed (**4a** and **4b**) also provided further confirmation of the stereochemistries at C-13 and C-13a, i.e., a *syn*-relationship between the two protons at these positions.

8-Me-THPB and 8-Me-PB alkaloids have been isolated from *C. ochotensis*, and one 8-Me-PB has been shown to act as an antileukemic agent.<sup>45</sup> Stereoselective routes to 8-Me THPBs are limited,<sup>46</sup> and routes to analogues are useful for drug discovery purposes. A chemical PS reaction in formic acid with acetaldehyde was therefore explored, with the aim of generating functionalized 8-THPBs. Crude, lyophilized **3c** was used again in a telescoped synthesis (0.050 mmol scale). Product **4c** was formed in a 56% conversion yield (starting material consumption by HPLC) and 21% isolated yield over two steps (Scheme 3). Two epimers at C-8 of the product were observed in a 3:1 ratio by NMR spectroscopy. The stereochemistries at C-13 and C-13a were retained, and the major epimer was assigned by NOESY analysis as (8*S*,13*R*,13*aS*)-**4c**. Modeling of both epimers was used to rationalize these results. In both cases, the C-ring is held in a half-chair conformation and, for the minor isomer, the two methyl groups are *cis* to each other in pseudoaxial orientations giving an unfavorable steric interaction (SI Figure S10). Therefore, the stereochemistry at C-13 can lead to stereocontrol at C-8 in the cyclization to give **4c**.

The regioselectivity of D-ring formation has been reported to be solvent-mediated,<sup>47,48</sup> with the other regioisomer of product preferentially formed in apolar, aprotic solvents such as toluene or dichloroethane. As **3c** has poor solubilities in such solvents, DMF was used with reactions performed at 120 °C on a 0.10 mmol scale (20 mg **1a**). Complete consumption

of **3c** was observed after 18 h, and interestingly **5** was formed via a cyclization and oxidation in 5% isolated yield (over three steps from **1a** and **2** with no purification of the intermediates). The oxidation is presumably mediated by trace oxidants in the crude material taken through and may prove useful in generating the PB scaffold from the analogous THPB, as the oxidation of the C-ring must occur after the PS reaction. It would seem that the presence of activating methoxy groups on the D-ring and the use of high temperatures also help to drive the reaction. The unpurified THPB **4b** (formed from **3c**) was also heated in DMF at 120 °C for 18 h with 86% of starting material converted by HPLC, and this gave **5** as the only product in 21% isolated yield. This is a useful route to PBs from the analogous THPB, avoiding the addition of oxidase enzymes<sup>49</sup> or oxidants (I<sub>2</sub>/EtOH, reflux).<sup>50</sup> Since the stereochemistry is lost during this oxidation, the racemic THIQ starting material could also be formed by a phosphate-mediated PS reaction which has been shown to be regioselective with  $\alpha$ -methyl substituted aldehydes.<sup>24</sup> The PB scaffold is synthetically useful, as the iminium ion is susceptible to nucleophilic attack, leading to C-8 functionalization.<sup>51</sup>

In summary, a range of novel (–)-13-Me-THPB alkaloids have been generated with opposing stereochemistries at C-13 and C-13a to the naturally occurring 13-Me-THPB alkaloids from *Corydalis* plants. Single regio- and diastereomer 13-Me-THPBs were formed through two subsequent PS reactions: one enzymatic reaction to generate a THIQ scaffold with two defined stereocenters followed by a chemical PS reaction to generate the tetracyclic scaffold. Regioselective methylation of a single hydroxyl group of the THIQ was also achieved using O-MTs. Heating the 13-Me-THPBs in DMF provided a facile route to the analogous 13-Me-PB. All enzymatic reactions were high yielding, and there was no requirement for the isolation of material at each reaction step. High stereoselectivity was also obtained without the need for chiral ligands or precursors, and toxic reagents.

## ■ ASSOCIATED CONTENT

### SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.1c02110>.

Experimental procedures and <sup>1</sup>H/<sup>13</sup>C NMR spectra for the compounds synthesized (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

Helen C. Hailes – Department of Chemistry, Christopher Ingold Building, University College London, London WC1H 0AJ, U.K.; [orcid.org/0000-0001-5574-4742](https://orcid.org/0000-0001-5574-4742);  
Email: [h.c.hailes@ucl.ac.uk](mailto:h.c.hailes@ucl.ac.uk)

### Authors

Rebecca Roddan – Department of Biological Sciences, Institute of Structural and Molecular Biology, Birkbeck College, London WC1E 7HX, U.K.; Department of Chemistry, Christopher Ingold Building, University College London, London WC1H 0AJ, U.K.

Fabiana Subrizi – Department of Chemistry, Christopher Ingold Building, University College London, London WC1H 0AJ, U.K.

Joseph Broomfield – Department of Chemistry, Christopher Ingold Building, University College London, London WC1H 0AJ, U.K.

John M. Ward – Department of Biochemical Engineering, Bernard Katz Building, University College London, London WC1E 6BT, U.K.

Nicholas H. Keep – Department of Biological Sciences, Institute of Structural and Molecular Biology, Birkbeck College, London WC1E 7HX, U.K.; [orcid.org/0000-0002-5042-1837](https://orcid.org/0000-0002-5042-1837)

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.orglett.1c02110>

## Author Contributions

The manuscript was written through contributions of all authors. R.R. and J.B. performed chemical syntheses. R.R. performed synthesis and characterization, enzyme expression and purification, and the chemoenzymatic cascades. F.S. provided input on enzymatic reactions. The project was supervised by H.C.H, N.H.K., and J.M.W. The manuscript was written by R.R. and H.C.H. All authors have given approval to the final manuscript.

## Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

The authors would like to thank Birkbeck College, for a Birkbeck Anniversary PhD scholarship to R.R. Also, the Biotechnology and Biological Sciences Research Council (BBRSC) (BB/R021643/1) and 17-ERACoBioTech are thanked for financial support of this work to F.S. as part of the project BioDiMet. We also thank Jennifer N. Alexander (Institute of Pharmaceutical Sciences, University of Freiburg) and Michael Richter (Fraunhofer Institute for Interfacial Engineering and Biotechnology, Straubing) for MTs (*Rn*COMT, *Mx*SafC), *Ec*MAT and *Ec*MTAN plasmids. Furthermore, we gratefully thank the UCL Mass Spectrometry and NMR Facilities in the Department of Chemistry and EPSRC (EP/P020410/1) for 700 MHz NMR equipment support.

## ■ REFERENCES

- (1) Newman, D. J.; Cragg, G. M. Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019. *J. Nat. Prod.* **2020**, *83*, 770–803.
- (2) Sasidharan, S.; Chen, Y.; Saravanan, D.; Sundram, K. M.; Latha, L. Y. Extraction, Isolation and Characterization of Bioactive Compounds from Plants' Extracts. *Afr. J. Tradit., Complementary Altern. Med.* **2010**, *8*, 1–10.
- (3) Baran, P. S. Natural Product Total Synthesis: As Exciting as Ever and Here to Stay. *J. Am. Chem. Soc.* **2018**, *140*, 4751–4755.
- (4) Hawkins, K. M.; Smolke, C. D. Production of Benzylisoquinoline Alkaloids in *Saccharomyces cerevisiae*. *Nat. Chem. Biol.* **2008**, *4*, 564–573.
- (5) Trenchard, I. J.; Siddiqui, M. S.; Thodey, K.; Smolke, C. D. De Novo Production of the Key Branch Point Benzylisoquinoline Alkaloid Reticuline in Yeast. *Metab. Eng.* **2015**, *31*, 74–83.
- (6) Wang, Y.; Tappertzhofen, N.; Méndez-Sánchez, D.; Bawn, M.; Lyu, B.; Ward, J. M.; Hailes, H. C. Design and Use of de Novo Cascades for the Biosynthesis of New Benzylisoquinoline Alkaloids. *Angew. Chem., Int. Ed.* **2019**, *58*, 10120–10125.

- (7) Li, J.; Amatuni, A.; Renata, H. Recent Advances in the Chemoenzymatic Synthesis of Bioactive Natural Products. *Curr. Opin. Chem. Biol.* **2020**, *55*, 111–118.
- (8) Zhao, J.; Lichman, B. R.; Ward, J. M.; Hailes, H. C. One-Pot Chemoenzymatic Synthesis of Trolline and Tetrahydroisoquinoline Analogues. *Chem. Commun.* **2018**, *54*, 1323–1326.
- (9) Pressnitz, D.; Fischereder, E.; Pletz, J.; Kofler, C.; Hammerer, L.; Hiebler, K.; Lechner, H.; Richter, N.; Eger, E.; Kroutil, W. Asymmetric Synthesis of (*R*)-1-Alkyl Substituted Tetrahydro- $\beta$ -Carbolines Catalyzed by Strictosidine Synthases. *Angew. Chem., Int. Ed.* **2018**, *57*, 10683–10687.
- (10) Kishimoto, S.; Sato, M.; Tsunematsu, Y.; Watanabe, K. Evaluation of Biosynthetic Pathway and Engineered Biosynthesis of Alkaloids. *Molecules* **2016**, *21*, 1078–1096.
- (11) Manske, R. H. F. The Alkaloids of Fumariaceae Plants. XLIX. Thalictricavine, A New Alkaloid from *Corydalis Tuberosa* DC. *J. Am. Chem. Soc.* **1953**, *75*, 4928–4929.
- (12) Yu, C. K.; MacLean, D. B.; Rodrigo, R. G. A.; Manske, R. H. F. Structural and Conformational Studies on Tetrahydroprotoberberine Alkaloids. *Can. J. Chem.* **1970**, *48*, 3673–3678.
- (13) Chow, Y. L.; Sogame, M.; Sato, F. 13-Methylberberine, a Berberine Analogue with Stronger Anti-Adipogenic Effects on Mouse 3T3-L1 Cells. *Sci. Rep.* **2016**, *6*, 38129.
- (14) Sethi, M. L. Enzyme Inhibition VI: Inhibition of Reverse Transcriptase Activity by Protoberberine Alkaloids and Structure–Activity Relationships. *J. Pharm. Sci.* **1983**, *72*, 538–541.
- (15) Wang, H. Q.; Hu, J.; Yan, H. Y.; Wu, S.; Li, Y. H. Corydaline Inhibits Enterovirus 71 Replication by Regulating COX-2 Expression. *J. Asian Nat. Prod. Res.* **2017**, *19*, 1124–1133.
- (16) Li, H. L.; Han, T.; Liu, R. H.; Zhang, C.; Chen, H. S.; Zhang, W. D. Alkaloids from *Corydalis Saxicola* and Their Anti-Hepatitis B Virus Activity. *Chem. Biodiversity* **2008**, *5*, 777–783.
- (17) Wu, L.; Zhang, W.; Qiu, X.; Wang, C.; Liu, Y.; Wang, Z.; Yu, Y.; Ye, R. D.; Zhang, Y. Identification of Alkaloids from *Corydalis yanhusuo* W. T. Wang as Dopamine D<sub>1</sub> Receptor Antagonists by Using CRE-Luciferase Reporter Gene Assay. *Molecules* **2018**, *23*, 2585.
- (18) Saá, C.; Guitián, E.; Castedo, L.; Suau, R.; Saá, J. M. A Regioselective Entry to 13-Substituted 8-Oxoprotoberberines. Total Synthesis of ( $\pm$ )-Corydaline. *J. Org. Chem.* **1986**, *51*, 2781–2784.
- (19) Cushman, M.; Dekow, F. W. A Total Synthesis of Corydaline. *Tetrahedron* **1978**, *34*, 1435–1439.
- (20) Hanaoka, M.; Yoshida, S.; Mukai, C. A Novel and Efficient Synthesis of 13-Methylprotoberberine Alkaloids. *J. Chem. Soc., Chem. Commun.* **1985**, *745*, 1257–1258.
- (21) Zhou, S.; Tong, R. Three-Step Catalytic Asymmetric Total Syntheses of 13-Methyltetrahydroprotoberberine Alkaloids. *Org. Lett.* **2017**, *19*, 1594–1597.
- (22) Mengozzi, L.; Gualandi, A.; Cozzi, P. G. A Highly Enantioselective Acyl-Mannich Reaction of Isoquinolines with Aldehydes Promoted by Proline Derivatives: An Approach to 13-Alkyl-Tetrahydroprotoberberine Alkaloids. *Chem. Sci.* **2014**, *5*, 3915–3921.
- (23) Roddan, R.; Ward, J. M.; Keep, N. H.; Hailes, H. C. Pictet–Spenglerases in Alkaloid Biosynthesis: Future Applications in Biocatalysis. *Curr. Opin. Chem. Biol.* **2020**, *55*, 69–76.
- (24) Roddan, R.; Gygli, G.; Sula, A.; Méndez-Sánchez, D.; Pleiss, J.; Ward, J. M.; Keep, N. H.; Hailes, H. C. Acceptance and Kinetic Resolution of  $\alpha$ -Methyl-Substituted Aldehydes by Norcoclaurine Synthases. *ACS Catal.* **2019**, *9*, 9640–9649.
- (25) Chang, D.-J.; An, H.; Kim, K.-S.; Kim, H. H.; Jung, J.; Lee, J. M.; Kim, N.-J.; Han, Y. T.; Yun, H.; Lee, S.; Lee, J. S.; Cha, J.-H.; Park, J.-H.; Park, J. W.; Lee, S.-C.; Kim, S. G.; Kim, J. H.; Lee, H.-Y.; Kim, K.-W.; Suh, Y.-G. Design, Synthesis, and Biological Evaluation of Novel Deguelin-Based Heat Shock Protein 90 (HSP90) Inhibitors Targeting Proliferation and Angiogenesis. *J. Med. Chem.* **2012**, *55*, 10863–10884.
- (26) Roddan, R.; Sula, A.; Méndez-Sánchez, D.; Subrizi, F.; Lichman, B. R.; Broomfield, J.; Richter, M.; Andexer, J. N.; Ward, J. M.; Keep, N. H.; Hailes, H. C. Single Step Syntheses of (1*S*)-Aryl-Tetrahydroisoquinolines by Norcoclaurine Synthases. *Commun. Chem.* **2020**, *3*, 170.
- (27) Pesnot, T.; Gershater, M. C.; Ward, J. M.; Hailes, H. C. The Catalytic Potential of *Coptis japonica* NCS2 Revealed - Development and Utilisation of a Fluorescamine-Based Assay. *Adv. Synth. Catal.* **2012**, *354*, 2997–3008.
- (28) Fuchs, C. S.; Hollauf, M.; Meissner, M.; Simon, R. C.; Besset, T.; Reek, J. N. H.; Riethorst, W.; Zepeck, F.; Kroutil, W. Dynamic Kinetic Resolution of 2-Phenylpropanal Derivatives to Yield  $\beta$ -Chiral Primary Amines via Bioamination. *Adv. Synth. Catal.* **2014**, *356*, 2257–2265.
- (29) Vanden Eynden, M. J.; Kunchithapatham, K.; Stambuli, J. P. Calcium-Promoted Pictet–Spengler Reactions of Ketones and Aldehydes. *J. Org. Chem.* **2010**, *75*, 8542–8549.
- (30) Ruff, B. M.; Bräse, S.; O'Connor, S. E. Biocatalytic Production of Tetrahydroisoquinolines. *Tetrahedron Lett.* **2012**, *53*, 1071–1074.
- (31) Subrizi, F.; Wang, Y.; Thair, B.; Méndez-Sánchez, D.; Roddan, R.; Cárdenas-Fernández, M.; Siegrist, J.; Richter, M.; Andexer, J. N.; Ward, J. M.; Hailes, H. C. Multienzyme One-pot Cascades Incorporating Methyltransferases for the Strategic Diversification of Tetrahydroisoquinoline Alkaloids. *Angew. Chem., Int. Ed.* **2021**, *60*, DOI: 10.1002/anie.202104476.
- (32) Nelson, J. T.; Lee, J.; Sims, J. W.; Schmidt, E. W. Characterization of SafC, a Catechol 4-O-Methyltransferase Involved in Saframycin Biosynthesis. *Appl. Environ. Microbiol.* **2007**, *73*, 3575–3580.
- (33) Creveling, C. R.; Morris, N.; Shimizu, H.; Ong, H. H.; Daly, J. Catechol O-Methyltransferase. *Mol. Pharmacol.* **1972**, *8*, 398–409.
- (34) Vidgren, J.; Svensson, L. A.; Liljas, A. Crystal Structure of Catechol O-Methyltransferase. *Nature* **1994**, *368*, 354–358.
- (35) Mordhorst, S.; Siegrist, J.; Müller, M.; Richter, M.; Andexer, J. N. Catalytic Alkylation Using a Cyclic S-Adenosylmethionine Regeneration System. *Angew. Chem., Int. Ed.* **2017**, *56*, 4037–4041.
- (36) Siegrist, J.; Aschwanden, S.; Mordhorst, S.; Thöny-Meyer, L.; Richter, M.; Andexer, J. N. Regio-complementary O-Methylation of Catechols by Using Three-Enzyme Cascades. *ChemBioChem* **2015**, *16*, 2576–2579.
- (37) Siu, K. K. W.; Asmus, K.; Zhang, A. N.; Horvatin, C.; Li, S.; Liu, T.; Moffatt, B.; Woods, V. L.; Howell, P. L. Mechanism of Substrate Specificity in 5'-Methylthioadenosine/S-Adenosylhomocysteine Nucleosidases. *J. Struct. Biol.* **2011**, *173*, 86–98.
- (38) Liao, C.; Seebeck, F. P. S-Adenosylhomocysteine as a Methyl Transfer Catalyst in Biocatalytic Methylation Reactions. *Nat. Catal.* **2019**, *2*, 696–701.
- (39) Sadler, J. C.; Humphreys, L. D.; Snajdrova, R.; Burley, G. A. A Tandem Enzymatic sp<sup>2</sup>-C-Methylation Process: Coupling in Situ S-Adenosyl-L-Methionine Formation with Methyl Transfer. *ChemBioChem* **2017**, *18*, 992–995.
- (40) Kutchan, T. M.; Dittrich, H. Characterization and Mechanism of the Berberine Bridge Enzyme, a Covalently Flavinyllated Oxidase of Benzophenanthridine Alkaloid Biosynthesis in Plants. *J. Biol. Chem.* **1995**, *270*, 24475–24481.
- (41) Schrittwieser, J. H.; Groenendaal, B.; Resch, V.; Ghislieri, D.; Wallner, S.; Fischereder, E.-M.; Fuchs, E.; Grischek, B.; Sattler, J. H.; Macheroux, P.; Turner, N. J.; Kroutil, W. Deracemization by Simultaneous Bio-Oxidative Kinetic Resolution and Stereo-inversion. *Angew. Chem., Int. Ed.* **2014**, *53*, 3731–3734.
- (42) Pesnot, T.; Gershater, M. C.; Ward, J. M.; Hailes, H. C. Phosphate Mediated Biomimetic Synthesis of Tetrahydroisoquinoline Alkaloids. *Chem. Commun.* **2011**, *47*, 3242–3244.
- (43) Lichman, B. R.; Lamming, E. D.; Pesnot, T.; Smith, J. M.; Hailes, H. C.; Ward, J. M. One-Pot Triangular Chemoenzymatic Cascades for the Syntheses of Chiral Alkaloids from Dopamine. *Green Chem.* **2015**, *17*, 852–855.
- (44) Qian, W.; Lu, W.; Sun, H.; Li, Z.; Zhu, L.; Zhao, R.; Zhang, L.; Zhou, S.; Zhou, Y.; Jiang, H.; Zhen, X.; Liu, H. Design, Synthesis, and Pharmacological Evaluation of Novel Tetrahydroprotoberberine

Derivatives : Selective Inhibitors of Dopamine D 1 Receptor. *Bioorg. Med. Chem.* **2012**, *20*, 4862–4871.

(45) Zee-Cheng, K. Y.; Paull, K. D.; Cheng, C. C. Experimental Antileukemic Agents. Coralyne, Analogs, and Related Compounds. *J. Med. Chem.* **1974**, *17*, 347–351.

(46) Baird, P. D.; Blagga, J.; Davies, S. G.; Sutton, K. H. The Stereospecific Synthesis of (–)-(8R) and (–)-8S-Methylcanadine. *Tetrahedron* **1988**, *44*, 171–186.

(47) Horst, B.; Wanner, M. J.; Jørgensen, S. I.; Hiemstra, H.; van Maarseveen, J. H. Total Synthesis of the Ortho-Hydroxylated Protoberberines (S)-Govaniadine, (S)-Caseamine, and (S)-Clarkeanidine via a Solvent-Directed Pictet–Spengler Reaction. *J. Org. Chem.* **2018**, *83*, 15110–15117.

(48) McMurtrey, K. D.; Meyerson, L. R.; Cashaw, J. L.; Davis, V. E. Kinetics and Product Distribution in Pictet–Spengler Cyclization of Tetrahydropapaveroline to Tetrahydroprotoberberine Alkaloids. *J. Org. Chem.* **1984**, *49*, 947–948.

(49) Daniel, B.; Konrad, B.; Toplak, M.; Lahham, M.; Messenlehner, J.; Winkler, A.; Macheroux, P. The Family of Berberine Bridge Enzyme-like Enzymes: A Treasure-Trove of Oxidative Reactions. *Arch. Biochem. Biophys.* **2017**, *632*, 88–103.

(50) Suau, R.; Silva, M. V.; Valpuesta, M. A Novel Oxidation Stage in the Chemistry of Protoberberine Alkaloids. Synthesis of 7,8-Dehydroberberines. *Tetrahedron* **1991**, *47*, 5841–5846.

(51) Grycová, L.; Dostál, J.; Marek, R. Quaternary Protoberberine Alkaloids. *Phytochemistry* **2007**, *68*, 150–175.