Nickel-catalysed Substitution Reactions of Allylic Compounds with Soft Nucleophiles: an Efficient Alternative to Palladium Catalysis

Hervé Bricout, Jean-François Carpentier and André Mortreux*

Laboratoire de Catalyse Hétérogène et Homogène, Groupe de Chimie Organique Appliquée, URA CNRS 402, ENSCL, BP 108—59652 Villeneuve d'Ascq Cedex, France

Substitution reactions of allyl alcohol derivatives **1a**-c with diethylamine, phenol and dimethyl malonate are efficiently carried out in the presence of Ni(dppb)₂ (dppb = 1,2-diphenylphosphinobutane) as catalyst and ammonium salts or bases as promoters or co-reagents.

Transition metal catalysed substitution reactions of allylic alcohol derivatives with nucleophilic reagents is an important carbon–carbon bond forming process. Most of the work in this field has been devoted to palladium complexes for the design of chemo-, regio-, stereo- and enantio-selective catalyst systems.¹ Nickel catalysts have also, to a lesser extent, attracted attention and have almost essentially been used with hard nucleophiles such as organometallic reagents.² Soft nucleophiles require much more drastic reaction conditions and afford quite poor results compared to palladium catalysts.³ Here we describe some nickel catalyst systems that enable the reaction of allylic alcohol derivatives with soft nucleophiles to proceed under mild conditions with high turnover frequencies.

The substitution reactions were first examined using allyl acetate **1a** or allyl phenyl ether **1b** and 2–3 equiv. of diethylamine **2** in the presence of 0.5 mol% of a nickel catalyst (Scheme 1). The results are summarised in Table 1. When the reaction was carried out with **1a** at 80 °C in THF in the presence of Ni(dppe)₂ (dppe = 1,2-diphenylphosphinoethane) or Ni(dppp)₂ (dppp = 1,2-diphenylphosphinopropane) (prepared *in situ* by mixing Ni(COD)₂ and 2 equiv. of phosphine ligand), the expected product **4** was obtained in a quantitative yield after 4.5 and 2.75 h respectively (entries 1 and 2). The use of Ni(dppb)₂ resulted in a large increase in the catalytic activity^{3d} and the same reaction proceeded under much milder conditions,

R ¹ 0 +	NuH [NiL ₂]	Nu	+ R ¹ OH or + CO ₂ + MeOH
1a R ¹ = Ac b R ¹ = Ph c R ¹ = MeO ₂ C	2 Nu = NEt ₂ 3 Nu = PhO	4 Nu = NEt ₂ 5 Nu = PhO	

Scheme 1

i.e. 50 °C within 2 h (entry 3). The catalytic activity of Ni(dppb)₂ was also greatly affected by the solvent nature. Namely, under comparable conditions, the time to complete the reaction was slightly increased by using toluene as the solvent (entry 4). Conversely, more interesting results were observed in the presence of DMF and acetonitrile (entries 5 and 6), despite the partial solubility of Ni(dppb)2 in acetonitrile. Owing to the apparent relation between the catalytic activity and polarity of the solvent and to the fact that all the Ni(dppb)₂-catalysed reactions exhibited increasing rates as far as the reaction proceeded,† the effect of added ammonium salts was investigated. The introduction of diethylammonium acetate (the salt normally produced by neutralisation of the acetic acid formed by excess diethylamine) in the early stages of the reaction between allyl acetate 1a and diethylamine 2 allowed the reaction time to be reduced (entry 7). Somewhat better results were obtained by using diethylammonium phenoxide (entry 8) or tetrabutylammonium hexafluorophosphate (entry 9).‡ In the latter case, the optimised time to perform the reaction was reduced from 90 to 14 min by using a minute amount of [NBu₄][PF₆] (2.5 mol% with respect to the allylic substrate) (compare entries 3 and 9). A further example (reaction of allyl phenyl ether 1b with diethylamine 2 shows a similar salt effect and the expected product 4 was obtained selectively in toluene at 50 °C within 2 min (entries 10 and 11).

The catalytic activity of Ni(dppb)₂ was further investigated by using other combinations of allylic substrates 1a-c and the nucleophiles 3 and 6 (Table 1). Substitution of methyl allyl carbonate 1c with phenol 3 in toluene also afforded a quantitative yield of 5 but the reaction was quite slow even at 80 °C (entry 12). In this case, the use of a polar solvent such as acetonitrile almost totally inhibited the process; this may be

Table 1 Nickel-catalysed substitution reactions of allyl compounds with soft nucleophiles

		Reagents							
Entry	Catalyst ^a	Allyl 1	NuH	Allyl : NuH : Ni	Solvent	Additive (equiv. Ni)	T/°C	Product ^b	t ^e / min
1	Ni(dppe) ₂	а	2	200:600:1	THF		80	4	270
2	$Ni(dppp)_2$	а	2	200:600:1	THF		80	4	165
3	Ni(dppb) ₂	а	2	200:600:1	THF		50	4	90
4	Ni(dppb) ₂	а	2	200:600:1	PhMe		50	4	110
5	Ni(dppb) ₂	а	2	200:600:1	DMF		50	4	6
6	$Ni(dppb)_2^d$	а	2	200:600:1	MeCN	_	50	4	< 2
7	$Ni(dppb)_2$	a	2	200:600:1	THF	$[NH_2Et_2][AcO]$ (100)	50	4	36
8	Ni(dppb) ₂	а	2	200:600:1	THF	[NH ₂ Et ₂][PhO] (100)	50	4	9
9	Ni(dppb) ₂	а	2	200:600:1	THF	$[NBu_4][PF_6]$ (5)	50	4	14
10	Ni(dppb) ₂	b	2	200:400:1	PhMe		50	4	11
11	Ni(dppb) ₂	b	2	200:400:1	PhMe	[NH ₂ Et ₂][PhO] (100)	50	4	< 2
12	Ni(dppb) ₂	с	3	300:200:1	PhMe ^e	_	80	5	210
13	Ni(dppb) ₂	с	3	300:200:1	PhMe ^e	NEt ₃ (200)	80	5	60
14	Ni(dppb) ₂	с	6	250:100:1	THF	_	80	8 /	150
15	Ni(dppb) ₂	а	6	200:300:1	THF	BSA ^g (300)	25	7/8 ^h	< 1
16	Ni(dppb) ₂	a	6	250:100:1	THF	BSA ^g (250)	25	8	50
17	$Ni(dppb)_2$	b	6	250:100:1	THF	BSA ^g (250)	25	8	< 2

^{*a*} Prepared *in situ* from Ni(COD)₂ and 2 equiv. of the corresponding phosphine ligand. ^{*b*} No side products were observed. ^{*c*} Optimized time for total conversion of the reagents to the desired product as indicated by GLC. ^{*d*} Catalyst was partially soluble. ^{*e*} Reaction carried out in 5 ml of solvent. ^{*f*} A mixture of 7 and 8 was obtained after 30 min. reaction in 20 and 75% GLC yields, respectively. ^{*g*} N,O- bis(trimethylsilyl)acetamide. ^{*h*} Compounds 7 and 8 were obtained in 92 and 8% GLC yields, respectively.

1864



explained by the higher acidity of phenol in acetonitrile than in toluene which contributes to the desctruction of the active Ni^0 species, as is evident by the rapid discolouration of the reaction medium. Actually, catalytic results were improved by introducing an equimolar amount of triethylamine with respect to phenol (entry 13).

The reaction of allylic substrates 1a-c with dimethyl malonate 6 afforded the mono- and/or di-allylated products (7 and 8, respectively), (Scheme 2). The reaction proceeded rather slowly with methyl allyl carbonate 1c (entry 14), but the combination of allyl acetate 1a or phenoxide 1b with BSA [*N*,*O*,-bis(trimethylsilyl)acetamide]⁴ considerably increased the reaction rates, and turnover frequencies higher than 12 000 mol (7/8).mol(Ni)⁻¹.h⁻¹ were observed at room temperature (entries 15–17). Thus, the synthesis of 7 was carried out in 92% yield from allyl acetate 1a, whereas the selective formation of the diallyl compound 8 was best performed from allyl phenoxide 1b.

The originality of this efficient nickel-based catalyst system compared to previous ones reported arises from (i) the use of a preformed zerovalent nickel complex, and (ii) the judicious choice of the polarity of the reaction medium.

Received, 22nd June 1995; Com. 5/04031K

J. CHEM. SOC., CHEM. COMMUN., 1995

Footnotes

 \dagger Exponential plots for the conversion to 4 vs. reaction time were observed for all Ni(dppb)₂-catalysed runs, whereas those conducted in the presence of dppe and dppp exhibited classical sigma profiles.

[‡] In a typical experiment, to Ni(COD)₂ (36 mg, 0.13 mmol) was added a solution of dppb (111 mg, 0.26 mmol) and [NBu₄][PF₆] (252 mg, 0.65 mmol) in THF (12.5 ml) under nitrogen. After stirring for 15 min, allyl acetate **1a** (2.60 g, 26 mmol), diethylamine **2** (5.70 g, 78 mmol) and heptane (1.00 g, 10 mmol) as an internal standard were added. The solution was stirred at 50 °C and the reaction was complete, compound **4** was isolated from the reaction mixture by distillation and fully characterised by NMR and MS techniques.

References

- For leading references, see: B. M. Trost and T. F. Verhoeven, in Comprehensive Organometallic Chemistry, ed. G. Wilkinson, Pergamon, Oxford, UK, 1982; R. F. Heck, Palladium Reagents in Organic Synthesis, Academic, New York, 1985; J. Tsuji, Tetrahedron, 1986, 42, 4361; J. P. Collman, L. S. Hegedus, J. R. Norton and R. G. Finke, Principles and Applications of Organotransition Metal Chemistry, University Science Books, Mill Valley, CA, 1987; B. M. Trost, Angew. Chem., Int. Ed. Engl., 1989, 28, 1173; G. Consiglio and R. Waymouth, Chem. Rev., 1989, 89, 257; C. G. Frost, J. Howarth and J. M. J. Williams, Tetrahedron: Asymmetry, 1992, 3, 1089.
- 2 For leading and recent references, see: C. Chuit, H. Felkin, C. Frajerman, G. Roussi and G. Swierczewski, J. Chem. Soc., Chem. Commun., 1968, 1604; M. Cherest, H. Felkin and J. O. Umpleby, J. Chem. Soc., Chem. Commun., 1981, 681; G. Consiglio, F. Morandini and O. Piccolo, J. Am. Chem. Soc., 1981, 103, 1846; G. Consiglio, O. Piccolo, L. Roncetti and F. Morandini, Tetrahedron, 1986, 42, 2043; Y. Kobayashi and E. Ikeda, J. Chem. Soc., Chem. Commun., 1994, 1789; S. K. Kang, D. G. Cho, C. H. Park, E. Y. Namkoong and J. S. Shin, Synth. Commun., 1995, 25, 1659.
- (a) J. Furukawa, J. Kiji, K. Yamamoto and T. Tojo, *Tetrahedron*, 1973,
 29, 3149; (b) T. Yamamoto, J. Ishizu and A. Yamamoto, J. Am. Chem.
 Soc., 1981, 103, 6863; (c) T. Cuvigny and M. Julia, J. Organomet. Chem.,
 1983, 250, C21; (d) T. Cuvigny and M. Julia, J. Organomet. Chem., 1986,
 317, 383; (e) E. Alvarez, T. Cuvigny and M. Julia, J. Organomet. Chem.,
 1988, 339, 199.
- 4 B. M. Trost and D. J. Murphy, Organometallics, 1985, 4, 1143.