

Communication

An organometallic complex formed by an unconventional radical SAM enzyme

Min Dong, Masaki Horitani, Boris Dzikovski, Maria-Eirini Pandelia,
Carsten Krebs, Jack H. Freed, Brian M. Hoffman, and Hening Lin

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.6b04155 • Publication Date (Web): 28 Jul 2016

Downloaded from <http://pubs.acs.org> on July 31, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



An organometallic complex formed by an unconventional radical SAM enzyme

Min Dong,^{1,#} Masaki Horitani,^{2,#} Boris Dzikovski,¹ Maria-Eirini Pandelia³, Carsten Krebs⁴, Jack H. Freed,¹ Brian M. Hoffman^{2,*} and Hening Lin^{1,5*}


¹Howard Hughes Medical Institute, Department of Chemistry and Chemical Biology, Cornell University, Ithaca, New York, 14853. ²Department of Chemistry, Northwestern University, Evanston, Illinois 60208. ³Department of Biochemistry, Brandeis University, MA 02453. ⁴Department of Chemistry and Department of Biochemistry and Molecular Biology, The Pennsylvania State University, University Park, Pennsylvania 16802. ⁵Howard Hughes Medical Institute, Cornell University, Ithaca, New York, 14853.

Supporting Information Placeholder

ABSTRACT: *Pyrococcus horikoshii* Dph2 (*PhDph2*) is an unusual radical SAM enzyme involved in the first step of diphthamide biosynthesis. It catalyzes the reaction by cleaving S-adenosylmethionine (SAM) to generate a 3-amino-3-carboxy propyl (ACP) radical. To probe the reaction mechanism, we synthesized a SAM analog (SAM_{CA}), in which the ACP group of SAM is replaced with a 3-carboxyallyl group. SAM_{CA} is cleaved by *PhDph2*, yielding a paramagnetic ($S = 1/2$) species, which is assigned to a complex formed between the reaction product, α -sulfinyl-3-butenic acid, and the [4Fe-4S] cluster. Electron-nuclear double resonance (ENDOR) measurements with ¹³C and ²H isotopically labeled SAM_{CA} support a π -complex between the C=C double bond of α -sulfinyl-3-butenic acid, and the unique iron of the [4Fe-4S] cluster. This is the first example of a radical SAM related [4Fe-4S]⁺ cluster forming an organometallic complex with an alkene, shedding additional light into the mechanism of *PhDph2* and expanding our current notions for the reactivity of [4Fe-4S] clusters in radical SAM enzymes.


Diphthamide is a post-translationally modified histidine residue on archaeal and eukaryotic translation elongation factor 2 (EF2), a protein essential for ribosomal protein synthesis. Its biosynthesis involves at least seven proteins.^{1,2} The first step involves the transfer of 3-amino-3-carboxypropyl (ACP) group derived from S-adenosyl-L-methionine (SAM) to the histidine residue of EF2. *In vitro*, the [4Fe-4S]-containing radical SAM (RS) enzyme *PhDph2*³ or the eukaryotic Dph1-Dph2 homodimer⁴ is sufficient for this step, with dithionite as the reducing agent. In contrast to classical RS enzymes that cleave the C_{5'}Ade-S bond of SAM to form 5'-deoxyadenosyl radical (5'-dA•) (Scheme 1A)⁵, *PhDph2* cleaves the C_{γ, Met}-S bond to form an ACP radical (Scheme 1B).³

Scheme 1. *PhDph2* is different from classical RS enzymes.



The formation of the ACP radical is supported by the *PhDph2*-catalyzed generation of 2-aminobutyric acid and homocysteine sulfinic acid in the absence of the substrate protein, *PhEF2*.³ However, the ACP radical has not been directly observed even when we carried out freeze quench-electron paramagnetic resonance (EPR) experiments (Figure S1). This is not surprising because the ACP radical is expected to be short-lived. Following the strategy developed by Frey and coworkers to generate a more stable allylic analog of the 5'-dA•,^{6,7} we synthesized a SAM analog, SAM_{CA}, in which the ACP group of SAM was replaced with a 3-carboxyallyl group (Scheme 2). If *PhDph2* could accept SAM_{CA} as a substrate, a 3-carboxyallyl radical would be generated, which should be more stable and allow direct observation by EPR. As shown below, even with SAM_{CA}, we could not detect the radical intermediate. However, surprisingly, we found that a dithionite quenched radical product formed an organometallic complex with the Fe-S cluster, revealing an interesting reactivity of the [4Fe-4S] cluster in RS enzymes.

Scheme 2. Structures of SAM_{CA} and aza-SAM_{CA}.



We initially examined whether SAM_{CA} could serve as a substrate of *PhDph2*. High-performance liquid chromatography (HPLC) showed that all the SAM_{CA} was converted to 5'-deoxy-5'-methylthioadenosine (MTA) within 20 min, whereas in the absence of dithionite or *PhDph2*, no SAM_{CA} was cleaved (Figure 1A). These results demonstrate that *PhDph2* is able to catalyze the cleavage of the 'correct' C-S bond of SAM_{CA}. In the presence of the substrate protein *PhEF2*, the

same reaction product was detected (Figure S2). However, PhEF2 was not modified by SAM_{CA}. This was likely because SAM_{CA} generated a more stable radical (see more experimental evidence for the stability below), which was not active enough to react with PhEF2.




Figure 1. A). HPLC traces of the SAM_{CA} cleavage reaction (reaction time 20 min) catalyzed by PhDph2 at room temperature. B). X-band CW EPR spectra of PhDph2 in the absence and presence of SAM_{CA}. A spectral simulation for the $S = 1/2$ species is shown as green dotted line. $T = 12$ K.

We then prepared freeze quench samples and used EPR to detect radical signals. The dithionite-reduced PhDph2 in the absence of SAM_{CA} showed a typical $[4Fe-4S]^{1+}$ EPR signal at 12 K (Figure 1B),³ which was temperature-dependent and broadened beyond detection at 30 K (Figure S1). When PhDph2 in the presence of dithionite was allowed to react with SAM_{CA} for 30 s, the $[4Fe-4S]^{1+}$ signal was replaced by a nearly axial signal, with a split feature at g_1 (Figure 1B). The ratio of this g_1 splitting at X- and Q-band scaled with the ratio of microwave frequencies (Figure S3), showing that the splitting represents two different conformations and not a hyperfine interaction with a paramagnetic nucleus. Simulation of the spectra gave for the one conformer principal g -values $g = [2.092, 2.010, 1.990]$, and for the second one $g = [2.074, 2.010, 1.990]$. Both signals exhibited similar relaxation properties and were hardly detectable above 70 K (Figure S1). The large g -anisotropy and the absence of proton hyperfine splittings suggest that it is not a free allyl radical.^{6,8} In contrast, the signals of the new species formed in the reaction of the reduced PhDph2 with SAM_{CA} exhibited an “isotropic” g -value, $g_{iso} = (g_{11} + g_{22} + g_{33})/3 > g_e = 2.0023$,⁹ which is reminiscent of that observed in high potential iron-sulfur proteins (HiPIP) and $[4Fe-4S]^{3+}$ intermediates trapped with the $[4Fe-4S]$ proteins IspG¹⁰ and IspH¹¹. With ⁵⁷Fe-enriched PhDph2, the EPR signal became broadened (Figure 1B), further supporting that the new species (termed CA) is associated with the $[4Fe-4S]$ cluster.

When we monitored the reaction at different time points, the EPR signal of CA persisted for >30 min, even after all SAM_{CA} was consumed (Figure S4). This suggested that CA was a complex of the reaction product with the $[4Fe-4S]^{1+}$ cluster, the latter regenerated by dithionite reduction of the $[4Fe-4S]^{2+}$ produced in the reaction. To test this hypothesis, protein-free aliquots of the reaction products were added to ‘fresh’ dithionite-reduced PhDph2. The same anisotropic EPR

signal as that of CA was obtained, supporting that CA is a product complex (Figure S5).

To understand the structure of CA, we analyzed the reaction products using ¹H-NMR. Several new peaks were observed from the reaction containing PhDph2, SAM_{CA}, and dithionite (Figure S6), which were absent in control reactions without dithionite. Some of the new peaks were assigned to MTA. Crotonic acid and its isomer, 2-butenic acid, which are the hydrogen abstraction products of the carboxyallyl radical, were not formed by comparison to the spectra of their standards. Three new peaks (a-c) were assigned to γ -sulfinylcrotonic acid (Figure S6) and the remaining three peaks (d, e and f) around 5.1 and 5.8 ppm were assigned to α -sulfinyl-3-butenic acid (Figure S6); the remaining proton on α -sulfinyl-3-butenic acid is localized at 3.4 ppm based on ¹H-¹H COSY (correlation spectroscopy) (Figure S14, S15). Using 1,1-²H₂-SAM_{CA} as the substrate, peaks c, e and f (Figure S7) disappeared, which supported the assignments. The assignments were confirmed by ¹H/¹³C HMBC (Heteronuclear Multiple Bond Correlation), HSQC (Heteronuclear Single Quantum Correlation) and ¹H-¹H COSY (Figure S8 - S15). The reaction products were also chemically modified to allow detection by LC-MS (Figure S16). Our data thus demonstrated the formation of γ -sulfinylcrotonic and α -sulfinyl-3-butenic acid, which likely resulted from the recombination of the carboxyallyl radical with the dithionite-derived SO₂[•] radical, similar to previously reported reactions of radicals with dithionite^{3,12,13}. The NMR signal intensities suggested roughly equal formation of the two products.

We then performed 35 GHz CW/pulse ¹³C, ¹H and ²H ENDOR with 1-¹³C, 2-¹³C, 3-¹³C, 1,1-²H₂ and 1,1,2,3-²H₄ labeled SAM_{CA} (see Scheme 2). Field-modulated 35 GHz CW ENDOR spectra obtained from reactions with 1-¹³C, 2-¹³C labeled SAM_{CA} both exhibit ¹³C doublets with broad component lines, split by the ¹³C hyperfine coupling, $A \sim 5$ -8 MHz (Figure 2). An estimate of the magnitude of the hyperfine coupling tensors for these two ¹³C sites of SAM_{CA} (Table S1) was obtained through analysis of 2D field-frequency patterns of ENDOR spectra collected across the EPR envelopes of the SAM_{CA} isotopologs (Figures S17, S18). The analysis showed comparable isotropic hyperfine tensors, $A_{iso} \sim 7$ MHz for both 1-¹³C and 2-¹³C. For 3-¹³C labeled SAM_{CA}, a considerably smaller isotropic coupling, $A_{iso} \sim 0.6$ MHz, was estimated using Mims pulsed ENDOR (Figure 2A, Inset).




Figure 2. (A) 35 GHz CW ¹³C ENDOR spectra for CA with 1-¹³C and 2-¹³C-labeled SAM_{CA} collected at g_2 ; the doublets are split by the hyperfine coupling, and centered at the Larmor

frequency as described in SI. **Inset:** Mims ENDOR spectra of $3\text{-}^{13}\text{C}$ SAM_{CA}. **(B)** ^1H Davies ENDOR difference spectra of CA generated by subtracting from the spectrum of ^1H -SAM_{CA} (natural abundance) the spectra for $1,1,2,3\text{-}^2\text{H}$ -labeled SAM_{CA} (upper) and $1,1\text{-}^2\text{H}$ -labeled SAM_{CA} (lower). Spectra were collected at g_2 and normalized by EPR echo height.

To further probe the structure of CA, ^1H and ^2H pulse ENDOR measurements were performed with $1,1\text{-}^2\text{H}_2$, and $1,1,2,3\text{-}^2\text{H}_4$ labeled SAM_{CA}. Only the pulsed ENDOR techniques resolved individual features, but the lower sensitivity of pulsed vs CW ENDOR allowed spectra to be collected only at the intensity maximum of the EPR envelope, near g_2 . The ^1H signals from the labeled sites were obtained by subtracting the spectra obtained using the natural-abundance SAM_{CA} (Figure 2B). After subtraction of the $1,1\text{-}^2\text{H}_2$ spectrum, a doublet of ^1H peaks assignable to the $1,1\text{-}^1\text{H}$ nuclei was obtained with coupling constant of $A \sim 7$ MHz; subtraction with the $1,1,2,3\text{-}^2\text{H}_4$ labeled SAM_{CA} showed an enhanced intensity for this doublet (Figure 2B), indicating that the $1,1\text{-}^1\text{H}$ and $2\text{-}^1\text{H}$ nuclei had comparable couplings (assigned based on $2\text{-}^{13}\text{C}$ data). In addition, weak broad features corresponding to a doublet with $A \sim 2$ MHz were assigned to $3\text{-}^1\text{H}$. Mims ^2H ENDOR spectra (Figure S19 and S20) provided support for this assignment. The substantial isotropic couplings to $1\text{-}^{13}\text{C}$ and $2\text{-}^{13}\text{C}$ ($A_{\text{iso}} \sim 7$ MHz), the deviations of the g -values from $g \sim 2$, and the ^{57}Fe broadening imply the spin is primarily localized on the cluster, but with close C-Fe interaction.

The similarity of $1\text{-}^{13}\text{C}$ and $2\text{-}^{13}\text{C}$ hyperfine couplings, as well as the similarity of the $1,1\text{-}^2\text{H}$ and the $2\text{-}^2\text{H}$ couplings, suggests that both 1-C and 2-C have similar direct interactions with the unique Fe of the $[\text{4Fe-4S}]$ cluster. In contrast, the weaker $3\text{-}^{13}\text{C}$ and $3\text{-}^2\text{H}$ hyperfine couplings show that 3-C does not experience a significant direct interaction with the Fe. These results suggest that CA only consists of one of the two products, α -sulfinyl-3-butenic acid, coordinated to the unique Fe site of the $[\text{4Fe-4S}]$ cluster. If CA also involved a complex with γ -sulfinylcrotonic acid, 2-C and 3-C should have similar strong hyperfine couplings, and 1-C should have weaker coupling, contrary to our observations (Figure 2A). We further confirmed this by utilizing the instability of α -sulfinyl-3-butenic acid. After incubating the protein-free aliquots of the reaction products at room temperature overnight, the peaks corresponding to α -sulfinyl-3-butenic acid disappeared in the ^1H -NMR spectrum (Figure S21), while the signals from γ -sulfinylcrotonic acid remained. The instability of α -sulfinyl-3-butenic acid is likely due to the acidity of the α -proton, which is adjacent to two strong electron-withdrawing groups.¹⁴ When the mixture containing only the γ -sulfinylcrotonic acid product was added to 'fresh' dithionite-reduced *PhDph2*, the signal associated with CA was not detected by EPR (Figure S22). This result supports the notion that γ -sulfinylcrotonic acid does not form an EPR-active complex with the $[\text{4Fe-4S}]$ cluster and that CA is best described as a complex formed between α -sulfinyl-3-butenic acid and the $[\text{4Fe-4S}]$ cluster.

We then probed whether the sulfinyl and carboxyl of α -sulfinyl-3-butenic acid groups are required for CA formation. We synthesized a non-cleavable analog of SAM_{CA}, aza-SAM_{CA} (Scheme 2)¹⁵. The freeze quench-EPR experiment

with aza-SAM_{CA} showed only a $[\text{4Fe-4S}]^+$ signal (Figure S23) that was similar to that of the reduced cluster in *PhDph2* in the absence of SAM_{CA}. Therefore, CA is not formed by binding of uncleaved SAM_{CA} to the $[\text{4Fe-4S}]$ cluster. Addition of crotonic acid to *PhDph2* also failed to generate the CA species. Thus, the sulfinic group in α -sulfinyl-3-butenic acid is likely involved in the formation of CA. Similarly, adding allylsulfenic acid to *PhDph2* did not generate the CA signal either. Thus, the carboxyl group in α -sulfinyl-3-butenic acid is also necessary. The results suggest that both the sulfinic group and the carboxylate of α -sulfinyl-3-butenic acid bind to the Fe-S cluster, facilitating the formation of a π -complex between that Fe and the C=C double bond of the product (Figure 3).




Figure 3. Plausible structures of CA and formation mechanism.

CA can have two formal resonance structures (Figure 3, structures **a** and **b**). To further probe the oxidation state of CA, we attempted to reduce CA by low-temperature γ -irradiation (cryoreduction)¹⁶ (Figure S24). The EPR signal did not decrease upon γ -irradiation, as would be expected for a $[\text{4Fe-4S}]^{3+}$ cluster. Similarly, extra dithionite could not reduce CA, which is inconsistent with the expectation for a $[\text{4Fe-4S}]^{3+}$ cluster (Figure S25)¹⁷. In contrast, when we tried to oxidize CA with $\text{K}_3[\text{Fe}(\text{CN})_6]$, CA was oxidatively degraded to a $[\text{3Fe-4S}]^+$ cluster (Figure S25), an outcome commonly observed in similar treatments of $[\text{4Fe-4S}]^+$ clusters of RS enzymes with chemical oxidants. Thus, the Fe-S core in CA behaves like a $[\text{4Fe-4S}]^+$ cluster. We also sought to use ^{57}Fe Mössbauer spectroscopy. However, owing to the rather modest yield of CA (50% of total Fe) and the presence of at least one additional EPR-silent Fe-containing product, the analysis of the spectra does not permit definitive conclusions about the electronic structure of CA (Figures S26, 27 and 28).

With the characterization of reaction products and the structure of CA, we proposed a plausible mechanism for the formation of CA (Figure 3). The $[\text{4Fe-4S}]^+$ cluster in *PhDph2* provides one electron to SAM_{CA}, cleaving the C_{CA}-S bond and generating a 3-carboxyallyl radical, MTA, and a $[\text{4Fe-4S}]^{2+}$ cluster. The 3-carboxyallyl radical reacts with a dithionite-derived $\text{SO}_2^{\bullet-}$ radical, forming γ -sulfinylcrotonic acid and α -sulfinyl-3-butenic acid. The sulfinic group, the carboxyl group, and the C=C double bond of α -sulfinyl-3-butenic acid, coordinate the $[\text{4Fe-4S}]^+$ cluster (reduced by extra dithionite) to generate the π -complex. The coordination of the sulfinic and carboxyl group helps to place the double bond in a favorable position for π interactions with the unique Fe. The two conformers revealed by EPR (Figure 1B) could be attributed to different binding orientations of the α -sulfinyl-

3-butenic acid to $[4\text{Fe-4S}]^+$ cluster (the three chelating functional groups may exchange positions).

In summary, using SAM_{CA} , we have detected an EPR-active species, CA, formed in the enzymatic reaction of *PhDph2*. Various experiments demonstrate that CA is a complex formed between the $[4\text{Fe-4S}]^{2+}$ cluster and one of the reaction products, α -sulfinyl-3-butenic acid, coordinated to the unique cluster Fe through the sulfinyl and carboxylate oxygens, with the C=C double bond forming an organometallic π -complex with the Fe. Although we could not accumulate the radical intermediate that we set out to detect with SAM_{CA} , formation of both α -sulfinyl-3-butenic acid and γ -sulfinylcrotonic acids provides additional support for the radical mechanism of *PhDph2* reaction, as a nucleophilic mechanism cannot account for their formation. Interestingly, although the EPR g tensors of CA shows similarity to those of $[4\text{Fe-4S}]^{3+}$ -like clusters observed in two enzymes involved in the non-mevalonate pathway, IspG and IspH, the cluster in CA behaves like a $[4\text{Fe-4S}]^+$ cluster.

The present case can be described as the reaction of the unique Fe of the $[4\text{Fe-4S}]^+$ cluster with the alkene product to form a stable organometallic complex. Such an alkene complex has reported for the nitrogenase FeMo-cofactor and the ethylenic product of alkyne reduction¹⁸, and for mutants of IspH and its substrate HMBPP.^{11,19} However, similar organometallic complex had not been demonstrated for RS enzymes until very recently, when the $[4\text{Fe-4S}]^{2+}$ cluster of the RS enzyme, pyruvate formate-lyase activating enzyme is shown to react with the 5'-dA• to form a highly reactive organometallic intermediate.²⁰ This study and our current result suggest that the unique iron of $[4\text{Fe-4S}]$ clusters in RS enzymes can readily form a C-Fe complex. Such a property not only may help to explain the unusual chemistries catalyzed by the $[4\text{Fe-4S}]$ -containing RS enzymes, but also may open up new avenues to tune the activity or develop new chemistry for this enzyme class or for other $[4\text{Fe-4S}]$ -containing enzymes.

ASSOCIATED CONTENT

Supporting Information

Experimental materials and methods, supporting figures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Authors

Brian M. Hoffman (bmh@northwestern.edu) and Hening Lin (hl379@cornell.edu).

Author Contributions

[#]MD and MH contributed equally to this work.

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

This work is supported by NIH (GM088276, P41-GM103521, and GM111097). We thank Dr. Ivan Keresztes and Mr. Anthony Condo for assistance with NMR; Prof. H. Halpern at University of Chicago for access to the ⁶⁰Co Gammacell irradiator.

REFERENCES

- (1) Su, X.; Lin, Z.; Lin, H. *Crit. Rev. Biochem. Mol. Biol.* **2013**, *48*, 515.
- (2) Schaffrath, R.; Abdel-Fattah, W.; Klassen, R.; Stark, M. J. *Mol. Microbiol.* **2014**, *94*, 1213.
- (3) Zhang, Y.; Zhu, X.; Torelli, A. T.; Lee, M.; Dzikovski, B.; Koralewski, R. M.; Wang, E.; Freed, J.; Krebs, C.; Ealick, S. E.; Lin, H. *Nature* **2010**, *465*, 891.
- (4) Dong, M.; Su, X.; Dzikovski, B.; Dando, E. E.; Zhu, X.; Du, J.; Freed, J. H.; Lin, H. *J. Am. Chem. Soc.* **2014**, *136*, 1754.
- (5) Broderick, J. B.; Duffus, B. R.; Duschene, K. S.; Shepard, E. M. *Chem. Rev.* **2014**, *114*, 4229.
- (6) Magnusson, O. T.; Reed, G. H.; Frey, P. A. *J. Am. Chem. Soc.* **1999**, *121*, 9764.
- (7) Horitani, M.; Byer, A. S.; Shisler, K. A.; Chandra, T.; Broderick, J. B.; Hoffman, B. M. *J. Am. Chem. Soc.* **2015**, *137*, 7111.
- (8) McManus, H. J.; Fessenden, R. W.; Chipman, D. M. *J. Phys. Chem.* **1988**, *92*, 3778.
- (9) Belinskii, M. *Chem. Phys.* **1993**, *172*, 189.
- (10) Wang, W.; Li, J.; Wang, K.; Huang, C.; Zhang, Y.; Oldfield, E. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 11189.
- (11) Wang, W.; Wang, K.; Span, I.; Jauch, J.; Bacher, A.; Groll, M.; Oldfield, E. *J. Am. Chem. Soc.* **2012**, *134*, 11225.
- (12) Ko, Y.; Ruzsyczky, M. W.; Choi, S. H.; Liu, H. W. *Angew. Chem. Int. Ed. Engl.* **2015**, *54*, 860.
- (13) Chandor-Proust, A.; Berteau, O.; Douki, T.; Gasparutto, D.; Olagnier-de-Choudens, S.; Fontecave, M.; Atta, M. *J. Biol. Chem.* **2008**, *283*, 36361.
- (14) Khatik, G. L.; Kumar, R.; Chakraborti, A. K. *Org. Lett.* **2006**, *8*, 2433.
- (15) Joce, C.; Caryl, J.; Stockley, P. G.; Warriner, S.; Nelson, A. *Org. Biomol. Chem.* **2009**, *7*, 635.
- (16) Davydov, R.; Hoffman, B. M. *Arch. Biochem. Biophys.* **2011**, *507*, 36.
- (17) de la Torre, A.; Lara, C.; Yee, B. C.; Malkin, R.; Buchanan, B. B. *Arch. Biochem. Biophys.* **1982**, *213*, 545.
- (18) Lee, H. I.; Igarashi, R. Y.; Laryukhin, M.; Doan, P. E.; Dos Santos, P. C.; Dean, D. R.; Seefeldt, L. C.; Hoffman, B. M. *J. Am. Chem. Soc.* **2004**, *126*, 9563.
- (19) Wang, W.; Wang, K.; Liu, Y. L.; No, J. H.; Li, J.; Nilges, M. J.; Oldfield, E. *Proc. Natl. Acad. Sci. U S A.* **2010**, *107*, 4522.
- (20) Horitani, M.; Shisler, K.; Broderick, W. E.; Hutcheson, R. U.; Duschene, K. S.; Marts, A. R.; Hoffman, B. M.; Broderick, J. B. *Science* **2016**, *352*, 822.

Table of content graphic

