

Design and synthesis of 14 and 15-membered macrocyclic scaffolds exhibiting inhibitory activities of hypoxia-inducible factor 1 α

Takahiro Shimizu, Norihito Takahashi, Vincent J. Huber, Yasunobu Asawa, Hiroki Ueda, Atsushi Yoshimori, Yukiko Muramatsu, Hiroyuki Seimiya, Hiroyuki Kouji, Hiroyuki Nakamura, Hiroki Oguri

PII: S0968-0896(20)30779-3
DOI: <https://doi.org/10.1016/j.bmc.2020.115949>
Reference: BMC 115949

To appear in: *Bioorganic & Medicinal Chemistry*

Received Date: 7 November 2020
Revised Date: 9 December 2020
Accepted Date: 9 December 2020

Please cite this article as: T. Shimizu, N. Takahashi, V.J. Huber, Y. Asawa, H. Ueda, A. Yoshimori, Y. Muramatsu, H. Seimiya, H. Kouji, H. Nakamura, H. Oguri, Design and synthesis of 14 and 15-membered macrocyclic scaffolds exhibiting inhibitory activities of hypoxia-inducible factor 1 α , *Bioorganic & Medicinal Chemistry* (2020), doi: <https://doi.org/10.1016/j.bmc.2020.115949>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



