÷

International Edition: DOI: 10.1002/anie.201603860 German Edition: DOI: 10.1002/ange.201603860

# Highly Selective Phosphinylphosphination of Alkenes with Tetraphenyldiphosphine Monoxide

Yuki Sato, Shin-ichi Kawaguchi,\* Akihiro Nomoto, and Akiya Ogawa\*

**Abstract:** In sharp contrast to tetraphenyldiphosphine, which does not add to carbon–carbon double bonds efficiently, its monoxide,  $[Ph_2P(O)PPh_2]$  can engage in a radical addition to various alkenes, thus affording the corresponding 1-phosphinyl-2-phosphinoalkanes regioselectively, and they can be converted into their sulfides by treatment with elemental sulfur. The phosphinylphosphination proceeds by the homolytic cleavage of the  $P^V(O)-P^{III}$  single bond of  $Ph_2P(O)PPh_2$ , followed by selective attack of the phosphinyl radical at the terminal position of the alkenes, and selective trapping of the resulting carbon radical by the phosphino group. Furthermore, the phosphinylphosphination product could be converted directly into its platinum complex with a hemilabile P,O chelation.

Organophosphorus compounds play a vital role in organic synthesis, catalysis, materials chemistry, medicinal chemistry, and coordination chemistry.<sup>[1]</sup> Therefore, the development of new synthetic methods to create organophosphorus species is of great importance. For the selective introduction of phosphorus moieties onto organic molecules, the addition reaction to carbon-carbon multiple bonds is one of the most useful and atom-economical methods.<sup>[2]</sup> In particular, the 1,2addition of phosphorus compounds, bearing a P-P single bond, to carbon-carbon multiple bonds is the most straightforward method for the preparation of vicinal bis(phosphine)s,<sup>[3]</sup> which are useful bidentate ligands in transition metal catalyzed reactions.<sup>[4]</sup> Although several attractive examples of the addition of phosphorus compounds to alkynes have been reported,<sup>[5]</sup> to the best of our knowledge, the corresponding addition to alkenes has not been achieved. with the exception of the addition of specified diphosphines, such as tetramethyldiphosphine [(Me<sub>2</sub>P)<sub>2</sub>],<sup>[6]</sup> tetrachlorodiphosphine  $[(Cl_2P)_2]$ ,<sup>[7]</sup> tetrafluorodiphosphine  $[(F_2P)_2]$ ,<sup>[8]</sup> and 1,1-diaminodiphosphine,<sup>[9]</sup> to only very a few alkenes.

We initiated our study by investigating the addition of tetraphenyldiphosphine (1) to 1-dodecene (2a) in either the

```
[*] Y. Sato, Dr. A. Nomoto, Prof. Dr. A. Ogawa
Department of Applied Chemistry, Graduate School of Engineering
Osaka Prefecture University
1-1 Gakuen-cho, Nakaku, Sakai, Osaka 599-8531 (Japan)
E-mail: ogawa@chem.osakafu-u.ac.jp
Dr. S-i. Kawaguchi
Center for Education and Research in Agricultural Innovation
Faculty of Agriculture, Saga University
152-1 Shonan-cho Karatsu, Saga 847-0021 (Japan)
E-mail: skawa@cc.saga-u.ac.jp
Supporting information and the ORCID identification number(s) for
the author(s) of this article can be found under http://dx.doi.org/10.
1002/anie.201603860.
```

Angew. Chem. Int. Ed. 2016, 55, 1-5

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

#### Wiley Online Library

These are not the final page numbers!

presence of a radical initiator [AIBN or V-40 (1,1'-azobis-(cyclohexane-1-carbonitrile)] or under photoirradiation (Scheme 1 a). The addition reaction did not proceed at all,

a) Radical addition of PIII-PIII linkage to 1-dodecene



b) Radical addition of Y-Y and Z-Z compounds to C-C multiple bonds

$$\begin{array}{c|c} R & \overbrace{reactivity toward} & Y & R & \overbrace{capturing ability of} & Z \\ \hline C-C multiple bonds & \hline Y & Y & R \\ \hline Y & Y & Z & Y & R \\ \hline C-C multiple bonds & \hline Y & Y & Z \\ \hline Y & Y & Z & Y & Y \\ \hline C-C multiple bonds & \hline Y & Y & Z \\ \hline Y & Y & Z & Z \\ \hline \end{array}$$

c) This work : Radical addition of PV(O)-PIII linkage to alkenes



**Scheme 1.** Radical addition of the diphosphine 1 (P'''-P''') and diphosphine monoxide 3 [P''(O)-P'''] to alkenes.

and **1** was recovered quantitatively. We previously reported that the regioselective thioselenation of alkenes proceeds smoothly using a disulfide–diselenide mixed system,<sup>[10]</sup> although the addition to alkenes did not proceed efficiently using either a disulfide or diselenide species alone. In such heteroatom-mixed systems, two different heteroatom-centered radicals (Y, Z) can be generated and two heteroatom-containing compounds (Y-Y, Z-Z) with different radical capturing abilities exist. The more reactive radical attacks carbon–carbon multiple bonds and the generated carbon radical is captured by the heteroatom-containing compound with the better capturing ability (Scheme 1b).

With this information in mind, we selected tetraphenyldiphosphine monoxide (**3**), bearing a  $P^{V}(O)-P^{III}$  single bond, as an initial phosphorus compound for the radical addition to alkenes. This choice was because the homolytic cleavage of the  $P^{V}(O)-P^{III}$  bond can afford two different phosphorus-centered radicals and **3** has pentavalent and trivalent phosphorus atoms, which might have different radical capturing abilities. The 1,2-addition of **3** to alkenes proceeded regioselectively to afford the 1-phosphinyl-2-phosphinoalkanes **4** (Scheme 1 c). Recently, organophosphines with vicinal soft (P<sup>III</sup>) and hard [P<sup>V</sup>(O)] Lewis-base centers<sup>[11]</sup> were used as hemilabile ligands in transition metal catalyzed reactions.<sup>[12]</sup> Furthermore, the reduction of **4** readily affords bidentate bis(phosphine) ligands. Therefore, we set out to investigate

the novel phosphinylphosphination of alkenes with diphosphine monoxides.

When a mixture of **3** (0.3 mmol), **2a** (0.3 mmol), and V-40 (0.03 mmol, 10 mol%) was dissolved in benzene and refluxed for 20 hours, the complete consumption of **2a** and the formation of the phosphinylphosphination product **4a**, which has a diphenylphosphinyl group at the terminal carbon atom, were confirmed by <sup>1</sup>H and <sup>31</sup>P NMR spectroscopy (Scheme 2). The regioisomer **5a** was not detected, and



Scheme 2. Phosphinylphosphination of 2 a with 3.

**4a** was converted into the bis(phosphino)alkane 1-*P*-oxide 2-*P*-sulfide (BPOS) **6a** upon treatment with elemental sulfur.

Since a small amount of the self-polymerization product of **2a** was observed, the ratio of **3** to **2a**, initiator, and temperature were screened (Table 1) to identify the optimal reaction conditions. When the reaction was conducted at low temperatures, such as 30 and 60 °C, the self-polymerization of **2a** preferentially occurred (entries 2 and 3). In contrast, at high temperature (80 °C) and with the use of excess **3** suppressed the polymerization effectively and thus improved the yield of **6a** (entry 4). The phosphinylphosphination did not take place without a radical initiator (entry 5). Photoirradiation with a xenon lamp (500 W) through a sealed Pyrex

**Table 1:** Optimization of the phosphinylphosphination of alkene  $\mathbf{2a}$  with  $\mathbf{3}^{[a]}$ 

( Ph <sub>2</sub> F	D →−PPh <sub>2</sub> +	initiator (10 mol%) benzene <i>T</i> , 20 h	S <sub>8</sub> (3 equiv) 60 °C, 5 h	Ph <sub>2</sub> P ,PPh <sub>2</sub> S 6a
Entry	<b>3</b> (mmol)	Initiator	T [°C]	Yield [%] <sup>[b]</sup>
1	0.30	V-40	80	50
2	0.30	AIBN	60	23
3 <sup>[c]</sup>	0.30	V-70	30	14
4	0.45	V-40	80	79 (72)
5	0.45	_	80	0
6 <sup>[c,d]</sup>	0.90	xenon lamp	20	78 (72)

[a] Reaction conditions: **2a** (0.3 mmol), benzene (0.6 mL). [b] Determined by <sup>1</sup>H NMR spectroscopy. Isolated yields are shown within parentheses. [c] The substrates were dissolved in CH<sub>2</sub>Cl<sub>2</sub>. [d] The mixture was irradiated with a xenon lamp (500 W) through a sealed Pyrex NMR tube for 55 h at room temperature. V-40 = 1,1'-azobis(cyclohexane-1-carbonitrile), V-70 = 2,2'-azobis(4-methoxy-2,4-dimethylvaleronitrile), AIBN = 2,2'-azobis (isobutyronitrile).

NMR tube afforded **6a** in a good yield, although a prolonged reaction time was required (entry 6).

The phosphinylphosphination of various alkenes with **3** was conducted in the presence of either a catalytic amount of V-40 (Conditions A) or under photoirradiation (Conditions B; Table 2). The vinyl ether **2b** reacted with **3** efficiently. After the mixture was treated with  $S_8$  (0.9 mmol) at 60 °C for 5 hours, the BPOS **6b** was obtained in a good yield. Allyl ethers afforded the corresponding 1,2-adducts **6c-h** in excellent yields under photoirradiation. The reaction was tolerant of various functional groups such as fluoro (**2e**), methoxy (**2f**), ester (**2g**), and nitrile (**2h**) groups. The use of the allyl ester **2i** also gave the corresponding 1,2-adduct **6i** in a good yield. Cyclohexane-containing species (**2j** and **2k**) gave the corresponding BPOSs in good yields. Styrene (**2l**) did not undergo the phosphinylphosphination at all, because

**Table 2:** Phosphinylphosphination of several alkenes **2** with diphosphine monoxide  $\mathbf{3}^{[a]}$ 



[a] Reaction conditions A: **3** (0.45 mmol), **2** (0.3 mmol), V-40 (0.03 mmol), benzene (0.6 mL), 80 °C, and 20 h. After the reaction, the resulting mixture was treated with  $S_8$  (0.9 mmol) at 60 °C for 5 h. Reaction conditions B: **3** (0.6 mmol), **2** (0.2 mmol),  $CD_2CI_2$  (0.4 mL), xenon lamp (500 W), 40 h, room temperature, and subsequent treatment with  $S_8$ . [b] The mixture was irradiated for 55 h. [c] The mixture was irradiated for 20 h. [d] Phenylacetylene (0.3 mmol) was added.

#### www.angewandte.org

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

**K** These are not the final page numbers!

of preference for the self-polymerization. In contrast, the reaction of allylbenzene (**2m**) proceeded to afford BPOS **6m** in a moderate yield. Furthermore, the terminal alkyne **2n** also served as a viable substrate to give the 1,2-adduct **6n** in a good yield with excellent stereoselectivity.

The present reaction is a novel and straightforward method to synthesize 1-phosphinyl-2-phosphinoalkanes, which can be used as ligands for transition metal catalyzed reactions.<sup>[12]</sup> Therefore, we set out to create a **4d**/transition metal complex using the present method (Scheme 3). When a mixture of **3** (0.6 mmol) and **2d** (0.2 mmol) was photo-irradiated, the 1-phosphinyl-2-phosphinoalkane **4d** was obtained in a high yield. Subsequently,  $PtCl_2(PhCN)_2$  (0.2 mmol) was added to the resulting reaction mixture, and the platinum complex **7d** was successfully isolated in 66% yield.



Scheme 3. The facile preparation of platinum(II)/bis(phosphine) monoxide complex 7d.

To shed light on the reaction pathway, the diene **8** was mixed with **3** in the presence of a catalytic amount of V-40 (Scheme 4). As a result, the cyclized product **9** (anti/syn = 5/1)



Scheme 4. Phosphinylphosphination of diene 8 with 3.

was obtained and the noncyclized 1,2-adducts were not detected at all. This result strongly indicates that the phosphinylphosphination proceeds by a radical pathway. In addition, the rate of carbon radical capture by **3** was estimated to be  $5 \times 10^3 \text{ s}^{-1}\text{M}^{-1}$  at most, because the rate for the 5-*exo* cyclization of 5-hexenyl radical is  $4 \times 10^5 \text{ s}^{-1}$ .<sup>[13]</sup> The carbon radical capture rate of **3** is very slow considering the rate constant for the radical attack of diphenylphosphinyl radical on isoprene  $[k = (1.4 \pm 0.3) \times 10^7 \text{ s}^{-1}\text{M}^{-1}]^{[14]}$  and phenyl vinyl ether  $[k = (2.6 \pm 0.2) \times 10^6 \text{ s}^{-1}\text{M}^{-1}]^{[15]}$  Therefore, the key step of the phosphinylphosphination is the carbon radical capture by **3**.

To clarify the difference in reactivity between 1 and 3, we conducted the reaction between 2-bromopropane (0.2 mmol) and either 1 or 3 (0.6 mmol) with hexabutylditin (0.2 mmol) under photoirradiation (Scheme 5). In this system, an isopropyl radical was generated in situ by the abstraction of bromine from 2-bromopropane, and it was caused by the Sn–Sn bond cleavage of hexabutylditin upon near-UV irradiation. In sharp



Scheme 5. Photoinduced homolytic substitution of 1 and 3 with isopropyl radical.

contrast to **1**, which did not react with the generated isopropyl radical, **3** caused homolytic substitution at the trivalent phosphorus  $atom^{[16]}$  to generate isopropyl-(diphenyl)phosphine in 40% yield. This result strongly indicates that the  $\pi$ -type alkyl radical capturing ability of **3**, such as isopropyl radicals, is much higher than that of **1**. So the phosphinylphosphination of various alkenes with **3** successfully proceeded.

A plausible reaction pathway for the present reaction is illustrated in Figure 1. The  $P^{V}(O)-P^{III}$  bond cleavage of **3** is induced by the radical initiator or near-UV light to generate the diphenylphosphinyl radical  $[Ph_2P(O)^{\cdot}]$ .<sup>[17]</sup> The generated radical reacts with **2** at the terminal carbon atom to generate the alkyl radical **10**. The alkyl radical **10** is captured by **3** to give **4** by an S<sub>H</sub>2-type mechanism.

Preliminary density functional theory (DFT) calculations (B3LYP/6-31G(d)) were performed to explain the observed regioselectivity of the reaction. Regarding the addition of **3** to **20** ( $\mathbf{R} = \mathbf{CH}_3$  in Figure 1 a), the single occupied molecular orbital (SOMO) level of **100** (-4.85 eV) was similar to the highest occupied molecular orbital (HOMO) level of **3** (-5.74 eV), which is localized on the trivalent phosphorus



**Figure 1.** a) A plausible reaction pathway for the highly regioselective phosphinylphosphination of **3** to alkenes. b) Kohn–Sham molecular orbitals (HOMO and LUMO) of **3** calculated at the B3LYP/6-31G(d) level of theory.

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

einheim www.angewandte.org

atom. The lowest unoccupied molecular orbital (LUMO) level of **3** was -1.10 eV. This result indicates that the reactive site of **3** to **100** is at the trivalent phosphorus atom. Therefore, the radical-chain reaction, involving both the radical attack of Ph<sub>2</sub>P(O)<sup>•</sup> to alkenes and the radical capture of **10** by **3**, proceeds to afford **4**, selectively.<sup>[18]</sup>

In summary, we have developed a protocol for the highly regioselective addition of tetraphenyldiphosphine monoxides to a variety of alkenes. This approach enables the regioselective introduction of diphenylphosphino and diphenylphosphinyl groups on various aliphatic alkenes and phenylacetylene. A plausible reaction pathway involving a regioselective radical capturing process was proposed. We believe this method will facilitate significant opportunities in the synthesis of bis(phosphine) monoxide ligands.

#### Acknowledgments

This research was supported by a Grant-in-Aid for Exploratory Research (26620149, A.O., 26860168, S.K.) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Keywords: phosphanes · phosphorus · radicals · regioselectivity

- [1] L. D. Quin, *Guide to Organophosphorus Chemistry*, Wiley, New York, **2000**.
- [2] For the addition reactions of *H*-phosphonates and phosphine oxides to carbon-carbon multiple bonds, see: a) Q. Xu, L.-B. Han, *J. Organomet. Chem.* 2011, 696, 130–140; b) T. Bunlaksananusorn, P. Knochel, *Tetrahedron Lett.* 2002, 43, 5817–5819; c) L. L. Khemchyan, J. V. Ivanova, S. S. Zalesskiy, V. P. Ananikov, I. P. Beletskaya, Z. A. Starikova, *Adv. Synth. Catal.* 2014, 356, 771–780.
- [3] H. Yorimitsu, Beilstein J. Org. Chem. 2013, 9, 1269-1277.
- [4] a) E. R. Fruchey, B. M. Monks, S. P. Cook, J. Am. Chem. Soc. 2014, 136, 13130-13133; b) J.-H. Fan, W.-T. Wei, M.-B. Zhou, R.-J. Song, J.-H. Li, Angew. Chem. Int. Ed. 2014, 53, 6650-6654; Angew. Chem. 2014, 126, 6768-6772; c) S. Uesugi, Z. Li, R. Yazaki, T. Ohshima, Angew. Chem. Int. Ed. 2014, 53, 1611-1615; Angew. Chem. 2014, 126, 1637-1641; d) M. Arisawa, T. Ichikawa, M. Yamaguchi, Chem. Commun. 2015, 51, 8821-8824.
- [5] a) A. Tzschach, S. Baensch, J. Prakt. Chem. 1971, 313, 254–258;
   b) A. Sato, H. Yorimitsu, K. Oshima, Angew. Chem. Int. Ed. 2005, 44, 1694–1696; Angew. Chem. 2005, 117, 1722–1724;

c) D. L. Dodds, M. F. Haddow, A. G. Orpen, P. G. Pringle, G. Woodward, *Organometallics* **2006**, *25*, 5937–5945; d) S-i. Kawaguchi, S. Nagata, T. Shirai, K. Tsuchii, A. Nomoto, A. Ogawa, *Tetrahedron Lett.* **2006**, *47*, 3919–3922.

- [6] A. B. Burg, J. Am. Chem. Soc. 1961, 83, 2226-2231.
- [7] M. Drieß, G. Haiber, Z. Anorg. Allg. Chem. 1993, 619, 215-219.
- [8] a) K. W. Morse, J. G. Morse, J. Am. Chem. Soc. 1973, 95, 8469– 8470; b) J. G. Morse, K. W. Morse, Inorg. Chem. 1975, 14, 565– 569.
- [9] I. Hajdók, F. Lissner, M. Nieger, S. Strobel, D. Gudat, Organometallics 2009, 28, 1644–1651.
- [10] a) A. Ogawa, H. Tanaka, H. Yokoyama, R. Obayashi, K. Yokoyama, N. Sonoda, *J. Org. Chem.* **1992**, *57*, 111–115; b) A. Ogawa, R. Obayashi, M. Doi, N. Sonoda, T. Hirao, *J. Org. Chem.* **1998**, *63*, 4277–4281; c) A. Ogawa, I. Ogawa, R. Obayashi, K. Umezu, M. Doi, T. Hirao, *J. Org. Chem.* **1999**, *64*, 86–92.
- [11] V. V. Grushin, Chem. Rev. 2004, 104, 1629-1662.
- [12] a) A. Côté, A. B. Charette, J. Org. Chem. 2005, 70, 10864-10867; b) A. Côté, V. N. G. Lindsay, A. B. Charette, Org. Lett. 2007, 9, 85-87; c) M. McConville, O. Saidi, J. Blacker, J. Xiao, J. Org. Chem. 2009, 74, 2692-2698; d) A. Hamada, P. Braunstein, Organometallics 2009, 28, 1688-1696; e) T. H. Wöste, M. Oestreich, Chem. Eur. J. 2011, 17, 11914-11918; f) J. Hu, Y. Lu, Y. Li, J. Zhou, Chem. Commun. 2013, 49, 9425-9427; g) C. Hahn, L. Cruz, A. Villalobos, L. Garza, S. Adeosun, Dalton Trans. 2014, 43, 16300-16309; h) C. Li, T. Chen, B. Li, G. Xiao, W. Tang, Angew. Chem. Int. Ed. 2015, 54, 3792-3796; Angew. Chem. 2015, 127, 3863-3867; i) J. Shi, T. Wang, Y. Huang, X. Zhang, Y.-D. Wu, Q. Cai, Org. Lett. 2015, 17, 840-843; j) G. M. Borrajo-Calleja, V. Bizet, C. Mazet, J. Am. Chem. Soc. 2016, 138, 4014-4017; k) Y. Fukuzaki, Y. Tomita, T. Terashima, M. Ouchi, M. Sawamoto, Macromolecules 2010, 43, 5989-5995; 1) B. P. Carrow, K. Nozaki, J. Am. Chem. Soc. 2012, 134, 8802-8805.
- [13] A. L. J. Beckwith, Tetrahedron 1981, 37, 3073-3100.
- [14] M. Kamachi, A. Kajiwara, K. Saegusa, Y. Morishima, *Macro-molecules* **1993**, 26, 7369–7371.
- [15] A. Kajiwara, Y. Konishi, Y. Morishima, W. Schnabel, K. Kuwata, M. Kamachi, *Macromolecules* **1993**, *26*, 1656–1658.
- [16] a) A. Sato, H. Yorimitsu, K. Oshima, J. Am. Chem. Soc. 2006, 128, 4240-4241; b) S. E. Vaillard, C. Mück-Lichtenfeld, S. Grimme, A. Studer, Angew. Chem. Int. Ed. 2007, 46, 6533-6536; Angew. Chem. 2007, 119, 6653-6656.
- [17] The diphenylphosphino radical (Ph<sub>2</sub>P·) is also generated by near-UV light. However, Ph<sub>2</sub>P(O)' is more reactive with alkenes than Ph<sub>2</sub>P· and initiates the radical-chain reaction. In this reaction, the P<sup>V</sup>(O)-P<sup>III</sup> bond cleavage, induced by near-UV light, is much slower than the radical chain reaction.
- [18] 1,2-Bisphosphinoalkane was obtained in <1% yield.

Received: April 21, 2016 Published online: ■■ ■■, ■■■

These are not the final page numbers!



## **Communications**



## **Communications**

### Phosphanes

Y. Sato, S-i. Kawaguchi,\* A. Nomoto, A. Ogawa\* \_\_\_\_\_

Highly Selective

Phosphinylphosphination of Alkenes with Tetraphenyldiphosphine Monoxide



 $P(O)P by: Ph_2P(O)PPh_2$  can engage in radical addition to various alkenes, thus affording the corresponding 1-phosphinyl-2-phosphinoalkanes regioselectively, and they can be converted into their sulfides by treatment with S<sub>8</sub>. The reaction proceeds by homolytic cleavage of the  $P^{V}(O)-P^{|||}$  bond of  $Ph_2P(O)PPh_2$ , selective attack of the phosphinyl radical at the terminal position of the alkene, and selective trapping of the resulting carbon radical by the phosphino group.

einheim www.angewandte.org These are not the final page numbers!

5