

Synthetic Methods


Control of Selectivity in Palladium-Catalyzed Oxidative Carbocyclization/Borylation of Allenynes**

Youqian Deng, Teresa Bartholomeyzik, and Jan-E. Bäckvall*

Organoboronates are convenient and versatile reagents owing to their comparatively low toxicity, high functional group compatibility, and good stability.^[1] Moreover, these compounds can easily be oxidized to alcohols^[2] or used to construct new C–C bonds by Suzuki–Miyaura cross-couplings.^[3] Because of the broad applications of these boronates, many borylation methods have been developed.^[4–7] Amongst the routes reported for C–B bond formation the most common are Miyaura borylation,^[4] hydroboration,^[5] and the reaction of lithium or magnesium organometallic compounds with borate esters.^[1] In addition, recent developments in transition-metal-catalyzed C–H borylation reactions have also provided efficient access to boronates.^[6]

Furthermore, by combining the borylation with a C–C bond-forming cyclization, complex molecules suitable for various further functionalizations could be obtained in one step.^[8–11,13b,d] Such borylating carbocyclizations have been successfully developed by the group of Cárdenas.^[9–11] Starting from unsaturated compounds, such as enynes,^[9] enediynes,^[10] enallenes,^[11] and allenynes,^[11] homoallylic or allylic boronates were prepared under palladium(0) catalysis. For instance, the non-oxidative borylating carbocyclization of allenynes **1** in the presence of bis(pinacolato)diboron (B₂pin₂) yielded two isomers (**2** and **3**), where borylation occurred at the allene (Scheme 1a).^[11]

In ongoing investigations our research group has been studying oxidative Pd^{II}-catalyzed carbocyclizations of various unsaturated molecules.^[12–15] Recently we accomplished the carbocyclization/arylation of allenynes with arylboronic acids.^[13b] Also, some preliminary results regarding carbocyclization/borylation were obtained with differently substituted 1,5-allenynes (**1**; R = H, Me, Ar), which only gave borylated triene products **4** (Scheme 1b).^[13b] However, under carbocy-



Scheme 1. Palladium-catalyzed borylating carbocyclizations of allenynes: a) under non-oxidative conditions;^[11] b) under oxidative conditions;^[13b] c) under selective oxidative conditions. E = CO₂Me.

clization/arylation conditions alkyl-substituted allenynes afforded two different constitutional isomers (arylated trienes and arylated vinylallenes) in a ratio determined by the substitution on the starting allenyne.^[13b] The aim of the present study was to develop a carbocyclization/borylation that can be directed towards either a borylated triene or a borylated vinylallene by control of the reaction conditions (Scheme 1c). We now report a highly selective oxidative carbocyclization/borylation of allenynes **1** with B₂pin₂ under Pd^{II} catalysis with *p*-benzoquinone (BQ) as the oxidant. The use of LiOAc·2H₂O in 1,2-dichloroethane (DCE) or BF₃·Et₂O in THF addressed the issue of selectivity, to give either borylated trienes **4** or borylated vinylallenes **5**, respectively.

We first studied the reaction of ethyl-substituted allenyne **1a** with B₂pin₂ under the original carbocyclization/borylation conditions (Scheme 1b).^[13b] The use of a catalytic amount of palladium acetate (2 mol %) and stoichiometric amounts of BQ (1.1 equiv) in THF at 50 °C led to an isomeric mixture of borylated triene **4a** and borylated vinylallene **5a** in 28 % and 14 % yield, respectively (Table 1, entry 1). Analyzing the effect of different solvents showed that a higher selectivity for **4a** was obtained when DCE was used as the solvent (in Table 1, entry 4 vs. entries 1–3). Furthermore, upon the addition of catalytic amounts (20 mol %) of a basic salt,

[*] Dr. Y. Deng, T. Bartholomeyzik, Prof. Dr. J.-E. Bäckvall
Department of Organic Chemistry, Arrhenius Laboratory
Stockholm University
SE-106 91 Stockholm (Sweden)
E-mail: jeb@organ.su.se

[**] Financial support from the European Research Council (ERC AdG 247014), the Swedish Research Council, the Berzelii Centre EXSELENT, the Knut and Alice Wallenberg Foundation, and the Wenner-Gren Foundation (Y.D.) is gratefully acknowledged.


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201301167>.

© 2013 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

moiety (**1c** and **1e**) a comparatively high selectivity for the triene products was observed (Table 2, entries 3 and 5). However, a mixture of borylated triene products was obtained, with a preference for formation of the products with the more substituted double bond. Products **4c** and **4e** were obtained as a mixture of *Z/E* isomers in a ratio of 3.3:1 and 3.5:1, respectively. The allenyne **1f** with a cyclohexylidene group on the allene cyclized to give a mixture of isomers **4f** and **4f'**, where the formation of **4f** could be explained by a Pd-catalyzed isomerization of **4f** (Table 2, entry 6; for the detailed mechanism, please see the Supporting Information). Moreover, the reaction of allenyne **1g**, having two benzyl ether groups on the linker part X, also gave borylated triene product **4g** selectively in 57% yield (Table 2, entry 7), thus proving that the malonate group of linker X is not necessary for a successful transformation.

The optimized reaction conditions for the selective formation of borylated vinylallene **5** (20 mol % of $\text{BF}_3 \cdot \text{Et}_2\text{O}$,


Table 3: Selective carbocyclization of allenyne **1** yielding borylated vinylallenes **5**.^[a]

				
Entry	Allenyne	Product	5/4 ^[b]	Yield of 5 [%] ^[c]
1			> 20:1	77
2			> 20:1	73
3			> 20:1	56
4			> 20:1	79 87 ^[d]
5			20:1	77
6			> 20:1	70
7 ^[e]			> 20:1	37

[a] Unless otherwise noted, **1** (0.1–0.2 mmol), B_2pin_2 (1.3 equiv), $\text{Pd}(\text{OAc})_2$ (2 mol %), BQ (1.1 equiv), and $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (20 mol %) were dissolved in THF (5 mL mmol⁻¹) and stirred at 50 °C for 20 h. [b] Ratio determined by ¹H NMR analysis of the crude reaction mixture. [c] Yield of the isolated product. [d] 1 mmol of **1d** was used. [e] 2 mol % of $[\text{Pd}(\text{CH}_3\text{CN})_4][(\text{BF}_4)_2]$ was used in place of $\text{Pd}(\text{OAc})_2$ and $\text{BF}_3 \cdot \text{Et}_2\text{O}$. E = CO_2Me .

Method B) were applied to various allenyne (Table 3). Allenyne **1a–1f** were transformed into vinylallenic boronates **5a–5f**; for most cases the yield was between 70% and 80% and the formation of the corresponding triene isomers **4a–4f** was efficiently suppressed. Even the methyl-substituted substrate **1b**, which intrinsically favors formation of triene **4b**,^[13b] displayed opposite selectivity under these reaction conditions, that is, favoring vinylallene formation (Table 3, entry 2). The reaction of allenyne **1g** under the standard conditions of Table 3 was sluggish and did not give the desired product **5g**, probably because of the incompatibility between the benzyl ether group and $\text{BF}_3 \cdot \text{Et}_2\text{O}$. However, by switching the palladium catalyst to $[\text{Pd}(\text{CH}_3\text{CN})_4][(\text{BF}_4)_2]$ and in the absence of $\text{BF}_3 \cdot \text{Et}_2\text{O}$, product **5g** was obtained in 37% yield (entry 7).


To gain further insights into the mechanism of the oxidative carbocyclization/borylation, kinetic deuterium isotope effects were studied (Scheme 2). An intermolecular competition experiment using **1d** and its hexadeuterated derivative $[\text{D}_6]\text{-1d}$ under the conditions for selective triene formation for 1 h provided a large intermolecular KIE value of 6.7^[17] (Scheme 2a). This result indicates that the allylic C–H bond cleavage involved has to occur prior to any irreversible step of the reaction, for example, the carbocyclization step.^[18] On the other hand, when a 1:1 mixture of **1d** and $[\text{D}_2]\text{-1d}$ was subjected to the conditions for selective vinylallene formation for 1 h the ratio between **5d** and $[\text{D}_1]\text{-5d}$ was 2.4, from which the KIE was determined to 2.7^[19] (Scheme 2b).^[17] The intrinsic KIE from intramolecular competition for vinylallene formation was determined to 5.3^[17] by the use of $[\text{D}_1]\text{-1d}$ as the allenyne substrate (Scheme 2c). The results in Scheme 2b and 2c indicate that the propargylic C–H bond cleavage does not fully determine the selectivity



Scheme 2. Kinetic isotope effect study. Reaction conditions: i) B_2pin_2 (1.3 equiv), $\text{Pd}(\text{OAc})_2$ (2 mol %), BQ (1.1 equiv), $\text{LiOAc} \cdot 2\text{H}_2\text{O}$ (20 mol %), DCE, 50 °C, 1 h. ii) B_2pin_2 (1.3 equiv), $\text{Pd}(\text{OAc})_2$ (2 mol %), BQ (1.1 equiv), $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (20 mol %), THF, 50 °C, 1 h for b) and 20 h for c). E = CO_2Me .

between **5d** and **[D₁]-5d** in the competitive experiment (Scheme 2b).^[18]


Three control experiments with allenyne **1d** and the corresponding deuterium-labeled allenynes **[D₂]-1d** and **[D₆]-1d** were conducted under palladium catalysis in the absence of any additional basic or acidic additive and using DCE as the solvent (Scheme 3). Under these conditions the reaction



Scheme 3. Effect of isotope substitution on product distribution. Reaction conditions: i) B_2pin_2 (1.3 equiv), $Pd(OAc)_2$ (2 mol %), BQ (1.1 equiv), DCE, 50 °C, 15 h. ii) B_2pin_2 (1.3 equiv), $Pd(OAc)_2$ (5 mol %), BQ (1.1 equiv), DCE, 50 °C, 15 h. E = CO_2Me .

of **1d** gave a mixture of **4d** and **5d** in a ratio of 3.8:1 (Scheme 3a). When substrate **[D₂]-1d** was employed (Scheme 3b) under the same reaction conditions, the ratio increased to 12.3:1.^[20] Allenyne **[D₆]-1d** showed the opposite selectivity, with **[D₅]-4d** and **[D₆]-5d** being formed in a ratio of 1:5.1^[20] (Scheme 3c).

The results in Scheme 2 and Scheme 3 indicate that competing allylic and propargylic C–H bond cleavage occurs in **1**, and this determines the ratio of boronates **4** and **5** (Scheme 4). The allene attack on Pd^{II} complex **A** through allylic C–H bond cleavage^[12,13a–d] would give **B** and subsequent alkyne insertion would generate intermediate **C**.



Scheme 4. Proposed mechanism for palladium-catalyzed oxidative selective carbocyclization/borylation of allenyne **1**.

Transmetalation of **C** with B_2pin_2 and reductive elimination would form product **4**. The competing alkyne attack through propargylic C–H bond cleavage in **A** would produce allenyl-palladium intermediate **D**. Intramolecular vinylpalladation of the allene moiety would generate $(\pi\text{-allyl})$ palladium intermediate **E**. Transmetalation with B_2pin_2 and subsequent reductive elimination would give **5**. The mechanism in Scheme 4 is supported by the kinetic isotope effects and the experiments with deuterium-labeled compounds (Scheme 2 and Scheme 3). The lower kinetic isotope effect observed for the competitive experiment in Scheme 2b compared to the intramolecular experiment in Scheme 2c may reflect that **1** and **A** are not in full equilibrium under the conditions for formation of **5**. In the path for formation of **5** it is likely that $BF_3 \cdot Et_2O$ creates a cationic palladium species, which interacts better with the acetylene compared to the allene in **A**.^[21]

In summary, we have developed an unprecedented selective Pd^{II} -catalyzed carbocyclization/borylation of allenynes under oxidative conditions. By controlling the reaction conditions the reaction can be directed to either the triene **4** or the vinylallene **5**. On the basis of the results of deuterium-labeling experiments, we propose that the reactions of allenynes proceed through competing allylic and propargylic C–H bond cleavage pathways to give borylated trienes and borylated vinylallenes, respectively.

Experimental Section

Typical experimental procedure for palladium-catalyzed oxidative borylating carbocyclization of allenyne **1** to boronate **4**: **1a** (26.0 mg, 0.10 mmol) and 0.5 mL of DCE were added to a mixture of B_2pin_2 (33.1 mg, 0.13 mmol), BQ (12.2 mg, 0.11 mmol), $Pd(OAc)_2$ (0.5 mg, 0.002 mmol), and $LiOAc \cdot 2H_2O$ (1.8 mg, 0.02 mmol) at RT. The reaction was stirred at 50 °C for 15 h. After the reaction was complete, as monitored by TLC, evaporation and column chromatography on silica gel (pentane/ethyl acetate = 10:1) afforded **4a** (27.9 mg, 73 %) as a liquid; 1H NMR (500 MHz, $CDCl_3$): δ = 5.93 (s, 1H), 5.02–5.00 (m, 2H), 3.72 (s, 6H), 3.20 (s, 2H), 2.19 (q, J = 7.5 Hz, 2H), 1.95 (s, 3H), 1.26 (s, 12H), 1.02 ppm (t, J = 7.5 Hz, 3H); ^{13}C NMR (125 MHz, $CDCl_3$): δ = 171.1, 151.1, 147.9, 139.8, 129.1, 116.6, 83.3, 63.0, 52.9, 37.5, 27.0, 25.2, 23.8, 13.4 ppm; HRMS (ESI): calc. for $C_{21}H_{31}BNaO_6$ [$M + Na$] $^+$: 413.2110; found: 413.2113.

Typical experimental procedure for palladium-catalyzed oxidative borylating carbocyclization of allenyne **1** to boronate **5**: **1a** (52.7 mg, 0.20 mmol) and 1.0 mL of THF were added to a mixture of B_2pin_2 (66.2 mg, 0.26 mmol), BQ (24.0 mg, 0.22 mmol), $Pd(OAc)_2$ (1.0 mg, 0.004 mmol), and $BF_3 \cdot Et_2O$ (6 μ L, 0.04 mmol) at RT. The reaction was stirred at 50 °C for 20 h. After the reaction was complete, as monitored by TLC, evaporation and column chromatography on silica gel (pentane/ethyl acetate = 10/1) afforded **5a** (59.6 mg, 77 %) as a liquid; 1H NMR (400 MHz, $CDCl_3$): δ = 5.55 (d, J = 1.6 Hz, 1H), 5.34–5.23 (m, 1H), 3.716 (s, 3H), 3.715 (s, 3H), 3.19–3.17 (m, 2H), 1.68 (d, J = 7.2 Hz, 3H), 1.18 (s, 12H), 1.17 (s, 3H), 1.13 ppm (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$): δ = 199.1, 171.5, 171.3, 153.8, 122.9, 107.1, 91.2, 83.1, 63.5, 52.7, 36.5, 25.0, 24.7, 24.5, 23.9, 23.8, 14.8 ppm; HRMS (ESI): calc. for $C_{21}H_{31}BNaO_6$ [$M + Na$] $^+$: 413.2110; found: 413.2103.

Received: February 8, 2013
Published online: May 17, 2013

Keywords: allenynes · boronates · cyclization · oxidation · palladium

- [1] *Boronic Acids: Preparation and Applications in Organic Synthesis and Medicine* (Ed.: D. G. Hall), Wiley-VCH, Weinheim, **2005**.
- [2] For selected references, see: a) H. C. Brown, B. C. S. Rao, *J. Am. Chem. Soc.* **1959**, *81*, 6434; b) T. Ishiyama, T. Ahiko, N. Miyaoura, *J. Am. Chem. Soc.* **2002**, *124*, 12414; c) J. W. J. Kennedy, D. G. Hall, *J. Am. Chem. Soc.* **2002**, *124*, 11586; d) R. E. Maleczka, Jr., F. Shi, D. Holmes, M. R. Smith III, *J. Am. Chem. Soc.* **2003**, *125*, 7792.
- [3] For reviews, see: a) N. Miyaoura, A. Suzuki, *Chem. Rev.* **1995**, *95*, 2457; b) A. Suzuki, *J. Organomet. Chem.* **1999**, *576*, 147; c) N. Miyaoura, *Top. Curr. Chem.* **2002**, *219*, 248; d) N. Miyaoura in *Metal-Catalyzed Cross-Coupling Reactions* (Eds.: A. de Meijere, F. Diederich), Wiley-VCH, Weinheim, **2004**, pp. 41–123.
- [4] For selected reviews, see: a) T. Ishiyama, N. Miyaoura, *Chem. Rev.* **2004**, *3*, 271; b) N. Miyaoura, *Top. Curr. Chem.* **2002**, *219*, 11.
- [5] For selected references for the Brown hydroboration, see: a) H. C. Brown, B. C. S. Rao, *J. Am. Chem. Soc.* **1959**, *81*, 6423; b) H. C. Brown, B. C. S. Rao, *J. Am. Chem. Soc.* **1959**, *81*, 6428; c) J. M. Clay, E. Vedejs, *J. Am. Chem. Soc.* **2005**, *127*, 5766.
- [6] For a review of borylation of C–H bonds, see: a) I. A. I. Mkhaliid, J. H. Barnard, T. B. Marder, J. M. Murphy, J. F. Hartwig, *Chem. Rev.* **2010**, *110*, 890; for recent selected reports of transition-metal-catalyzed C–H borylations, see: b) T. Ishiyama, H. Isou, T. Kikuchi, N. Miyaoura, *Chem. Commun.* **2010**, *46*, 159; c) K. Yamazaki, S. Kawamorita, H. Ohmiya, M. Sawamura, *Org. Lett.* **2010**, *12*, 3978; d) N. Selander, B. Willy, K. J. Szabó, *Angew. Chem.* **2010**, *122*, 4145; *Angew. Chem. Int. Ed.* **2010**, *49*, 4051; e) H. Itoh, T. Kikuchi, T. Ishiyama, N. Miyaoura, *Chem. Lett.* **2011**, *40*, 1007; f) A. Ros, B. Estepa, R. López-Rodríguez, E. Álvarez, R. Fernández, J. M. Lassaletta, *Angew. Chem.* **2011**, *123*, 11928; *Angew. Chem. Int. Ed.* **2011**, *50*, 11724; g) S. Kawamorita, T. Miyazaki, H. Ohmiya, T. Iwai, M. Sawamura, *J. Am. Chem. Soc.* **2011**, *133*, 19310; h) H.-X. Dai, J.-Q. Yu, *J. Am. Chem. Soc.* **2012**, *134*, 134; i) B. Xiao, Y.-M. Li, Z.-J. Liu, H.-Y. Yang, Y. Fu, *Chem. Commun.* **2012**, *48*, 4854.
- [7] For a review of catalytic carboborations, see: a) M. Suginoe, *Chem. Rev.* **2010**, *10*, 348; b) M. Daini, M. Suginoe, *J. Am. Chem. Soc.* **2011**, *133*, 4758; c) M. Raducan, R. Alam, K. J. Szabó, *Angew. Chem.* **2012**, *124*, 13227; *Angew. Chem. Int. Ed.* **2012**, *51*, 13050.
- [8] a) A. R. Burns, J. S. González, H. W. Lam, *Angew. Chem.* **2012**, *124*, 10985; *Angew. Chem. Int. Ed.* **2012**, *51*, 10827; b) H. Ito, T. Toyoda, M. Sawamura, *J. Am. Chem. Soc.* **2010**, *132*, 5990; c) T. Miura, Y. Takahashi, M. Murakami, *Org. Lett.* **2008**, *10*, 1743; d) A. M. Camelio, T. Barton, F. Guo, T. Shaw, D. Siegel, *Org. Lett.* **2011**, *13*, 1517; e) W. Gong, R. R. Singidi, J. C. Gallucci, T. V. RajanBabu, *Chem. Sci.* **2012**, *3*, 1221; f) R. R. Singidi, A. M. Kutney, J. C. Gallucci, T. V. RajanBabu, *J. Am. Chem. Soc.* **2010**, *132*, 13078; g) S. Onozawa, Y. Hatanaka, N. Choi, M. Tanaka, *Organometallics* **1997**, *16*, 5389; h) S. Onozawa, Y. Hatanaka, M. Tanaka, *Chem. Commun.* **1997**, 1229.
- [9] a) J. Marco-Martínez, V. López-Carrillo, E. Buñuel, R. Simancas, D. J. Cárdenas, *J. Am. Chem. Soc.* **2007**, *129*, 1874; b) V. Pardo-Rodríguez, E. Buñuel, D. Collado-Sanz, D. J. Cárdenas, *Chem. Commun.* **2012**, *48*, 10517.
- [10] a) J. Marco-Martínez, E. Buñuel, R. Muñoz-Rodríguez, D. J. Cárdenas, *Org. Lett.* **2008**, *10*, 3619; b) J. Marco-Martínez, E. Buñuel, R. López-Durán, D. J. Cárdenas, *Chem. Eur. J.* **2011**, *17*, 2734.
- [11] V. Pardo-Rodríguez, J. Marco-Martínez, E. Buñuel, D. J. Cárdenas, *Org. Lett.* **2009**, *11*, 4548.
- [12] a) J. Löfstedt, J. Franzén, J.-E. Bäckvall, *J. Org. Chem.* **2001**, *66*, 8015; b) J. Franzén, J.-E. Bäckvall, *J. Am. Chem. Soc.* **2003**, *125*, 6056; c) J. Piera, K. Närhi, J.-E. Bäckvall, *Angew. Chem.* **2006**, *118*, 7068; *Angew. Chem. Int. Ed.* **2006**, *45*, 6914; d) J. Piera, A. Persson, X. Caldentey, J.-E. Bäckvall, *J. Am. Chem. Soc.* **2007**, *129*, 14120; e) E. V. Johnston, E. A. Karlsson, S. A. Lindberg, B. Åkermarck, J.-E. Bäckvall, *Chem. Eur. J.* **2009**, *15*, 6799; f) A. K. Å. Persson, J.-E. Bäckvall, *Angew. Chem.* **2010**, *122*, 4728; *Angew. Chem. Int. Ed.* **2010**, *49*, 4624; for a DFT calculation for allene attack on a Pd^{II} center, see: g) E. A. Karlsson, J.-E. Bäckvall, *Chem. Eur. J.* **2008**, *14*, 9175.
- [13] a) Y. Deng, J.-E. Bäckvall, *Angew. Chem.* **2013**, *125*, 3299; *Angew. Chem. Int. Ed.* **2013**, *52*, 3217; b) Y. Deng, T. Bartholomeyzik, A. K. Å. Persson, J. Sun, J.-E. Bäckvall, *Angew. Chem.* **2012**, *124*, 2757; *Angew. Chem. Int. Ed.* **2012**, *51*, 2703; c) T. Jiang, A. K. Å. Persson, J.-E. Bäckvall, *Org. Lett.* **2011**, *13*, 5838; d) A. K. Å. Persson, T. Jiang, M. T. Johnson, J.-E. Bäckvall, *Angew. Chem.* **2011**, *123*, 6279; *Angew. Chem. Int. Ed.* **2011**, *50*, 6155; e) M. Jiang, T. Jiang, J.-E. Bäckvall, *Org. Lett.* **2012**, *14*, 3538.
- [14] For selected recent oxidative carbocyclization reactions by other groups, see: a) K.-T. Yip, D. Yang, *Org. Lett.* **2011**, *13*, 2134; b) B. S. Matsuura, A. G. Condie, R. C. Buff, G. J. Karahalidis, C. R. J. Stephenson, *Org. Lett.* **2011**, *13*, 6320; c) T. Wu, X. Mu, G. Liu, *Angew. Chem.* **2011**, *123*, 12786; *Angew. Chem. Int. Ed.* **2011**, *50*, 12578; d) X. Mu, T. Wu, H. Wang, Y. Guo, G. Liu, *J. Am. Chem. Soc.* **2012**, *134*, 878; e) R. Zhu, S. L. Buchwald, *Angew. Chem.* **2012**, *124*, 1962; *Angew. Chem. Int. Ed.* **2012**, *51*, 1926; f) Y. Wei, I. Deb, N. Yoshikai, *J. Am. Chem. Soc.* **2012**, *134*, 9098.
- [15] For selected reviews involving palladium-catalyzed oxidative carbocyclizations, see: a) E. M. Beccalli, G. Broggini, M. Martinelli, S. Sottocornola, *Chem. Rev.* **2007**, *107*, 5318; b) F. Dénès, A. Pérez-Luna, F. Chemla, *Chem. Rev.* **2010**, *110*, 2366; c) C. S. Yeung, V. M. Dong, *Chem. Rev.* **2011**, *111*, 1215; d) Y. Deng, A. K. Å. Persson, J.-E. Bäckvall, *Chem. Eur. J.* **2012**, *18*, 11498.
- [16] J. M. Larsson, K. J. Szabó, *J. Am. Chem. Soc.* **2013**, *135*, 443.
- [17] A small normal secondary isotope effect will affect the value.
- [18] E. M. Simmons, J. F. Hartwig, *Angew. Chem.* **2012**, *124*, 3120; *Angew. Chem. Int. Ed.* **2012**, *51*, 3066.
- [19] At the end of the reaction the starting material ratio was estimated to 1:1.2 and the product ratio was 2.43:1. Therefore, the isotope effect calculated from the product ratio and the change of the starting material ratio is approximately $2.43 \times [(1 + 1.2)/2] = 2.7$ (see the Supporting Information).
- [20] A normal secondary deuterium isotope affects the ratio to some extent.
- [21] For another possible mechanism involving a pallada(IV)cyclopentene intermediate^[13a,b] (see the Supporting Information).