

Subscriber access provided by - Access paid by the | UCSB Libraries

### Transition-Metal-free Synthesis of Ynones via Decarboxylative Alkynylation of #-Keto Acids under Mild Conditions

Peng-Fei Wang, Yi-Si Feng, Zhi-Fei Cheng, Qiu-Min Wu, Guang-Yu Wang, Liang-Liang Liu, Jian-Jun Dai, Jun Xu, and Hua-Jian Xu

J. Org. Chem., Just Accepted Manuscript • DOI: 10.1021/acs.joc.5b01387 • Publication Date (Web): 24 Aug 2015 Downloaded from http://pubs.acs.org on August 25, 2015

#### Just Accepted

Note

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



The Journal of Organic Chemistry is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties. Page 1 of 23

## Transition-Metal-free Synthesis of Ynones via Decarboxylative Alkynylation of α-Keto Acids under Mild Conditions

Peng-Fei Wang,<sup>†</sup> Yi-Si Feng,<sup>\*†,‡</sup> Zhi-Fei Cheng,<sup>†</sup> Qiu-Min Wu,<sup>†</sup> Guang-Yu Wang,<sup>†</sup>

Liang-Liang Liu,<sup>†</sup> Jian-Jun Dai,<sup>§</sup> Jun Xu,<sup>§</sup> and Hua-Jian Xu\*<sup>‡,§</sup>

<sup>†</sup>School of Chemistry and Chemical Engineering, Hefei University of Technology, Hefei 230009, P. R. China

<sup>‡</sup>Anhui Key Laboratory of Controllable Chemical Reaction & Material Chemical Engineering, Hefei 230009, P. R. China

<sup>§</sup>School of Medical Engineering, Hefei University of Technology, Hefei 230009, P. R. China



**ABSTRACT:** A transition-metal-free synthetic method of various ynones via decarboxylative alkynylation of  $\alpha$ -keto acids is described. The reaction is carried out under mild conditions and exhibits remarkable tolerance of functional groups. The mechanism of a radical process is proposed in the reaction.

Ynones are highly valuable and amusing compounds that are widely employed as fascinating precursors of heterocyclic compounds in organic synthesis.<sup>1</sup> Typical approaches for the synthesis of ynones involve Sonogashira couplings reactions of terminal alkynes with acyl chlorides utilizing Pd/Cu as catalysts<sup>2</sup> or carbonylative couplings reactions between terminal alkynes and aryl or alkyl halides,<sup>3</sup> aryl triflates,<sup>4</sup> aryl amines<sup>5</sup> with CO insertion. Recently, other methods have also emerged as new strategies for the synthesis of ynones. For example, Hashmi et al reported a gold-catalyzed dehydrogenative Meyer-Schuster-like rearrangement to produce

ynones (Scheme 1, eq 1).<sup>6</sup> Huang et al reported a general synthesis of ynones from aldehydes via oxidative C–C bond cleavage under aerobic conditions (Scheme 1, eq 2).<sup>7</sup> More recently, direct alkynylation of aldehydes to synthesize ynones via nucleophilic addition/Oppenauer oxidation and C-H activation have also been reported.<sup>8</sup> When our study work is accomplished, Chen et al reported that dual hypervalent iodine(III) reagents and photoredox catalysis enabled decarboxylative ynonylation (Scheme 1, eq 3).<sup>9</sup> Almost meanwhile, Wang et al reported a sunlight-driven decarboxylative alkynylation of a-keto acids with bromoacetylenes by hypervalent iodine reagent catalysis (Scheme 1, eq 4).<sup>10</sup> Although considerable efforts and significant progress have been made to prepare ynones, most synthetic methods have some shortcomings, such as the usage of toxic CO, harsh reaction conditions, limited substrate scope and expensive catalysts. Therefore, further development of preparing ynones is of great demand to enable mild reaction conditions as well as available raw materials. In order to achieve this goal, it is greatly important to explore new approaches for the efficient synthesis of ynones.

Transition-metal-catalyzed decarboxylative cross-coupling reactions of carboxylic acids represent one of most important methods for the construction of C-C and C-heteroatom bonds.<sup>11-16</sup> These reactions don't require expensive organometallic reagents and produce nontoxic CO<sub>2</sub>. In 2002, Myers first reported a palladium-catalyzed Heck-type olefination of arene carboxylates,<sup>11</sup> subsequently considerable studies on the decarboxylative coupling reactions are conducted by Forgione,<sup>12</sup> Goossen,<sup>13</sup> Liu<sup>14</sup> and others.<sup>15-16</sup> These reactions reveal that decarboxylative couplings provide an alternative method for catalytic synthesis purpose. Although many significant contributions are achieved in this field, decarboxylation of  $\alpha$ -keto acids receives relatively less attention. Decarboxylation of a-keto acids can afford acyl donors

which it is crucial for the preparation of carbonyl compounds. We are interested in developing new strategies to prepare ynones from  $\alpha$ -keto acids under mild conditions. In particular, the reactions that are conducted under transition-metal-free condition are highly desirable to enable potential application because of the significance of ynones. Herein, we reported a general and efficient synthesis of ynones via K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>-promoted decarboxylative alkynylation of  $\alpha$ -keto acids under mild conditions (Scheme 1 eq 5).<sup>17</sup>

Scheme 1. The Synthetic methods of Ynones.



We initiated optimization of the reaction conditions by selecting benzoylformic acid (1a) with 1-[(triisopropylsilyl)ethynyl]-1,2-benziodoxol-3(1H)-one (2a) as model substrates (Table 1). With AgNO<sub>3</sub> (20 mol %) as a catalyst and  $K_2S_2O_8$  (3 equiv) as an oxidant, it was delighted to find that CH<sub>3</sub>CN was used as a solvent to give the desired product **3a** in 12% yield at 50 °C under an argon atmosphere (entry 1). Next, other solvents such as H<sub>2</sub>O, acetone/H<sub>2</sub>O (V:V = 1:1), CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O (V:V = 1:1) and CH<sub>3</sub>CN/H<sub>2</sub>O (V:V = 1:1) were examined (entries 2-5). It was found that CH<sub>3</sub>CN/H<sub>2</sub>O was the optimal medium for the transformation, which brought about an obviously increasing yield of 88%. Then the use of other silver salts such as AgBF<sub>4</sub>, Ag<sub>2</sub>CO<sub>3</sub>, Ag<sub>2</sub>SO<sub>4</sub> also

showed the similar yields (entries 6-8). Furthermore, the use of  $Na_2S_2O_8$  and  $(NH_4)_2S_2O_8$  as oxidants led to the reduction of yield (entries 9-10). Reducing or increasing the reaction temperature didn't improve the yield of **3a** (entries 11-13). The control experiment proved that  $K_2S_2O_8$  was essential, and the reaction still could be conducted to give the desired product **3a** in high yield in the absence of AgNO<sub>3</sub> (entries 14-15). Finally, by changing the amount of  $K_2S_2O_8$ , the yield of **3a** was further improved to 92% (entries 16-19).

 Table 1. Optimization of the Reaction Conditions<sup>a</sup>

	ОН	TIPS	-O Conditions		28
	1a	2a		3a	
entry	catalyst	oxidant (equiv)	solvent	temp (°C)	yield $(\%)^b$
1	AgNO <sub>3</sub>	$K_2S_2O_8(3)$	CH <sub>3</sub> CN	50	12
2	AgNO <sub>3</sub>	$K_2S_2O_8(3)$	$H_2O$	50	trace
3	AgNO <sub>3</sub>	$K_2S_2O_8(3)$	acetone/H <sub>2</sub> O	50	64
4	AgNO <sub>3</sub>	$K_2S_2O_8(3)$	$CH_2Cl_2/H_2O$	50	trace
5	AgNO <sub>3</sub>	$K_2S_2O_8(3)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	88
6	AgBF <sub>4</sub>	$K_2S_2O_8(3)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	87
7	Ag <sub>2</sub> CO <sub>3</sub>	$K_2S_2O_8(3)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	85
8	$Ag_2SO_4$	$K_2S_2O_8(3)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	84
9	AgNO <sub>3</sub>	$Na_2S_2O_8(3)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	79
10	AgNO <sub>3</sub>	$(NH_4)_2S_2O_8(3)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	77
11	AgNO <sub>3</sub>	$K_{2}S_{2}O_{8}(3)$	CH <sub>3</sub> CN/H <sub>2</sub> O	rt(24)	28
12	AgNO <sub>3</sub>	$K_{2}S_{2}O_{8}(3)$	CH <sub>3</sub> CN/H <sub>2</sub> O	40	70
13	AgNO <sub>3</sub>	$K_{2}S_{2}O_{8}(3)$	CH <sub>3</sub> CN/H <sub>2</sub> O	60	82
14	AgNO <sub>3</sub>	-	CH <sub>3</sub> CN/H <sub>2</sub> O	50	-
15	-	$K_{2}S_{2}O_{8}(3)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	89
16	-	$K_2S_2O_8(0.2)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	58
17	-	$K_2S_2O_8(0.4)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	87
18	-	$K_2S_2O_8(0.6)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	92
19	-	$K_2S_2O_8(0.8)$	CH <sub>3</sub> CN/H <sub>2</sub> O	50	91

<sup>*a*</sup>Reaction conditions: **1a** (0.2 mmol), **2a** (0.3 mmol, 1.5 equiv),  $K_2S_2O_8$  (1 equiv = 0.2 mmol), silver salt catalyst (20 mol %), solvent (2 mL), 12 h under an argon atmosphere. <sup>*b*</sup>Isolated yields based on **1a**. TIPS = triisopropylsilyl

Scheme 2. Scope of a-Keto Acids<sup>*a,b*</sup>

#### The Journal of Organic Chemistry



<sup>*a*</sup>Reaction conditions: **1** (0.2 mmol), **2a** (0.3 mmol, 1.5 equiv),  $K_2S_2O_8$  (0.12 mmol, 0.6 equiv),  $CH_3CN/H_2O$  (V:V = 1:1, 2 mL), 50 °C for 12 h under an argon atmosphere. <sup>*b*</sup>Isolated yields based on **1**. <sup>*c*</sup>The reaction was carried out for 24 h.

With the optimal conditions in hand, we explored the scope of  $\alpha$ -keto acids to examine the compatibility of the present reaction (Scheme 2). It was found that various  $\alpha$ -keto acids could smoothly perform decarboxylative alkynylation to generate the desired ynones products in modest to high yields. Both electron-donating substituents such as methy (**3b**, **3k**, **3p**, **3u**), ethyl (**3e**), methoxyl (**3c**, **3l**, **3q**, **3v**), and electron-withdrawing groups such as trifluoromethyl (**3j**) could be well tolerated in current reaction system. It is interesting to note that the reaction could tolerate an unprotected OH group (**3o**), and the corresponding product could further perform cyclization to led to the formation of chromone.<sup>8b</sup> Additionally, the halide groups, such as F (**3f**, **3m**, **3r**), Cl (**3g**, **3n**, **3s**), Br (**3h**, **3t**), I (**3i**) were also well compatible with the reaction, enabling possibly further functionalization at these positions. The  $\alpha$ -keto acids bearing *ortho*- and *meta*-substituents could work well to give the corresponding ynones of 29-87% yields (**3k-t**). The reactions of disubstituted  $\alpha$ -keto acids gave the desired products in 68-82% yields under the above conditions

(**3u-y**). Furthermore, the reaction could also tolerate heterocyclic  $\alpha$ -keto acids such as thiophene, benzo[b]thiophene and indole, giving the desired products in low yields of 31-36% (**3z**, **3aa-ab**). However, no product was detected when 2-(furan-2-yl)-2-oxoacetic acid was employed as a heterocyclic substrate (**3ac**). On the other hand, when aliphatic  $\alpha$ -keto acids were employed for the transformation, the corresponding products were obtained in 48-72% yields (**3ad-ag**).

Next, we further explored the scope of various hypervalent iodine alkynyl reagents (Scheme 3). Alkyne bearing alkyl-substituent such as linear hexyl group could well react with benzoylformic acid **1a** to give the desired products in 52% yield (**4a**). Substrates with bulky silyl groups including SiEt<sub>3</sub>, SiMe<sub>2</sub><sup>t</sup>Bu, Si<sup>t</sup>BuPh<sub>2</sub> groups could be also well applicable for the reaction (**4b-d**). Furthermore, various aryl-substituent alkynes, bearing electron-rich methyl, methoxyl, phenyl, diphenyl and electron-deficient chloride, nitrile, ketone, aldehyde, could successfully convert to the corresponding ynones in 25-81% yields (**4e-l**).

Scheme 3. Scope of Alkynyl Reagents<sup>*a*, *b*</sup>



<sup>*a*</sup>Reaction conditions: **1a** (0.2 mmol), **2** (0.3 mmol, 1.5 equiv),  $K_2S_2O_8$  (0.12 mmol, 0.6 equiv), CH<sub>3</sub>CN/H<sub>2</sub>O (V:V = 1:1, 2 mL), 50 °C for 12 h under an argon atmosphere. <sup>*b*</sup>Isolated yields based on **1a**.

To elucidate the plausible mechanism of reaction, the following experiment was conducted (Scheme 4). Under the optimal conditions, when a radical-trapping reagent TEMPO (2,2,6,6-tetramethyl-1-oxylpiperidine) was added to the reaction, no product was detected. The result suggested that the reaction may proceed by a radical pathway.

#### Scheme 4. Investigation of the Mechanism



According to the above experimental result and the previous studies,<sup>18</sup> we proposed a possible reaction mechanism shown in scheme 5. Firstly, peroxydisulfate ion,  $S_2O_8^{2-}$  performs homolysis to give sulfate ion radical, which reacts with  $\alpha$ -keto acids 1 to produce carboxyl radicals. Carboxyl radical undergoes fast decarboxylation to generate acyl radical, and then attacks the triple bonds of hypervalent iodine alkynyl reagents 2 to achieve the radical intermediate I. Subsequently, I occurs  $\beta$ -elimination to give the desired products and benziodoxolonyl radical II. Finally, 2-iodobenzoic acid is afforded via reduction-protonation of benziodoxolonyl radical II.

#### **Scheme 5. Proposed Reaction Mechanism**



In conclusion, we developed a new efficient reaction system for decarboxylative alkynylation of  $\alpha$ -keto acids using  $K_2S_2O_8$  as an oxidant. The present reaction could tolerate many important functional groups and be conducted under very mild conditions. Furthermore, the reaction

provided a general and efficient method for the synthesis of ynones without transition-metal catalysts.

#### **Experimental section**

#### **General Information and Materials**

All the reactions were conducted in oven-dried Schlenk tubes under an argon atmosphere. All solvents were obtained from commercial suppliers and used without further purification. α-Keto Acids were obtained from commercial suppliers or prepared according to the literature reported procedures.<sup>19</sup> Hypervalent iodine alkynyl reagents were prepared according to the literature reported procedure.<sup>20</sup> <sup>1</sup>H NMR, <sup>13</sup>C NMR, <sup>19</sup>F NMR spectra were recorded on a 600 MHz spectrometer in CDCl<sub>3</sub>. Data for <sup>1</sup>H NMR are reported as follows: chemical shift (δppm), multiplicity, integration, and coupling constant (Hz). Data for <sup>13</sup>C NMR are reported in terms of chemical shift (δppm), multiplicity. High-resolution mass spectra were obtained by ESI on a TOF mass analyzer.

# Typical procedure for the synthesis of ynones via decarboxylative alkynylation of $\alpha$ -keto acids

A 25 mL Schlenk tube equipped with a magnetic stirring bar was charged with benzoylformic acid **1a** (0.2 mmol), 1-[(triisopropylsilyl)ethynyl]-1,2-benziodoxol-3(1H)-one **2a** (1.5 equiv), and  $K_2S_2O_8$  (0.6 equiv). The tube was evacuated and backfilled with argon (this operation was repeated three times). CH<sub>3</sub>CN (1 mL) and H<sub>2</sub>O (1 mL) were added by syringe under a flow of argon. The tube was sealed and the mixture was allowed to stir at 50 °C for 12 h. After completion of the reaction, the mixture was cooled to room temperature. Then the mixture was diluted with EtOAc, and washed with water. The organic phase was separated and the aqueous phase was extracted with EtOAc in three times. The combined organic phase was dried over anhydrous  $MgSO_4$  and the solvent was removed under vacuo. The residue was purified by column chromatography on silica gel to give the desired product **3a**.

#### Characterization data for the products

**1-phenyl-3-(triisopropylsilyl)prop-2-yn-1-one** (**3a**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (52.8 mg, 92% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.19 – 8.15 (m, 2H), 7.60 (t, *J* = 6.8 Hz, 1H), 7.48 (t, *J* = 7.8 Hz, 2H), 1.25 – 1.14 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  177.5, 136.8, 134.0, 129.5, 128.6, 103.1, 98.0, 18.6, 11.1.

**1-(p-tolyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3b**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (52.3 mg, 87% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.06 (d, *J* = 7.8 Hz, 2H), 7.27 (d, *J* = 7.9 Hz, 2H), 2.42 (s, 3H), 1.23 – 1.05 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  177.2, 145.1, 134.5, 129.7, 129.3, 103.2, 97.3, 21.8, 18.6, 11.1.

**1-(4-methoxyphenyl)-3-(triisopropylsilyl)prop-2-yn-1-one (3c)**:<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a light yellow oil (55.8 mg, 88% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.13 (d, J = 8.6 Hz, 2H), 6.94 (d, J = 8.6 Hz, 2H), 3.87 (s, 3H), 1.22 – 1.08 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 176.2, 164.4, 131.9, 130.2, 113.8, 103.2, 96.8, 55.6, 18.6, 11.1.

**1-([1,1'-biphenyl]-4-yl)-3-(triisopropylsilyl)prop-2-yn-1-one (3d)**: The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a light yellow oil

(66.1 mg, 91% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.26 – 8.21 (m, 2H), 7.71 (d, J = 8.1 Hz, 2H), 7.64 (d, J = 7.4 Hz, 2H), 7.47 (t, J = 7.5 Hz, 2H), 7.40 (t, J = 7.3 Hz, 1H), 1.26 – 1.11 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  177.1, 146.7, 139.7, 135.7, 130.1, 129.0, 128.4, 127.3, 127.2, 103.2, 98.0, 18.6, 11.2. HRMS calcd for C<sub>24</sub>H<sub>31</sub>OSi [M+H]<sup>+</sup> 363.2144; found: 363.2140.

**1-(4-ethylphenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3e**): The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (56.7 mg, 90% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.09 (d, *J* = 8.1 Hz, 2H), 7.30 (d, *J* = 8.0 Hz, 2H), 2.71 (q, *J* = 7.6 Hz, 2H), 1.26 (t, *J* = 7.6 Hz, 3H), 1.23 – 1.08 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  177.2, 151.3, 134.7, 129.8, 128.1, 103.2, 97.3, 29.1, 18.6, 15.1, 11.1. HRMS calcd for C<sub>20</sub>H<sub>31</sub>OSi [M+H]<sup>+</sup> 315.2144; found: 315.2142.

**1-(4-fluorophenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3f**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (53.6 mg, 88% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.18 (dd, J = 8.3, 5.7 Hz, 2H), 7.15 (t, J = 8.5 Hz, 2H), 1.25 – 1.01 (m, 21H).<sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  175.9, 166.4 (d, J = 256.5 Hz), 133.3 (d, J = 2.8 Hz), 132.2 (dd, J = 9.6, 3.3 Hz), 115.8 (d, J = 22.2 Hz), 102.7, 98.4, 18.6, 11.1. <sup>19</sup>F NMR (564 MHz, CDCl<sub>3</sub>)  $\delta$  -103.27 – -103.39 (m).

**1-(4-chlorophenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3g**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (53.3 mg, 83% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.09 (d, *J* = 8.4 Hz, 2H), 7.45 (d, *J* = 8.5 Hz, 2H), 1.25 - 1.01 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  176.1, 140.7, 135.2, 130.8, 129.0, 102.7, 98.8, 18.6, 11.1.

#### The Journal of Organic Chemistry

**1-(4-bromophenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3h**):<sup>17</sup>The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (56.3 mg, 77% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.01 (d, *J* = 8.2 Hz, 2H), 7.63 (d, *J* = 8.2 Hz, 2H), 1.25 – 1.01 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  176.3, 135.6, 131.9, 130.9, 129.5, 102.6, 98.9, 18.6, 11.1.

**1-(4-iodophenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3i**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (60.9 mg, 74% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.86 (s, 4H), 1.21 – 1.08 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  176.7, 138.0, 136.1, 130.7, 102.6, 102.5, 98.9, 18.6, 11.1.

**1-(4-(trifluoromethyl)phenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3j**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a light yellow oil (26.2 mg, 37% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.27 (d, *J* = 8.1 Hz, 2H), 7.76 (d, *J* = 8.2 Hz, 2H), 1.25 – 1.08 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  176.2, 139.2, 135.2 (d, *J* = 32.8 Hz), 129.8 (d, *J* = 2.4 Hz), 125.7 (q, *J* = 3.6 Hz), 123.5 (d, *J* = 272.8 Hz), 102.5, 99.9, 18.6, 11.1. <sup>19</sup>F NMR (564 MHz, CDCl<sub>3</sub>)  $\delta$  -63.22 (s).

**1-(o-tolyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3k**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (48.1 mg, 80% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.28 (d, *J* = 7.8 Hz, 1H), 7.43 (t, *J* = 7.5 Hz, 1H), 7.32 (t, *J* = 7.6 Hz, 1H), 7.23 (s, 1H), 2.63 (s, 3H), 1.22 – 1.05 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  179.2, 140.6, 135.4, 133.4, 132.9, 132.1, 125.8, 104.5, 96.3, 22.0, 18.6, 11.1.

**1-(2-methoxyphenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**31**):<sup>21</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a colorless oil

(55.1mg, 87% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.02 (d, *J* = 7.7 Hz, 1H), 7.50 (t, *J* = 7.8 Hz, 1H), 7.02 – 6.95 (m, 2H), 3.89 (s, 3H), 1.20 – 1.04 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 176.4, 159.7, 134.8, 132.9, 126.7, 120.1, 112.0, 105.1, 96.0, 55.7, 18.5, 11.1.

**1-(2-fluorophenyl)-3-(triisopropylsilyl)prop-2-yn-1-one (3m)**:<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (35.9 mg, 59% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.06 (dd, J = 10.8, 4.3 Hz, 1H), 7.55 (dd, J = 13.1, 7.2 Hz, 1H), 7.23 (d, J = 7.4 Hz, 1H), 7.14 (dd, J = 10.4, 8.8 Hz, 1H), 1.23 – 1.07 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  173.8, 162.1 (d, J = 262.5 Hz), 135.5 (d, J = 9.2 Hz), 132.0, 125.5 (d, J = 7.8 Hz), 124.1 (d, J = 4.0 Hz), 117.1 (d, J = 21.8 Hz), 104.2, 98.2 (d, J = 3.0 Hz), 18.5, 11.1. <sup>19</sup>F NMR (564 MHz, CDCl<sub>3</sub>)  $\delta$  -110.86 – -111.15 (m).

**1-(2-chlorophenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3n**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (44.9 mg, 70% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.06 (t, *J* = 6.7 Hz, 1H), 7.45 (d, *J* = 4.0 Hz, 2H), 7.36 (dt, *J* = 8.3, 4.3 Hz, 1H), 1.21 – 1.07 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  176.2, 135.5, 133.6, 133.3, 132.8, 131.5, 126.7, 104.0, 99.1, 18.5, 11.1.

**1-(2-hydroxyphenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3o**):<sup>8b</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a light yellow oil (17.6 mg, 29% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 11.67 (s, 1H), 8.07 (d, J = 8.0 Hz, 1H), 7.50 (t, J = 7.8 Hz, 1H), 6.96 (dd, J = 13.8, 7.8 Hz, 2H), 1.26 – 1.07 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 181.7, 162.9, 137.1, 133.0, 120.7, 119.4, 118.1, 101.6, 101.5, 18.6, 11.1.

**1-(m-tolyl)-3-(triisopropylsilyl)prop-2-yn-1-one (3p)**: The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (38.5 mg,

64% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.00 – 7.96 (m, 2H), 7.41 (d, *J* = 7.5 Hz, 1H), 7.36 (t, *J* = 7.6 Hz, 1H), 2.41 (s, 3H), 1.24 – 1.09 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 177.7, 138.4, 136.8, 134.8, 130.0, 128.5, 126.9, 103.2, 97.7, 21.2, 18.6, 11.1. HRMS calcd for C<sub>19</sub>H<sub>29</sub>OSi [M+H]<sup>+</sup> 301.1988; found:301.1985. **1-(3-methoxyphenyl)-3-(triisopropylsilyl)prop-2-yn-1-one (3q)**:<sup>17</sup> The product was isolated by

T-(3-methoxypnenyl)-3-(trusopropylshyl)prop-2-yn-1-one (3q): The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a light yellow oil (34.8 mg, 55% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.79 (d, J = 7.5 Hz, 1H), 7.66 (s, 1H), 7.38 (t, J = 7.9 Hz, 1H), 7.15 (dd, J = 8.2, 2.5 Hz, 1H), 3.85 (s, 3H), 1.23 – 1.09 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  177.2, 159.7, 138.1, 129.6, 122.6, 121.3, 112.7, 103.1, 97.8, 55.4, 18.6, 11.1.

**1-(3-fluorophenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3r**): The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (38.4 mg, 63% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.96 (d, *J* = 7.7 Hz, 1H), 7.82 (d, *J* = 9.2 Hz, 1H), 7.46 (td, *J* = 7.9, 5.6 Hz, 1H), 7.30 (td, *J* = 8.2, 2.4 Hz, 1H), 1.23 – 1.08 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  176.1 (d, J = 2.6 Hz), 162.7 (d, J = 248.4 Hz), 138.8 (d, J = 6.6 Hz), 130.3 (d, J = 7.6 Hz), 125.4 (d, J = 2.9 Hz), 121.1 (d, J = 21.6 Hz), 115.9 (d, J = 22.8 Hz), 102.6, 99.0, 18.5, 11.1. <sup>19</sup>F NMR (564 MHz, CDCl<sub>3</sub>)  $\delta$  -111.75 – -111.85 (m). HRMS calcd for C<sub>18</sub>H<sub>26</sub>FOSi 305.1737; found: 305.1739.

**1-(3-chlorophenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3s**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (48.1 mg, 75% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.14 (s, 1H), 8.04 (d, *J* = 7.6 Hz, 1H), 7.57 (dd, *J* = 7.9, 0.7 Hz, 1H), 7.42 (t, *J* = 7.9 Hz, 1H), 1.25 – 1.09 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  176.0, 138.2, 134.9, 133.9, 129.9, 129.6, 127.5, 102.5, 99.3, 18.5, 11.1.

**1-(3-bromophenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3t**):<sup>17</sup>The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a yellow oil (36.5 mg, 50% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.31 (s, 1H), 8.08 (d, *J* = 7.7 Hz, 1H), 7.72 (d, *J* = 7.9 Hz, 1H), 7.36 (t, *J* = 7.9 Hz, 1H), 1.23 – 1.09 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  175.9, 138.4, 136.8, 132.6, 130.2, 127.9, 122.8, 102.5, 99.5, 18.6, 11.1.

**1-(2,4-dimethylphenyl)-3-(triisopropylsilyl)prop-2-yn-1-one (3u**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a light yellow oil (44.7 mg, 71% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.19 (d, J = 8.0 Hz, 1H), 7.12 (d, J = 7.9 Hz, 1H), 7.05 (s, 1H), 2.60 (s, 3H), 2.36 (s, 3H), 1.22 – 1.05 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 178.8, 143.8, 140.8, 133.9, 132.94, 132.93, 126.5, 104.6, 95.5, 22.0, 21.5, 18.6, 11.1.

**1-(3,4-dimethoxyphenyl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3v**): The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:30) as a colorless oil (51.3 mg, 74% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.87 (dd, *J* = 8.3, 1.7 Hz, 1H), 7.64 (d, *J* = 1.5 Hz, 1H), 6.93 (d, *J* = 8.4 Hz, 1H), 3.95 (s, 3H), 3.92 (s, 3H), 1.21 – 1.08 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  176.1, 154.2, 148.9, 130.3, 125.2, 110.6, 110.2, 103.1, 96.8, 56.1, 56.0, 18.6, 11.1. HRMS calcd for C<sub>20</sub>H<sub>31</sub>O<sub>3</sub>Si [M+H]<sup>+</sup> 347.2042; found:347.2041.

**1-(naphthalen-2-yl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3w**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a yellow oil (50.5 mg, 75% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.79 (s, 1H), 8.15 (dd, *J* = 8.6, 1.3 Hz, 1H), 7.95 (d, *J* = 8.1 Hz, 1H), 7.89 (t, *J* = 7.8 Hz, 2H), 7.62 (t, *J* = 7.5 Hz, 1H), 7.56 (t, *J* = 7.5 Hz, 1H), 1.28 – 1.14 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 177.4, 136.1, 134.4, 133.0, 132.4, 129.8, 129.0, 128.4, 127.9, 126.9, 123.8, 103.2, 97.9, 18.6, 11.2.

 **1-(naphthalen-1-yl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3x**):<sup>17</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a yellow oil (45.8 mg, 68% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 9.20 (d, *J* = 8.7 Hz, 1H), 8.62 (d, *J* = 7.3 Hz, 1H), 8.07 (d, *J* = 8.1 Hz, 1H), 7.89 (d, *J* = 8.1 Hz, 1H), 7.67 – 7.63 (m, 1H), 7.56 (td, *J* = 7.4, 4.1 Hz, 2H), 1.24 – 1.11 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 179.2, 135.0, 134.6, 133.8, 132.8, 130.7, 128.9, 128.5, 126.7, 126.0, 124.4, 104.6, 96.2, 18.6, 11.2.

**1-(benzo[d][1,3]dioxol-5-yl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3y**): The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a colorless oil (54.2 mg, 82% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.83 (dd, *J* = 8.1, 1.4 Hz, 1H), 7.56 (d, *J* = 1.3 Hz, 1H), 6.87 (d, *J* = 8.2 Hz, 1H), 6.06 (s, 2H), 1.25 – 1.03 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  175.6, 152.8, 148.2, 132.0, 127.1, 108.3, 108.0, 103.0, 102.1, 97.1, 18.6, 11.1. HRMS calcd for C<sub>19</sub>H<sub>27</sub>O<sub>3</sub>Si [M+H]<sup>+</sup> 331.1729; found: 331.1728.

**1-(thiophen-3-yl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3z**):<sup>8d</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a colorless oil (18.2 mg, 31% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.28 – 8.25 (m, 1H), 7.61 (t, *J* = 4.5 Hz, 1H), 7.32 (dd, *J* = 4.9, 3.1 Hz, 1H), 1.21 – 1.08 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  170.9, 143.0, 135.4, 126.77, 126.75, 103.6, 95.9, 18.6, 11.1.

**1-(benzo[b]thiophen-2-yl)-3-(triisopropylsilyl)prop-2-yn-1-one** (**3aa**):<sup>8d</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a yellow oil (24.7 mg, 36% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.18 (s, 1H), 7.92 (d, *J* = 8.0 Hz, 1H), 7.86 (d, *J* = 8.2 Hz, 1H), 7.48 (t, *J* = 7.6 Hz, 1H), 7.42 (t, *J* = 7.6 Hz, 1H), 1.24 – 1.11 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  170.7, 144.3, 143.1, 138.7, 132.6, 128.0, 126.4, 125.2, 123.1,

102.4, 97.6, 18.6, 11.1.

1-(1H-indol-3-yl)-3-(triisopropylsilyl)prop-2-yn-1-one (3ab): The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:5) as a light yellow solid (21.5 mg, 33% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  9.62 (s, 1H), 8.41 – 8.38 (m, 1H), 8.11 (d, *J* = 2.9 Hz, 1H), 7.48 – 7.45 (m, 1H), 7.29 (m, 2H), 1.18 – 1.07 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  171.9, 136.8, 135.8, 125.0, 124.2, 123.1, 122.2, 119.8, 111.9, 104.2, 92.1, 18.6, 11.1. HRMS calcd for C<sub>20</sub>H<sub>28</sub>NOSi [M+H]<sup>+</sup> 326.1940; found: 326.1943.

**5-phenyl-1-(triisopropylsilyl)pent-1-yn-3-one (3ad)**.<sup>8d</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a colorless oil (43.4 mg, 69% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.28 (t, J = 7.5 Hz, 2H), 7.20 (t, J = 7.3 Hz, 3H), 3.01 (t, J = 7.7 Hz, 2H), 2.90 (t, J = 7.7 Hz, 2H), 1.15 – 1.02 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  186.5, 140.2, 128.5, 128.3, 126.3, 104.0, 96.1, 47.2, 30.1, 18.5, 11.0.

**4-(triisopropylsilyl)but-3-yn-2-one** (**3ae**):<sup>17</sup>The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a colorless oil (21.5 mg, 48% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  2.34 (d, *J* = 4.6 Hz, 3H), 1.14 – 1.03 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  184.2, 104.7, 95.2, 32.8, 18.4, 11.0.

**1-(triisopropylsilyl)pent-1-yn-3-one (3af)**: The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a colorless oil (24.8 mg, 52% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  2.58 (q, *J* = 7.4 Hz, 2H), 1.15 (dd, *J* = 9.8, 5.0 Hz, 3H), 1.14 – 1.03 (m, 21H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  188.3, 104.0, 95.5, 38.9, 18.5, 11.0, 8.2. HRMS calcd for C<sub>14</sub>H<sub>27</sub>OSi [M+H]<sup>+</sup> 239.1831; found: 239.1829.

1-cyclopropyl-3-(triisopropylsilyl)prop-2-yn-1-one (3ag):<sup>8d</sup> The product was isolated by

#### The Journal of Organic Chemistry

column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a colorless oil (36.1 mg, 72% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  2.02 (ddd, J = 12.6, 8.1, 4.6 Hz, 1H), 1.26 (td, J = 7.3, 3.5 Hz, 2H), 1.14 – 1.00 (m, 23H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  188.1, 101.9, 94.9, 24.6, 18.5, 11.0, 10.8.

**1-phenylnon-2-yn-1-one** (**4a**):<sup>22</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a light yellow oil (22.3 mg, 52% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.13 (d, *J* = 7.7 Hz, 2H), 7.59 (t, *J* = 7.3 Hz, 1H), 7.47 (t, *J* = 7.7 Hz, 2H), 2.49 (t, *J* = 7.1 Hz, 2H), 1.70 – 1.64 (m, 2H), 1.47 (dt, *J* = 14.7, 7.3 Hz, 2H), 1.35 – 1.30 (m, 4H), 0.93 – 0.86 (m, 3H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 178.3, 136.9, 133.8, 129.5, 128.5, 96.9, 79.7, 31.2, 28.6, 27.8, 22.5, 19.2, 14.0.

**1-phenyl-3-(triethylsilyl)prop-2-yn-1-one** (**4b**):<sup>23</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a light yellow oil (24.1 mg, 49% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.15 (d, *J* = 7.5 Hz, 2H), 7.60 (t, *J* = 7.4 Hz, 1H), 7.48 (t, *J* = 7.6 Hz, 2H), 1.07 (t, *J* = 7.9 Hz, 9H), 0.74 (q, *J* = 7.9 Hz, 6H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  177.6, 136.6, 134.1, 129.6, 128.6, 102.1, 98.8, 7.4, 4.0.

**3-(tert-butyldimethylsilyl)-1-phenylprop-2-yn-1-one** (**4c**):<sup>24</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a colorless oil (35.2 mg, 72% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.14 (d, *J* = 7.7 Hz, 2H), 7.60 (t, *J* = 7.3 Hz, 1H), 7.48 (t, *J* = 7.7 Hz, 2H), 1.02 (s, 9H), 0.25 (s, 6H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 177.6, 136.6, 134.1, 129.6, 128.6, 101.6, 99.4, 26.1, 16.7, -5.1.

**3-(tert-butyldiphenylsilyl)-1-phenylprop-2-yn-1-one** (**4d**): The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a light yellow oil (47.2

mg, 64% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.23 (d, J = 7.4 Hz, 2H), 7.83 (d, J = 6.8 Hz, 4H), 7.63 (t, J = 7.4 Hz, 1H), 7.50 (t, J = 7.7 Hz, 2H), 7.43 (dt, J = 22.6, 7.3 Hz, 6H), 1.19 (s, 9H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  177.3, 136.7, 135.6, 134.3, 131.5, 130.1, 129.6, 128.7, 128.0, 104.4, 96.0, 27.1, 18.9. HRMS calcd for C<sub>25</sub>H<sub>25</sub>OSi [M+H]<sup>+</sup> 369.1675; found:369.1673.

**1,3-diphenylprop-2-yn-1-one** (**4e**):<sup>24</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:100) as a light yellow oil (21.1 mg, 51% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.22 (d, *J* = 7.8 Hz, 2H), 7.68 (d, *J* = 7.5 Hz, 2H), 7.63 (t, *J* = 7.4 Hz, 1H), 7.50 (dt, *J* = 20.2, 7.6 Hz, 3H), 7.42 (t, *J* = 7.6 Hz, 2H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  178.0, 136.9, 134.1, 133.1, 130.8, 129.6, 128.7, 128.6, 120.1, 93.1, 86.9.

**1-phenyl-3-(p-tolyl)prop-2-yn-1-one** (**4f**):<sup>10</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a white solid (35.7 mg, 81% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.22 (d, *J* = 7.7 Hz, 2H), 7.62 (t, *J* = 7.3 Hz, 1H), 7.58 (d, *J* = 8.0 Hz, 2H), 7.51 (t, *J* = 7.7 Hz, 2H), 7.22 (d, *J* = 7.9 Hz, 2H), 2.40 (s, 3H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  178.1, 141.5, 137.0, 134.0, 133.1, 129.53, 129.47, 128.6, 117.0, 93.8, 86.8, 21.8.

**3-(3-methoxyphenyl)-1-phenylprop-2-yn-1-one (4g)**:<sup>9</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a yellow oil (11.8 mg, 25% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.21 (d, *J* = 8.0 Hz, 2H), 7.63 (t, *J* = 7.4 Hz, 1H), 7.52 (t, *J* = 7.7 Hz, 2H), 7.35 – 7.26 (m, 2H), 7.19 (s, 1H), 7.05 – 7.00 (m, 1H), 3.84 (s, 3H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 178.0, 159.4, 136.8, 134.1, 129.8, 129.6, 128.6, 125.6, 121.0, 117.56, 117.55, 93.0, 86.5, 55.4.

**3-(4-chlorophenyl)-1-phenylprop-2-yn-1-one** (**4h**):<sup>9</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a light yellow solid (27.4)

#### The Journal of Organic Chemistry

mg, 57% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.20 (d, J = 7.6 Hz, 2H), 7.62 (dd, J = 17.5, 7.9 Hz, 3H), 7.51 (t, J = 7.7 Hz, 2H), 7.40 (d, J = 8.2 Hz, 2H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  177.8, 137.2, 136.7, 134.24, 134.22, 129.6, 129.2, 128.7, 118.6, 91.6, 87.6.

**2-(3-oxo-3-phenylprop-1-yn-1-yl)benzonitrile** (**4i**):<sup>9</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:20) as a white solid (12.5 mg, 27% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.33 (d, *J* = 7.9 Hz, 2H), 7.81 (d, *J* = 7.8 Hz, 1H), 7.76 (d, *J* = 7.7 Hz, 1H), 7.69 – 7.62 (m, 2H), 7.59 (t, *J* = 7.7 Hz, 1H), 7.54 (t, *J* = 7.6 Hz, 2H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 177.5, 136.3, 134.6, 134.4, 133.0, 132.7, 130.8, 129.9, 128.8, 124.1, 117.1, 116.3, 91.2, 86.9.

**3-(4-acetylphenyl)-1-phenylprop-2-yn-1-one** (**4j**):<sup>9</sup> The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:20) as a light yellow solid (21.4 mg, 43% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.20 (d, *J* = 7.3 Hz, 2H), 7.99 (d, *J* = 8.3 Hz, 2H), 7.76 (d, *J* = 8.3 Hz, 2H), 7.64 (t, *J* = 7.4 Hz, 1H), 7.52 (t, *J* = 7.7 Hz, 2H), 2.63 (s, 3H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 197.1, 177.7, 138.0, 136.6, 134.4, 133.1, 129.6, 128.7, 128.4, 124.7, 91.1, 88.7, 26.7.

**4-(3-oxo-3-phenylprop-1-yn-1-yl)benzaldehyde** (**4k**):<sup>9</sup>The product was isolated by column chromatography on silica gel (ethyl acetate: petroleum ether = 1:20) as a light yellow solid (21.6 mg, 46% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 10.06 (s, 1H), 8.20 (d, *J* = 7.5 Hz, 2H), 7.93 (d, *J* = 8.0 Hz, 2H), 7.83 (d, *J* = 8.1 Hz, 2H), 7.65 (t, *J* = 7.4 Hz, 1H), 7.53 (t, *J* = 7.7 Hz, 2H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 191.1, 177.6, 137.1, 136.5, 134.5, 133.4, 129.6, 128.7, 126.1, 90.7, 89.1.

3-([1,1'-biphenyl]-4-yl)-1-phenylprop-2-yn-1-one (4l):9 The product was isolated by column

chromatography on silica gel (ethyl acetate: petroleum ether = 1:50) as a light yellow solid (23.2 mg, 41% yield); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.24 (d, *J* = 7.6 Hz, 2H), 7.76 (d, *J* = 8.2 Hz, 2H), 7.67 – 7.60 (m, 5H), 7.53 (t, *J* = 7.7 Hz, 2H), 7.47 (t, *J* = 7.6 Hz, 2H), 7.40 (t, *J* = 7.4 Hz, 1H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 178.0, 143.6, 139.8, 136.9, 134.1, 133.6, 129.6, 129.0, 128.6, 128.2, 127.3, 127.1, 118.8, 93.2, 87.6.

#### **AUTHOR INFORMATION**

#### Corresponding Authors

E-mail: hjxu@hfut.edu.cn; fengyisi2008@126.com

#### ACKNOWLEDGE

We gratefully acknowledge financial support from the National Natural Science Foundation of China (Nos. 21472033, 21371044, 21272050) and the Program for New Century Excellent Talents in University of the Chinese Ministry of Education (NCET-11-0627).

#### ASSOCIATED CONTENT

#### **Supporting Information**

<sup>1</sup>H NMR, <sup>13</sup>C NMR and <sup>19</sup>F NMR spectra of the products. This material is available free of charge via the Internet at <u>http://pubs.acs.org/</u>.

#### REFERENCES

1 (a) Van den Hoven, B. G.; Ali, B. E.; Alper, H. J. Org. Chem. 2000, 65, 4131. (b) Jeevanandam,

A.; Narkunan, K.; Ling, Y.-C. J. Org. Chem. 2001, 66, 6014. (c) Xiong, X.; Bagley, M. C.;

Chapaneri, K. Tetrahedron Lett. 2004, 45, 6121. (d) Liang, J. T.; Mani, N. S.; Jones, T. K. J. Org.

Chem. 2007, 72, 8243. (e) Kirkham, J. D.; Edeson, S. J.; Stokes, S.; Harrity, J. P. A. Org. Lett.

2012, 14, 5354.

2.(a) Karpov, A. S.; Müller, T. J. J. *Org. Lett.* **2003**, *5*, 3451. (b) Chen, L.; Li, C.-J. *Org. Lett.* **2004**, *6*, 3151.

3. (a) Mohamed Ahmed, M. S.; Mori, A. Org. Lett. 2003, 5, 3057. (b) Karpov, A. S.; Merkul, E.;

Rominger, F.; Müller, T. J. J. Angew. Chem., Int. Ed. 2005, 44, 6951. (c) Fusano, A.; Fukuyama,

T.; Nishitani, S.; Inouye, T.; Ryu, I. Org. Lett, 2010, 12, 2410.

4. Wu, X.-F.; Sundararaju, B.; Neumann, H.; Dixneuf, P. H.; Beller, M. *Chem. Eur. J.* **2011**, *17*, 106.

5. Wu, X.-F.; Neumann, H.; Beller, M. Angew. Chem. Int. Ed. 2011, 50, 11142.

6. Yu, Y.; Yang, W.; Pflästerer, D.; Hashmi, A. S. K. Angew. Chem. Int. Ed. 2014, 53, 1144.

7. Wang, Z.; Li, L.; Huang, Y. J. Am. Chem. Soc. 2014, 136, 12233.

8. (a) Yuan, J.; Wang, J.; Zhang, G.; Liu, C.; Qi, X.; Lan, Y.; Miller, J. T.; Kropf, A. J.; Bunel, E.

E.; Lei, A. Chem. Commun. 2015, 51, 576. (b) Wang, H.; Xie, F.; Qi, Z.; Li, X. Org. Lett. 2015,

17, 920. (c) Ai, W.; Wu, Y.; Tang, H.; Yang, X.; Yang, Y.; Li, Y.; Zhou, B. Chem. Commun. 2015,

51, 7871. (d) Liu, X.; Yu, L.; Luo, M.; Zhu, J.; Wei, W. Chem. Eur. J. 2015, 21, 8745.

9. Huang, H.; Zhang, G.; Chen, Y. Angew. Chem. Int. Ed. 2015, 54, 7872.

10. Tan, H.; Li, H.; Ji, W.; Wang, L. Angew. Chem. Int. Ed. 2015, 54, 8374.

11. Myers, A. G.; Tanaka, D.; Mannion, M. R. J. Am. Chem. Soc. 2002, 124, 11250.

12. Forgione, P.; Brochu, M.-C.; St-Onge, M.; Thesen, K. H.; Bailey, M. D.; Bilodeau, F. J. Am. Chem. Soc. 2006, 128, 11350.

13. (a) Goossen, L. J.; Rodriguez, N.; Melzer, B.; Linder, C.; Deng, G. J.; Levy, L. M. J. Am.

Chem. Soc. 2007, 129, 4824. (b) Goossen, L. J.; Rodriguez, N.; Linder, C. J. Am. Chem. Soc. 2008,

130, 15248. (c) Goossen, L. J.; Rodriguez, N.; Lange, P. P.; Linder, C. Angew. Chem. Int. Ed.
2010, 49, 1111.

14. (a) Shang, R.; Fu, Y.; Wang, Y.; Xu, Q.; Yu, H.-Z.; Liu, L. Angew. Chem. Int. Ed. 2009, 48,

9350. (b) Shang, R.; Yang, Z.-W.; Wang, Y.; Zhang, S.-L.; Liu, L. J. Am. Chem. Soc. 2010, 132,

14391. (c) Shang, R.; Ji, D.-S.; Chu, L.; Fu, Y.; Liu, L. Angew. Chem. Int. Ed. 2011, 50, 4470. (d)

Shang, R.; Xu, Q.; Jiang, Y.-Y.; Wang, Y.; Liu, L. Org. Lett. 2010, 12, 1000. (e) Shang, R.; Huang, Z.; Chu, L.; Fu, Y.; Liu, L. Org. Lett. 2011, 13, 4240.

15. Some examples: (a) Burger, E. C.; Tunge, J. A. J. Am. Chem. Soc. 2006, 128, 10002. (b) Becht,

J.-M.; Catala, C.; Drian, C. L.; Wagner, A. Org. Lett, 2007, 9, 1781. (c) Voutchkova, A.; Coplin,

A.; Leadbeater, N. E.; Crabtree, R. H. Chem. Commun. 2008, 6312. (d) Hu, P.; Zhang, M.; Xie, X.;

Su, W. Angew. Chem. Int. Ed. 2012, 51, 227. (e) Liu, J.; Liu, Q.; Yi, H.; Qin, C.; Bai, R.; Qi, X.;

Lan, Y.; Lei, A. Angew. Chem. Int. Ed. 2014, 53, 502.

3054.

16. Our previous work: (a) Feng, Y.-S.; Wu, W.; Xu, Z.-Q.; Li, Y.; Li, M.; Xu, H.-J. *Tetrahedron*.
2012, *68*, 2113. (b) Li, Y.; Chen, H.-H.; Wang, C.-F.; Xu, X.-L.; Feng, Y.-S. *Tetrahedron Lett*.
2012, *53*, 5796. (c) Feng, Y.-S.; Xu, Z.-Q.; Mao, L.; Zhang, F.-F.; Xu, H.-J. *Org. Lett*. 2013, *15*, 1472. (d) Wang, P.-F.; Wang, X.-Q.; Dai, J.-J.; Feng, Y.-S.; Xu, H.-J. *Org. Lett*. 2014, *16*, 4586.
17. During the preparation of our manuscript, a similar work was reported by *Org. Lett*. 2015, *17*,

(a) Anderson, J. M.; Kochi, J. K. J. Am. Chem. Soc. 1970, 92, 1651. (b) Liu, X.; Wang, Z.;
 Cheng, X.; Li, C. J. Am. Chem. Soc. 2012, 134, 14330.

Wadhwa, K.; Yang, C.; West, P. R.; Deming, K, C.; Chemburkar, S. R.; Reddy, R. E. Synth.
 *Commun.* 2008, 38, 4434.

1
2
3
4
5
5
6
7
8
à
10
10
11
12
13
14
15
10
16
17
18
19
20
20
21
22
23
24
25
20
26
27
28
29
30
30
31
32
33
34
35
30
36
37
38
39
40
40
41
42
43
44
45
16
40
47
48
49
50
50 51
51
52
53
54
55
56
50
5/
58
59
60
00

- Huang, H.; Zhang, G.; Gong, L.; Zhang, S.; Chen, Y. J. Am. Chem. Soc. 2014, 136, 2280.
- 21. Feng, L.; Kumar, D.; Kerwin, S. M. J. Org. Chem. 2003, 68, 2234.
- 22. Reddy, K. R.; Suresh, M.; Kantam, M. L.; Bhargava, S, K.; Srinivasu, P. Ind. Eng. Chem. Res.

**2014,** *53*, 18630.

- 23. Shintani, R.; Okamoto, K.; Hayashi, T. J. Am. Chem. Soc. 2005, 127, 2872.
- 24. Cox, R. J.; Ritson, D. J.; Dane, T. A.; Berge, J.; Charmant, J. P. H; Kantacha, A. Chem.

Commun. 2005, 1037.