

Copper-catalyzed Conjugate Additions of Alkylboranes to Aryl α,β -Unsaturated Ketones

Hirohisa Ohmiya,* Yoshinori Shido, Mika Yoshida, and Masaya Sawamura*

Department of Chemistry, Faculty of Science, Hokkaido University, Sapporo, Hokkaido 060-0810

(Received April 13, 2011; CL-110311; E-mail: ohmiya@sci.hokudai.ac.jp, sawamura@sci.hokudai.ac.jp)


Conjugate addition of alkylboron compounds (alkyl-9-BBN) to aryl α,β -unsaturated ketones proceeded in the presence of a catalytic amount (10 mol %) of $[(\text{IPr})\text{CuCl}]$ and $t\text{-BuOK}$. The alkylboranes are available through alkene hydroboration, and thus the overall process represents a reductive conjugate addition of alkenes to enone derivatives. A variety of functional groups are tolerated in both the alkenes and the α,β -unsaturated ketones.

Transition-metal (Rh, Pd, Ni, etc.)-catalyzed conjugate additions of organoboron compounds to α,β -unsaturated carbonyl compounds are useful carbon–carbon bond formation methods because of their broad substrate scope, functional group compatibility and applicability to asymmetric reactions.^{1–4} Unfortunately however, the organoboron reagents that are generally usable for these methods are limited to aryl-, alkenyl-, and allylboron compounds, and the reactions of alkylboron derivatives are relatively unexplored.^{2–7}

In this context, we reported recently that conjugate addition of alkylboron compounds (alkyl-9-BBN) to imidazol-2-yl α,β -unsaturated ketones proceeded in the presence of a catalytic amount (10 mol %) of a Cu/*N*-heterocyclic carbene (NHC) and $t\text{-BuOK}$.^{8,9} Here we report that the copper-catalyzed conjugate addition of alkylboranes has been expanded to the reaction with aryl α,β -unsaturated ketones.^{10–12} The wide and easy availability of alkylboranes via the established alkene hydroboration is an attractive feature of this transformation and thus the overall process represents a reductive conjugate addition of alkenes to enone derivatives. A variety of functional groups are tolerated in both the alkenes and the α,β -unsaturated ketones.

Specifically, alkylborane **2a** in THF solution was prepared via hydroboration of styrene (**1a**) with 9-borabicyclo[3.3.1]nonane dimer [9-BBN- H]₂ (**1a**/B 1.05:1) at 60 °C (Scheme 1). Subsequently, the resulting THF solution of **2a** (0.3 mmol) was added to a THF solution of $[(\text{IPr})\text{CuCl}]$ ¹³ (10 mol %) and $t\text{-BuOK}$ (10 mol %). Chalcone (**3a**) (0.2 mmol) was then added to the mixture, which was heated at 80 °C for 8 h. After hydrolytic workup, the conjugate addition product **4aa** was obtained in 83% isolated yield.¹⁴

Several observations concerning the optimum reaction conditions are to be noted (Scheme 1). Unlike the conjugate addition of alkylboranes to imidazol-2-yl α,β -unsaturated ketones, IPr ligand was effective for the present protocol. While SIPr was as effective as IPr (80% yield), other NHC ligands such as IMes and ICy resulted in no reactions under otherwise identical conditions.¹³ The reaction without a ligand afforded a complex mixture with no addition product. The use of $\text{Cu}(\text{O}t\text{-Bu})/\text{IPr}$ instead of $[(\text{IPr})\text{CuCl}]/t\text{-BuOK}$ was also effective to produce **3aa** in 85%, suggesting that KCl present in the optimal conditions is not essential for the catalysis. No reaction occurred when $[(\text{IPr})\text{CuCl}]/t\text{-BuOK}$ was omitted.¹⁵ Alkyl α,β -unsaturat-



Scheme 1. Cu-catalyzed conjugate addition of alkylborane.

ed ketones did not afford the corresponding products under similar reaction conditions: the lower reactivity of the alkyl ketones would be attributed to their lower electrophilicity.¹⁶

Various alkenes **1** and aryl α,β -unsaturated ketones **3** were subjected to the reductive conjugate addition protocol (Table 1).¹⁷ Functional groups such as ester, acetal, silyl ether, methoxy, trifluoromethyl, phthalimide, bromo, and amido moieties were tolerated in the reaction (Entries 2–11). Both alkyl- and aryl substituents were compatible at the β -position in the α,β -unsaturated ketones. The substrate **3k** bearing a 2-thienyl group at the β -position underwent the conjugate addition (Entry 11). The substitution of the aromatic ring of the aryl ketone with either MeO or CF₃ groups caused only marginal and capricious effects on the reactivity (Entries 5 and 6). The 2-furyl ketone substrate **3g** served as a substrate (Entry 7).

The tolerance of the reaction toward steric demand in alkylboranes **2** and α,β -unsaturated ketones **3** is also shown in Table 1. The sterically more demanding alkylborane **2e**, which was derived from a terminal alkene **1e** with a tertiary alkyl substituent, underwent coupling with **3i** to afford the corresponding product **4ei** in good yield (Entry 9). The reaction of the β -branched alkylborane **2h**, which was prepared from α -methylstyrene **1h**, also proceeded smoothly to produce **4ha** as a mixture of diastereomers (79:21) (Entry 12). No reaction occurred with secondary alkylboranes (data not shown). A sterically more demanding γ -alkyl substituent such as a cyclohexyl group was tolerated (Entry 3).

In summary, the scope of the copper-catalyzed conjugate addition of alkylboranes (alkyl-9-BBN) has been expanded toward aryl α,β -unsaturated ketones. The alkylboranes are easily and widely available through alkene hydroboration and thus the overall process represents a reductive conjugate addition of alkenes to enone derivatives. A variety of functional groups

Table 1. Substrate scope of the Cu-catalyzed conjugate addition of alkylboranes^a

Entry	Alkene	Ketone	Product	Yield/% ^b
1	1a	3b	4ab	73
2	1b	3c	4bc	69
3	1a	3d	4ad	62 ^c
4	1c	3a	4ca	84
5	1c	3e	4ce	93 ^d
6	1c	3f	4cf	82
7	1a	3g	4ag	78
8	1d	3h	4dh	93
9	1e	3i	4ei	74
10	1f	3j	4fj	71
11	1g	3k	4gk	75 ^d
12	1h	3a	4ha	74 ^c

^aThe reaction was carried out with **3** (0.4 mmol, *E/Z* > 99:1), alkylborane **2** (Entries 1–3, 0.6 mmol; Entries 4–12, 0.48 mmol), [(IPr)CuCl] (10 mol %) and *t*-BuOK (10 mol %) in THF (0.8 mL) at 80 °C for 8 h. Alkylborane **2** was prepared in advance from **1**. ^bIsolated yield based on **3**. ^cNMR yield. ^dThe isolated product was contaminated with a trace amount of an unidentified material. ^eDiastereomer ratio 79:21.

are tolerated in both the alkenes and the α,β -unsaturated ketones.

This work was supported by Grants-in-Aid for Scientific Research (B) and for Young Scientists (B), JSPS. We thank MEXT for financial support in the form of a Global COE grant (Project No. B01: Catalysis as the Basis for Innovation in Materials Science).

This paper is in celebration of the 2010 Nobel Prize awarded to Professors Richard F. Heck, Akira Suzuki, and Ei-ichi Negishi.

References and Notes

- For reviews on transition-metal-catalyzed conjugate additions, see: a) N. Krause, A. Hoffmann-Röder, *Synthesis* **2001**, 171. b) A. Alexakis, J. E. Bäckvall, N. Krause, O. Pàmies, M. Diéguez, *Chem. Rev.* **2008**, *108*, 2796. c) S. R. Harutyunyan, T. den Hartog, K. Geurts, A. J. Minnaard, B. L. Feringa, *Chem. Rev.* **2008**, *108*, 2824. d) J. Christoffers, G. Koripelly, A. Rosiak, M. Rössle, *Synthesis* **2007**, 1279. e) C. Hawner, A. Alexakis, *Chem. Commun.* **2010**, 46, 7295.
- For selected references on Rh-catalyzed conjugate additions with aryl- and alkenylboron compounds, see: a) M. Sakai, H. Hayashi, N. Miyaura, *Organometallics* **1997**, *16*, 4229. b) Y. Takaya, M. Ogasawara, T. Hayashi, M. Sakai, N. Miyaura, K. Ueyama, N. Tokunaga, K. Yoshida, *J. Am. Chem. Soc.* **2003**, *125*, 11508. For reviews, see: d) K. Fagnou, M. Lautens, *Chem. Rev.* **2003**, *103*, 169. e) T. Hayashi, K. Yamasaki, *Chem. Rev.* **2003**, *103*, 2829.
- For selected references on Pd-catalyzed conjugate additions with aryl- and alkenylboron compounds, see: a) C. S. Cho, K. Tanabe, S. Uemura, *Tetrahedron Lett.* **1994**, *35*, 1275. b) C. S. Cho, S. Motofusa, K. Ohe, S. Uemura, S. C. Shim, *J. Org. Chem.* **1995**, *60*, 883. c) C. S. Cho, S. Motofusa, K. Ohe, S. Uemura, *Bull. Chem. Soc. Jpn.* **1996**, *69*, 2341. d) F. Gini, B. Hessen, A. J. Minnaard, *Org. Lett.* **2005**, *7*, 5309. e) P. He, Y. Lu, C.-G. Dong, Q.-S. Hu, *Org. Lett.* **2007**, *9*, 343. f) X. Lu, S. Lin, *J. Org. Chem.* **2005**, *70*, 9651. g) S. Lin, X. Lu, *Tetrahedron Lett.* **2006**, *47*, 7167. h) S. Lin, X. Lu, *Org. Lett.* **2010**, *12*, 2536. i) Y. Yamamoto, T. Nishikata, N. Miyaura, *Pure Appl. Chem.* **2008**, *80*, 807, and references therein.
- For Ni-catalyzed conjugate additions with arylboron compounds, see: E. Shirakawa, Y. Yasuhara, T. Hayashi, *Chem. Lett.* **2006**, *35*, 768.
- For conjugate additions with allylboron compounds, see: a) J. D. Sieber, S. Liu, J. P. Morken, *J. Am. Chem. Soc.* **2007**, *129*, 2214. b) J. D. Sieber, J. P. Morken, *J. Am. Chem. Soc.* **2008**, *130*, 4978. c) M. B. Shaghafi, B. L. Kohn, E. R. Jarvo, *Org. Lett.* **2008**, *10*, 4743.
- For conjugate additions with alkylboron compounds, see: a) N. Miyaura, M. Itoh, A. Suzuki, *Tetrahedron Lett.* **1976**, *17*, 255. b) K. Hirano, H. Yorimitsu, K. Oshima, *Org. Lett.* **2007**, *9*, 1541.
- For a review on transition-metal-catalyzed cross-coupling reactions using alkyl-organometallic reagents, see: R. Jana, T. P. Pathak, M. S. Sigman, *Chem. Rev.* **2011**, *111*, 1417.
- H. Ohmiya, M. Yoshida, M. Sawamura, *Org. Lett.* **2011**, *13*, 482.
- For Cu-catalyzed conjugate addition of arylboronic acid to alkynoates, see: Y. Yamamoto, N. Kirai, Y. Harada, *Chem. Commun.* **2008**, 2010.
- For Cu-catalyzed γ -selective and stereospecific allyl-alkyl and allyl-aryl couplings with organoboron compounds, see: a) H. Ohmiya, U. Yokobori, Y. Makida, M. Sawamura, *J. Am. Chem. Soc.* **2010**, *132*, 2895. b) H. Ohmiya, N.

- Yokokawa, M. Sawamura, *Org. Lett.* **2010**, *12*, 2438. c) A. M. Whittaker, R. P. Rucker, G. Lalic, *Org. Lett.* **2010**, *12*, 3216. For Cu-catalyzed carboxylations with alkylboron compounds (alkyl-9-BBN) to carbon dioxide, see: d) H. Ohmiya, M. Tanabe, M. Sawamura, *Org. Lett.* **2011**, *13*, 1086.
- 11 Knochel et al. reported the copper-mediated conjugate addition of dialkylzinc reagents, prepared by a hydroboration/boron–zinc exchange sequence, to ethyl acrylate, but the application to β -substituted α,β -unsaturated esters is underdeveloped while benzylidene malonate has successfully been used. Furthermore, the cumbersome procedure and low atom efficiency, using large excess substrate and reagents, hampers the wide application of this method. See: a) F. Langer, L. Schwink, A. Devasagayaraj, P.-Y. Chavant, P. Knochel, *J. Org. Chem.* **1996**, *61*, 8229. b) E. Hupe, M. I. Calaza, P. Knochel, *Tetrahedron Lett.* **2001**, *42*, 8829. c) E. Hupe, M. I. Calaza, P. Knochel, *Chem.—Eur. J.* **2003**, *9*, 2789. d) E. Hupe, M. I. Calaza, P. Knochel, *J. Organomet. Chem.* **2003**, *680*, 136.
- 12 For selected references on Cu-catalyzed enantioselective conjugate additions with alkylmagnesium- or alkylzinc reagents, see: a) K.-s. Lee, M. K. Brown, A. W. Hird, A. H. Hoveyda, *J. Am. Chem. Soc.* **2006**, *128*, 7182. b) D. Martin, S. Kehrli, M. d'Augustin, H. Clavier, M. Mauduit, A. Alexakis, *J. Am. Chem. Soc.* **2006**, *128*, 8416. c) S. R. Harutyunyan, F. López, W. R. Browne, A. Correa, D. Peña, R. Badorrey, A. Meetsma, A. J. Minnaard, B. L. Feringa, *J. Am. Chem. Soc.* **2006**, *128*, 9103. d) S.-Y. Wang, S.-J. Ji, T.-P. Loh, *J. Am. Chem. Soc.* **2007**, *129*, 276. See also refs. 1b and 1c.
- 13 IPr: 1,3-Bis(2,6-diisopropylphenyl)imidazol-2-ylidene; SIPr: 1,3-Bis(2,6-diisopropylphenyl)imidazolidin-2-ylidene; IMes: 1,3-Bis(2,4,6-trimethylphenyl)imidazol-2-ylidene; ICy: 1,3-Dicyclohexylimidazol-2-ylidene; For reviews on *N*-heterocyclic carbenes (NHCs), see: a) *N-Heterocyclic Carbenes in Transition Metal Catalysis in Topics in Organometallic Chemistry*, ed. by F. Glorius, Springer, Heidelberg, **2007**, Vol. 21. doi:10.1007/978-3-540-36930-1 b) *N-Heterocyclic Carbenes in Synthesis*, ed. by S. P. Nolan, Wiley-VCH, Weinheim, **2006**. c) W. A. Herrmann, *Angew. Chem., Int. Ed.* **2002**, *41*, 1290. d) S. Díez-González, N. Marion, S. P. Nolan, *Chem. Rev.* **2009**, *109*, 3612.
- 14 When the catalytic conjugate addition was quenched with D₂O instead of H₂O, 77% deuterium was incorporated at the α -position of **4aa**. This suggests that the product of the catalytic reaction is in a form of an enolate, but the attempted enolate trap with aldehydes was unsuccessful.
- 15 The reaction between **2a** and **3a** in the presence of the radical inhibitor, TEMPO (2,2,6,6-tetramethylpiperidine 1-oxyl) under otherwise identical conditions afforded **3aa** in 83% yield. Thus, the radical mechanism should be ruled out. For a conjugate addition of trialkylboranes to α,β -unsaturated aldehydes and ketones under radical conditions, see: H. C. Brown, G. W. Kabalka, *J. Am. Chem. Soc.* **1970**, *92*, 714.
- 16 A reaction mechanism should include the formation of alkylcopper(I) species through B/Cu transmetalation between an alkylborane and a copper(I) alkoxide and subsequent conjugate addition of the alkylcopper(I) species to an enone. See, refs. 8, 10a, and 10d.
- 17 Typical experimental procedure for the copper-catalyzed conjugate addition (Table 1, Entry 10): In a glove box, (9-BBN-H)₂ (58.6 mg, 0.24 mmol), alkene **1f** (145.5 mg, 0.504 mmol), and THF (0.20 mL) were placed in a vial containing a magnetic stirring bar. Then, the vial was sealed with a cap equipped with a Teflon-coated silicon rubber septum. The vial was removed from the glove box. The mixture was stirred at 60 °C for 1 h to prepare alkylborane **2f**. Also in the glove box, [(IPr)CuCl] (19.5 mg, 0.04 mmol), *t*-BuOK (4.5 mg, 0.04 mmol), and THF (0.20 mL) were placed in another vial and the vial was sealed with a cap equipped with a Teflon-coated silicon rubber septum. After the vial was removed from the glove box, the mixture was stirred at 25 °C for 1 h. Next, the alkylborane was transferred to the vial containing a Cu–IPr complex. Finally, aryl α,β -unsaturated ketone **3j** (106.5 mg, 0.40 mmol) in THF (0.4 mL) was added. After 8 h stirring at 80 °C, H₂O was added to the reaction mixture. Then, the mixture was diluted and extracted with EtOAc (2 mL $\times$ 3). The combined organic layer was dried over MgSO₄. Then, the drying agent was removed by filtration, and the resulting solution was evaporated under reduced pressure. The residue was purified by flash chromatography on silica gel (5–15% EtOAc/hexanes) to afford **4fj** in 71% yield (158.4 mg, 0.284 mmol). White solid. *R*_f 0.3 (20% EtOAc/hexanes). ¹H NMR (300 MHz, CDCl₃): δ 1.11–1.31 (m, 2H), 1.38 (s, 9H), 1.53–1.85 (m, 4H), 2.82 (t, *J* = 7.5 Hz, 2H), 3.27–3.30 (m, 2H), 3.40 (m, 1H), 3.91 (s, 3H), 4.84 (s, 2H), 7.20–7.26 (m, 5H), 7.35 (m, 1H), 7.41–7.46 (m, 3H), 7.54 (m, 1H), 7.84–7.91 (m, 4H). ¹³C NMR (75.4 MHz, CDCl₃): δ 24.92, 26.99, 27.72, 36.11, 38.02, 40.82, 45.48, 47.15, 51.97, 83.03, 127.07, 127.51, 127.69, 128.07, 128.30, 128.40, 128.52, 128.60, 130.36, 132.79, 133.08, 137.10, 138.41, 145.28, 153.16, 167.31, 176.10, 198.79. Mp 102.1–103.5 °C. Anal. Calcd for C₃₄H₃₉NO₆: C, 73.23; H, 7.05%. Found: C, 73.40; H, 7.30%.