

- Pearson, W. H., Lian, B. W. & Bergmeier, S. C. (1996). *Comprehensive Heterocyclic Chemistry II*, Vol. 1A, edited by A. Padwa, pp. 1–60. New York: Pergamon Press.
- Sheldrick, G. M. (1996a). *SADABS. Program for Empirical Absorption Correction of Area Detector Data*. University of Göttingen, Germany.
- Sheldrick, G. M. (1996b). *SHELXTL Reference Manual*. Version 5. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1996a). *SAINT Software Reference Manual*. Version 4. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1996b). *SMART Software Reference Manual*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Sriraghavan, K. & Ramakrishnan, V. T. (1998). Unpublished work.
- Tanner, D. (1994). *Angew. Chem. Int. Ed. Eng.* **33**, 599–619.

Acta Cryst. (1998). **C54**, 1301–1302

N-(3-Hydroxyphenyl)-*p*-toluenesulfonamide

SHYAMAPROSAD GOSWAMI,^a AJIT KUMAR MAHAPATRA,^b
GUR DAYAL NIGAM,^c KANDASAMY CHINNAKALI^{d†} AND
HOONG-KUN FUN^d

^aDepartment of Chemistry, Bengal Engineering College, Botanical Garden, Howrah 711 103, India, ^bDepartment of Chemistry, Indian Institute of Technology, Kharagpur 721 302, West Bengal, India, ^cDepartment of Physics and Meteorology, Indian Institute of Technology, Kharagpur 721 302, West Bengal, India, and ^dX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia. E-mail: hkfun@usm.my

(Received 5 March 1998; accepted 1 April 1998)


Abstract

In the title compound, C₁₃H₁₃NO₃S, the dihedral angle between the toluene and phenol moieties is 69.48 (5)°. The molecules form columns linked by N—H···O and O—H···O hydrogen bonds.

Comment

Molecular recognition is central to the understanding of biological activity and enzymatic systems (Goswami & Ghosh, 1997). The title compound, (I), contains a potentially active sulfonamide moiety. Sulfonamides have been extensively used against many common gram-positive bacterial infections like pneumonia, meningitis, dysentery and urinary-tract infection (Kar, 1993). Hydrogen bonding is expected to play an important role

in governing the crystal structure of the title compound (Aoyama *et al.*, 1995). The X-ray structure determination was carried out in order to elucidate the molecular conformation and hydrogen bonding.



Bond lengths agree with other reported values (Allen *et al.*, 1987). The sum of the bond angles around the N1 atom [349 (1)°] indicates that it is in a pyramidal configuration. The dihedral angle between the toluene and phenol moieties is 69.48 (5)°. Chains of molecules along *b* are linked by O—H···O hydrogen bonds, and glide-related chains are interlinked by N—H···O hydrogen bonds (see Table 2 for details). These two chains form a column-like structure along the *b* axis, with the phenol groups occupying a plane and the toluene groups projecting outwards from it on either side. The toluene groups of neighbouring columns are arranged close to each other, but are not stacked to allow any π – π interactions.




Fig. 1. The structure and atom-numbering scheme of the title compound, shown with 50% probability displacement ellipsoids.

Experimental

For the preparation of the title compound, *p*-toluenesulfonyl chloride (875 mg, 4.5 mmol) was added to a solution of *m*-aminophenol (500 mg, 4.5 mmol) and dry pyridine (0.5 ml) in dry CH₂Cl₂ solvent at 283 K. The reaction mixture was stirred for 15 min at this temperature and for a further 1 h at room temperature. After the usual work-up, the desired compound was obtained in 80% yield (Kurita, 1974). Single crystals were grown by slow evaporation of a 4:1 dichloromethane–methanol solution of the compound.

Crystal data

C₁₃H₁₃NO₃S
M_r = 263.30

Mo K α radiation
 λ = 0.71073 Å

† On leave from: Department of Physics, Anna University, Chennai 600 025, India.

Monoclinic

C2/c

 $a = 23.9096(7) \text{ \AA}$ $b = 9.5192(3) \text{ \AA}$ $c = 11.9629(3) \text{ \AA}$ $\beta = 112.136(1)^\circ$ $V = 2522.07(13) \text{ \AA}^3$ $Z = 8$ $D_x = 1.387 \text{ Mg m}^{-3}$ D_m not measured

Cell parameters from 6375

reflections

 $\theta = 1.5\text{--}33.0^\circ$ $\mu = 0.256 \text{ mm}^{-1}$ $T = 293(2) \text{ K}$

Block

 $0.52 \times 0.36 \times 0.30 \text{ mm}$

Colourless

Data collection: *SMART* (Siemens, 1996b). Cell refinement: *SAINT* (Siemens, 1996a). Data reduction: *SAINT*. Program(s) used to solve structure: *SHELXTL* (Sheldrick, 1996b). Program(s) used to refine structure: *SHELXTL*. Molecular graphics: *SHELXTL*. Software used to prepare material for publication: *SHELXTL* and *PARST* (Nardelli, 1995).

The authors would like to thank the Malaysian Government and Universiti Sains Malaysia for research grant R&D No. 190-9609-2801. AKM is thankful to the Department of Science and Technology, New Delhi, for the award of a Senior Research Fellowship. KC thanks the Universiti Sains Malaysia for a Visiting Postdoctoral Fellowship.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: HA1219). Services for accessing these data are described at the back of the journal.

Data collection

Siemens SMART CCD area-detector diffractometer

 ω scans

Absorption correction:

empirical (*SADABS*;

Sheldrick, 1996a)

 $T_{\min} = 0.871$, $T_{\max} = 0.955$

8060 measured reflections

2898 independent reflections

2474 reflections with

 $I > 2\sigma(I)$ $R_{\text{int}} = 0.016$ $\theta_{\max} = 27.5^\circ$ $h = -30 \rightarrow 28$ $k = 0 \rightarrow 12$ $l = 0 \rightarrow 15$

Refinement

Refinement on F^2 $R[F^2 > 2\sigma(F^2)] = 0.036$ $wR(F^2) = 0.102$ $S = 1.058$

2897 reflections

215 parameters

All H atoms refined

 $w = 1/[\sigma^2(F_o^2) + (0.0496P)^2$ $+ 1.4298P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.218 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.280 \text{ e \AA}^{-3}$

Extinction correction: none

Scattering factors from

*International Tables for**Crystallography* (Vol. C)

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Aoyama, Y., Imai, Y., Endo, K. & Kobayashi, K. (1995). *Tetrahedron*, **51**, 343–352.
- Goswami, S. P. & Ghosh, K. (1997). *Tetrahedron Lett.* **38**, 4503–4506.
- Kar, A. (1993). In *Medicinal Chemistry*. New Delhi: Wiley Eastern.
- Kurita, K. (1974). *Chem. Ind. (London)*, p. 345.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Sheldrick, G. M. (1996a). *SADABS. Program for Empirical Absorption Correction of Area Detector Data*. University of Göttingen, Germany.
- Sheldrick, G. M. (1996b). *SHELXTL Reference Manual*. Version 5. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1996a). *SAINT Software Reference Manual*. Version 4. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1996b). *SMART Software Reference Manual*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Table 1. Selected geometric parameters ($\text{\AA}$, $^\circ$)

S—O2	1.4259 (13)	S—C7	1.760 (2)
S—O3	1.4347 (12)	O1—C1	1.371 (2)
S—N1	1.6214 (13)	N1—C5	1.432 (2)
O2—S—O3	119.55 (8)	O3—S—C7	107.62 (8)
O2—S—N1	108.65 (8)	N1—S—C7	107.96 (7)
O3—S—N1	104.59 (7)	C5—N1—S	123.62 (11)
O2—S—C7	107.97 (8)		
C4—C5—N1—S	52.0 (2)	N1—S—C7—C8	−74.3 (2)
C5—N1—S—C7	64.19 (14)		

Table 2. Hydrogen-bonding geometry ($\text{\AA}$, $^\circ$)

D—H...A	D—H	H...A	D...A	D—H...A
O1—H1O1...O3 ⁱ	0.79 (3)	1.95 (3)	2.727 (2)	171 (2)
N1—H1N1...O1 ⁱⁱ	0.85 (2)	2.14 (2)	2.983 (2)	174 (2)

Symmetry codes: (i) $x, 1+y, z$; (ii) $\frac{1}{2} - x, y - \frac{1}{2}, \frac{1}{2} - z$.

The data collection covered over a hemisphere of reciprocal space by a combination of three sets of exposures; each set had a different φ angle ($0, 88$ and 180°) for the crystal and each exposure of 30 s covered 0.3° in ω . The crystal-to-detector distance was 4 cm and the detector swing angle was -35° . Coverage of the unique set is over 99% complete. Crystal decay, monitored by repeating 30 initial frames at the end of data collection and analysing the duplicate reflections, was found to be negligible. The structure was solved by direct methods and refined by full-matrix least-squares techniques. All H atoms were located from a difference Fourier map and refined isotropically.

Acta Cryst. (1998). **C54**, 1302–1304

Cyclodecanol

EDWARD J. VALENTE,^a DIWAKAR M. PAWAR^b AND
ERIC A. NOE^b

^aDepartment of Chemistry, Mississippi College, Clinton, MS 39058-4065, USA, and ^bDepartment of Chemistry, Jackson State University, Jackson, MS 39217-0510, USA. E-mail: valente@mc.edu

(Received 12 January 1998; accepted 24 February 1998)

Abstract

Cyclodecanol crystals, C₁₀H₂₀O, grown by slow sublimation, occur in the monoclinic system in space group