# Accepted Manuscript

Synthesis of four diastereomers and structural revision of tetradenolide

Maki Tokuda, Yuji Kurogome, Rieko Katoh, Yukie Nohara, Yasunao Hattori, Hidefumi Makabe

PII: DOI: Reference:	S0040-4039(14)00940-X http://dx.doi.org/10.1016/j.tetlet.2014.05.109 TETL 44700
To appear in:	Tetrahedron Letters
Received Date:	16 April 2014
Revised Date:	22 May 2014
Accepted Date:	28 May 2014



Please cite this article as: Tokuda, M., Kurogome, Y., Katoh, R., Nohara, Y., Hattori, Y., Makabe, H., Synthesis of four diastereomers and structural revision of tetradenolide, *Tetrahedron Letters* (2014), doi: http://dx.doi.org/ 10.1016/j.tetlet.2014.05.109

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Graphical Abstract** To create your abstract, type over the instructions in the template box below. Fonts or abstract dimensions should not be changed or altered.

Synthesis of four diastereomers and structural revision of	Leave this area blank for abstract info.
tetradenolide	
Maki Tokuda, Yuji Kurogome, Rieko Katoh, Yukie Nohara, Ya	sunao Hattori, and Hidefumi Makabe*
o C	
	о он
о́н	ОН ОН



Tetrahedron Letters

journal homepage: www.elsevier.com

## Synthesis of four diastereomers and structural revision of tetradenolide

Maki Tokuda,<sup>b</sup> Yuji Kurogome,<sup>a</sup> Rieko Katoh,<sup>a</sup> Yukie Nohara,<sup>b</sup> Yasunao Hattori,<sup>c</sup> and Hidefumi Makabe<sup>a</sup>\*

<sup>a</sup>Graduate School of Agriculture, Sciences of Functional Foods, Shinshu University, 8304 Minami-minowa Kami-ina, Nagano, 399-4598, Japan <sup>b</sup>Department of Bioscience and Biotechnology, Faculty of Agriculture, Shinshu University, 8304 Minami-minowa Kami-ina, Nagano, 399-4598, Japan <sup>c</sup>Department of Medicinal Chemistry, Kyoto Pharmaceutical University, Yamashina-ku, Kyoto 607-8412, Japan

## ARTICLE INFO

ABSTRACT

Article history: Received Received in revised form Accepted Available online

Keywords: tetradenolide α-pyrone structural revision total synthesis Four diastereomers of tetradenolide, a cytotoxic  $\alpha$ -pyrone isolated from *Tetradenia riparia*, were synthesized stereoselectively using Z-selective Horner-Emmons reaction followed by acid catalyzed lactonization. Making comparison of the <sup>1</sup>H and <sup>13</sup>C NMR spectral data of the four diastereomers with those of the reported value of natural product did not lead to determine the relative stereochemistry of the natural tetradenolide. Thus detailed investigation of the spectral data of the related compounds led us to revise the structure of tetradenolide as deacetylboronolide.

2009 Elsevier Ltd. All rights reserved.

Tetradenolide is an antimicrobial, phytotoxic and cytotoxic αpyrone isolated by Luc Van Puyvelde and De Kimpe from Tetradenia riparia.<sup>1</sup> The related compound (+)deacetylboronolide was isolated Luc Van Puyvelde and coworkers from the same plant, Tetradenia riparia (Hochst) N. E. Br. (Labiatae)<sup>2</sup> and (+)-boronolide was isolated from *Tetradenia* fruticosa Benth and Tetradenia barbera (N. E. Br.) Codd, respectively.<sup>3,4</sup> As to the biological activity of boronolides, antimalarial activity was reported (Figure 1).<sup>2,5</sup> Many reports of the syntheses of (+)-boronolide and (+)-deacetylboronolide have been published so far.<sup>6</sup> However, synthetic study of teradenolide has not been reported yet. The reported structure of tetradenolide was shown in Figure 1 and the relative stereochemistry and optical rotation value of this compound were not reported.<sup>1</sup>



Figure 1. The reported structure of tetradenolide and related compounds.

The synthesis of the four diastereomers and making comparison of their <sup>1</sup>H and <sup>13</sup>C NMR spectral data with those of natural tetradenolide would be a crew to determine its relative stereochemistry. Therefore we planned to synthesize four diastereomers of tetradenolide (Figure 2).





At first we began to synthesize (6S, 1'S, 2'R)-tetradenolide **(1a)** starting from 1,3-propanediol **(2)**. Mono-protection of 1,3propanediol **(2)** with benzyl bromide in the presence of NaH followed by oxidation with SO<sub>3</sub> • pyridine afforded aldehyde **3**. Alkylation of the 1-hexyne with aldehyde **3** using *n*-BuLi gave **4** which was subjected to reduction with LiAlH<sub>4</sub> in THF under reflux to furnish *E*-allylic alcohol **5**. Oxidation of **5** with Dess-Martin periodonane led to enone **6**.<sup>7</sup> Enone **6** was subjected to the Sharpless asymmetric dihydroxylation using AD mix  $\beta$  to give diol **7** in 97% ee analyzed by Mosher ester derivative.<sup>8</sup> Protection of the resulting diol with 2,2-dimethoxypropane in the presence of PPTS afforded **8**. Reduction of the ketone **8** with L-Selectride gave **9** in a single diastereomer.<sup>9</sup> To confirm the stereochemistry of newly formed oxymethine carbon, advanced Mosher method was used (supplementary content).<sup>10</sup> We confirmed that this carbinol center possessed *S* configuration. Secondary hydroxy

\* Corresponding author. Tel. +81 265 77 1630; fax +81 265 77 1700; e-mail: makabeh@shinshu-u.ac.jp

## Tetrahedron Letters

group of **9** was protected as ethoxyethoxy group to afford **10** subsequent removal of benzyl group with Na in liquid NH<sub>3</sub> followed by oxidation with Dess-Martin periodonane furnished aldehyde **11**. The resultant aldehyde **11** was subjected to Z-selective Horner-Emmons reaction using diphenyl phosphonoacetate to afford **12** with 1:19 of *E:Z* ratio.<sup>11</sup> Finally, **12** was treated with acid in THF-H<sub>2</sub>O under reflux to give (6S, 1'S, 2'R)-tetradenolide (**1a**) (Scheme 1).



Scheme 1. Synthesis of (6S,1'S,2'R)-tetradenolide (1). Reagents and conditions: (a) (i) BnBr, NaH, NaI, (ii) SO<sub>3</sub> • pyridine, DMSO, 35%; (b) *n*-BuLi, 1-hexyne, 90%; (c) LiAlH<sub>4</sub>, THF reflux, 77%; (d) Dess-Martin periodinane, 68%; (e) AD mix $\beta$ , MeSO<sub>2</sub>NH<sub>2</sub>, 78%; (f) 2,2-dimethoxypropane, PPTS, quant.; (g) L-Selectride, 90%; (h) ethyl vinyl ether, PPTS, 96%; (i) (i) Na, NH<sub>3</sub>, (ii) Dess-Martin periodinane, 60%; (j) (PhO)<sub>2</sub>P(O)CH<sub>2</sub>CO<sub>2</sub>Et, NaH, 66%; (k) HCl, THF-H<sub>2</sub>O, 51%

Synthesis of (6R, 1'S, 2'R)-tetradenolide (**1b**) was started from **9** which was an intermediate for the synthesis of **1a**. The secondary hydroxy group of **9** was inverted using Mitsunobu reaction with *p*-nitrobenzoic acid in the presence of DEAD and PPh<sub>3</sub>.<sup>12</sup> Hydrolysis of *p*-nitrobenzoyl ester with methanolic NaOH gave **13**. Compound **13** was converted to (6R, 1'S, 2'R)-tetradenolide (**1b**) as described in the synthesis of **1a** (Scheme 2).



Scheme 2. Synthesis of (6R, 1'S, 2'R)-tetradenolide (1b). Reagents and conditions: (a) (i) *p*-NBA, DEAD, PPh<sub>3</sub>, (ii) NaOH, MeOH, 85%; (b) ethyl vinyl ether, PPTS, 96%; (c) (i) Na, NH<sub>3</sub>, (ii) Dess-Martin periodinane, 60%; (d) (PhO)<sub>2</sub>P(O)CH<sub>2</sub>CO<sub>2</sub>Et, NaH, 66%; (k) HCl, THF-H<sub>2</sub>O, 55%.

Synthesis of (6S, 1'R, 2'R)-tetradenolide (**1c**) was started from Dribose (**17**).<sup>13</sup> Protection of the diol at 2,3-position with acetone in the presence of sulfric acid followed by Wittig reaction with CH<sub>3</sub>P<sup>+</sup>Ph<sub>3</sub>T afforded diol **18**. Introduction of the triisopropyl benzenesulfonyl (Tris) group at primary hydroxy group under basic condition followed by treating with NaH gave terminal epoxide **19**. Alkylation of **19** using  $C_3H_7MgCl$  in the presence of CuI gave **20**. Protection of the resulting secondary hydroxy group as MOM ether followed by hydroboration using 9-BBN and treatment with NaOH and  $H_2O_2$  furnished **21**. Compound **21** was converted to  $(6S, 1^{\prime}R, 2^{\prime}R)$ -tetradenolide (**1c**) as described in the synthesis of **1a** (Scheme 3).



Scheme 3. Synthesis of  $(6S, 1^{\prime}R, 2^{\prime}R)$ -tetradenolide (1c). Reagents and conditions: (a) (i) H<sub>2</sub>SO<sub>4</sub>, acetone, (ii) CH<sub>3</sub>P<sup>+</sup>Ph<sub>3</sub>Γ, *t*-BuOK, 71%; (b) (i) TrisCl, pyridine; (ii) NaH, 63%; (c) C<sub>3</sub>H<sub>7</sub>MgCl, CuI; 69%; (d) (i) MOMCl, *i*-Pr<sub>2</sub>NEt, (ii) 9-BBN, NaOH, H<sub>2</sub>O<sub>2</sub>; 44%; (e) (i) Dess-Martin periodinane, (ii) (PhO)<sub>2</sub>P(O)CH<sub>2</sub>CO<sub>2</sub>Et, NaH, 30%; (f) HCl, THF-H<sub>2</sub>O, 60%

Next, we began to synthesize  $(6R, 1^{2}R, 2^{2}R)$ -tetradenolide (1d). Alkylation of 4-butyn-1-ol (23) with 1-pentanal (24) using n-BuLi gave 25 which was subjected to reduction with LiAlH<sub>4</sub> in THF under reflux to furnish E-allylic alcohol 26. Protection of the primary hydroxy group of **26** with TBDPSCl in the presence of Et<sub>3</sub>N and DMAP afforded 30. Oxidation of the secondary hydroxy group of 27 with Dess-Martin periodinane gave enone 28. Enone 28 was subjected to the Sharpless asymmetric dihydroxylation using AD mix  $\beta$  to give diol **29** in 97% ee analyzed by Mosher ester derivative.<sup>8</sup> Protection of the resulting diol with 2,2-dimethoxypropane in the presence of PPTS afforded 30. Reduction of the ketone 30 with L-Selectride gave 31. To confirm stereochemistry of the newly formed oxymethine carbon, advanced Mosher method was used and this carbinol center was confirmed to be S configuration (supplementary content).<sup>10</sup> The secondary hydroxy group of **31** was inverted using Mitsunobu reaction with p-nitrobenzoic acid in the presence of DEAD and PPh<sub>3</sub>. Hydrolysis of *p*-nitrobenzoyl ester with methanolic NaOH gave 32. Protection of the resulting secondary hydroxy group as MOM ether followed by deprotection of the TBDPS group using TBAF gave 33 Compound 33 was converted to  $(6R, 1^{\prime}R, 2^{\prime}R)$ -tetradenolide (1d) as described in the synthesis of 1a (Scheme 4).





Scheme 4. Synthesis of (6R, 1'R, 2'R)-tetradenolide (1d). Reagents and conditions: (a) *n*-BuLi, 64%; (b) LiALH<sub>4</sub>, THF reflex, 69%; (c) TBDPSCl, Et<sub>3</sub>N, DMAP, 88%; (d) Dess-Martin periodinane, 83%; (e) AD mix $\beta$ , MeSO<sub>2</sub>NH<sub>2</sub>, 84%; (f) 2,2-dimethoxypropane, PPTS, 89%; (g) L-Selectride, 96%; (h) (i) *p*-NBA, DEAD, PPh<sub>3</sub>, (ii) NaOH, MeOH, 73%; (i) (i) MOMBr, *i*-Pr<sub>2</sub>NEt, (ii) TBAF, 93%; (j) (i) Dess-Martin periodinane, (ii) (PhO)<sub>2</sub>P(O)CH<sub>2</sub>CO<sub>2</sub>Et, NaH, 44%; (k) HCl, THF-H<sub>2</sub>O, 75%.

As to the compounds **1a** and **1b**, structurally similar compounds **35** and **36** were synthesized by Rúveda and coworkers who described the stereoselective transformation of the naturally abundant argentilactone in 1988.<sup>14</sup> The <sup>13</sup>C NMR spectral data of carbinol centers of **1a** and **1b** were quite similar to those of **35** and **36**, respectively (Figure 3).



Figure 3. The related compounds 35 and 36 which were synthesized by Rúveda and co-workers.

As four diastereomers of tetradenolide were in hand, we compared their spectral data with those of the reported values to determine the relative stereochemistry. As to the <sup>1</sup>H NMR spectra, it is rather difficult to compare because the range of the chemical shifts of reported <sup>1</sup>H NMR spectra had a wide range (see supplementary information). Thus we focused on analyzing <sup>13</sup>C NMR spectra. The chemical shifts of <sup>13</sup>C NMR of natural one and synthetic **1a-1d** were different especially C-6 and C-2' carbons. Therefore we could not determine the relative stereochemistry of tetradenolide (Figure 4).



**Figure 4.** The difference of chemical shifts of <sup>13</sup>C NMR data of synthetic **1a-1d** and natural tetradenolide at C-6 and C-2' position.

Thus we carefully examined the <sup>13</sup>C NMR data of the related compounds of tetradenolide.<sup>2</sup> We noticed that the <sup>13</sup>C NMR data of deacetylboronolide synthesized by Trost and co-worker were quite similar to those of tetradenolide (Table 1).<sup>15</sup> Thus we synthesized deacetylboronlide to examine its <sup>13</sup>C NMR spectra in

detail.<sup>16</sup> We found that the signals of <sup>13</sup>C NMR at C-1' and C-2' positions of deacetylboronolide were completely overlapped using HMQC experiment.<sup>16</sup> Therefore, it was possible that the <sup>13</sup>C NMR of deacetylboronolide was misread as tetradenolide.<sup>1</sup>

As to the EI-MS spectrum of natural tetradenolide, Puyvelde et al. reported that M<sup>+</sup> could not be observed. We also measured EI-MS spectra of **1a-1b** and found that only tiny signals of M<sup>+</sup> were observed. Puyvelde et al. prepared bis-O-TMS derivative of natural tetradenolide and reported that  $M^+$  (358, 0.2%) was observed at 70 eV.1 We prepared bis-O-TMS derivative of 1b using bis(trimethylsilyl)acetamide (BSA) and measured EI-MS at 70eV (Figure 4). However, M<sup>+</sup> (358, 0.2%) was not detected but  $(M-Me)^+$  (343),  $(M-C_4H_9)^+$  (301), and  $(M-C_4H_9-2Me)^+$  (272) were observed. The pattern of fragmentation was different from that of reported one of natural tetradenolide derivative.<sup>1</sup> We also prepared tris-O-TMS derivative of (+)-deacetylboronolide and measured EI-MS at 70 eV.<sup>1</sup> As Puyvelde et al. reported, m/z 358 (0.2%) was observed and the pattern of fragmentation was similar to that of natural tetradenolide derivative except that the signals of  $(M-Me)^+$  (445),  $(M-C_4H_9)^+$ (403),  $(M-C_4H_9-2Me)^+$  (374) were observed.



Figure 5. Preparation of TMS derivatives of 1b and (+)-deacetylboronolide.

Next, we measured CI-MS spectra of **1a-1d** and (+)deacetylboronolide. We could observe  $(M+H)^+$  clearly. We found that CI-MS was effective to obtain satisfied data for hydroxylated  $\delta$ -lactones. The melting point value of reported tetradenolide was (102.8-103°C) also very close to that of (+)-deacetylboronolide synthesized by us (101-102 °C<sup>16</sup>) lit.{ (99-100°C,<sup>2a</sup> 101°C<sup>17</sup>)}.<sup>1</sup> The sample of natural tetradenolide and its spectral data are no longer available,<sup>18</sup> however, it is highly possible that the structure of tetradenolide is (+)-deacetylboronolide from reasons as follows: 1) <sup>13</sup>C NMR spectral data of natural tetradenolide was identical with those of (+)-deacetylboronolide; 2) The melting point value of natural tetradenolide was very close to that of (+)deacetylboronolide; 3) natural tetradenolide and (+)deacetylboronolide were isolated from the same plant, *Tetradenia riparia*.<sup>12</sup>

In conclusion, four diastereomers of tetradenolide, a cytotoxic  $\alpha$ -pyrone isolated from *Tetradenia riparia*, were synthesized. The <sup>1</sup>H and <sup>13</sup>C NMR spectral data of the four diastereomers were not consistent with those of reported values of natural tetradenolide. The detailed investigation of the <sup>13</sup>C NMR data and melting point data of the related compounds led us to revise the structure of tetradenolide as (+)-deacetylboronolide.

13 -----

## Tetrahedron

Table 1.	<sup>13</sup> C NMR	data of	tetradenolide an	nd (+)-dead	etylboronolide
----------	---------------------	---------	------------------	-------------	----------------

position	tetradenolide	position	(+)-deacetylboronolide (Trost et al.) <sup>a</sup>	(+)-deacetylboronolide (by us)	
2	164.02	2	163.98	164.10	
3	120.95	3	120.90	120.84	
4	146.11	4	146.09	146.21	
5	25.79	5	25.75	25.73	
6	77.16	6	77.12	77.15	
		1'	76.74 (overlapped with solvent)	74.24 (overlapped with 2' carbon)	
1'	74.36	2'	74.30	74.24 (overlapped with 1' carbon)	$\bigcirc$
2'	70.20	3'	70.14	70.20	K
3'	33.46	4'	33.42,	33.36	
4'	27.67	5'	27.64	27.64	
5'	22.62	6'	22.59	22.59	
6'	14.00	7'	13.98	13.99	

<sup>a</sup>Trost and co-worker reported that the peak at 76.74 ppm was observed.<sup>15</sup> However, we could not observe this signal in the HMQC spectrum and found that both of C1' and C-2' signals were overlapped at 74.24 ppm.<sup>1</sup>

### Acknowledgments

We thank Prof. Dr. De Kimpe, N. for his valuable information about α-pyrone natural products. This work was supported in part by JSPS KAKENHI Grant Number 24580160 to H. M.

### **References and notes**

- Van Puyvelde, L.; De Kimpe, N. Phytochemistry 1998, 49, 1157. 1
- 2. (a) Van Puyvelde, L.; Dube, S.; Uwimana, E.; Uwera, C.: Domisse, R. A.; Esmans, E. L.; Van Schoor, O.; Vlietinck, A. Phytochemistry 1979, 18, 1215; (b) Van Puyvelde, L.; De Kimpe, N.; Dube, S.; Chagnon-Dube, M.; Boily, Y.; Borremans, F.; Schamp, N.; Anteunis, M. J. O. Phytochemistry 1981, 20, 2753.
- 3. Franca, N. C.; Polonsky, J. C. R. Hebd. Seances, Acad. Sci., Ser. C. 1971. 273. 439.
- Davies-Coleman, M. T.; Rivett, D. E. A. Phytochemistry 1987, 26, 4. 3047.
- Watt, J. M.; Brandwijk, M. G. B. The Medicinal and Poisonous 5. Plants of Southern and Eastern Africa, Livingston: Edinburgh, 1962, p516.
- (a) Nagano, H.; Yasui, H. Chem. Lett. 1992, 1045. (b) Honda, T.; 6. Horiuchi, S.; Mizutani, H.; Kanai, K. J. Org. Chem. 1996, 61, 4944. (c) Ghosh, A. K.; Bilcer, G. Tetrahedron Lett. 2000, 41, 1003. (d) Trost, B. M.; Yeh, V. S. C. Org. Lett. 2002, 4, 3513. (e) Carda, M.; Rodoríguez, S.; Segovia, B.; Marco, J. A. J. Org. Chem. 2002, 67, 6560. (f) Chandrasekhar, M.; Chandra, K. L.; Singh, V. K. J. Org. Chem. 2003, 68, 4039. (g) Hu, S. -G.; Hu, T. -S.; Wu, Y. -L. Org. Biomol. Chem. 2004, 2, 2305. (h) Naidu, S. V.; Gupta, P.; Kumar, P. Tetrahedron Lett. 2005, 46, 2129. (i) Boruwa, J.; Barua, N. C. Tetrahedron 2006, 62, 1193. (j) Kumar, P.; Naidu, S. V. J. Org. Chem. 2006, 71, 3935. (k) Prasad, K. R.; Anbarasan, P. Tetrahedron: Asymmetry 2006, 17, 1146. (1) Raghavan, S.; Krishnaiah, V. Tetrahedron Lett. 2006, 47, 7611. (m) Yamauchi, S. ; Isozaki, Y.; Nishimura, Y.; Tsuda, H.; Nishiwaki, T.; Shuto, Y. Biosci. Biotechnol. Biochem. 2012, 76, 1708. (n) Reddy, B. V. S.; Reddy, V. V.; Praneeth, K. Tetrahedron Lett. 2014, 55, 1398.
- Dess, D. B.; Martin, J. C. J. Am. Chem. Soc. 1991, 113, 7277.
- 8 Kolb, H.; VanNiewenhze, M. S.; Sharpless, K. B. Chem. Rev. **1994**, *94*, 2483.
- Ghosh A. K.; Bilcer, G. Tetrahedron Lett. 2000, 41, 1003. 9
- 10. Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. J. Am. Chem. Soc. 1991, 113, 4092.
- 11. (a) Ando, K. Tetrahedron Lett. 1995, 36, 410; (b) Ando, K. J. Org. Chem. 1997, 62, 1934.
- 12. Mitsunobu, O. Synthesis 1981, 1.

- 13. (a) Srihari, P.; Kumaraswamy, B.; Yadav, J. S. Terahedron 2009, 65, 6304; (b) Srihari, P.; Kumaraswamy, B.; Rao, G. M.; Yadav, J. S. Terahedron: Asymmetry 2010, 21, 106.
- Signorella, S. R.; Cravero, R. M.; Sala, L. F.; Rúveda, E. A. Synth. 14. Commun. 1988, 18, 1935.
- 15. Trost, B. M.; Yeh, V. S. C. Org. Lett. 2002, 4, 3513.
- Kurogome, Y.; Hattori, Y.; Makabe, H. Tetrahedron Lett. 2014, 16. 55, 2822.
- Chandrasekhar, M.; Chandra, K. L.; Singh, V. K. J. Org. Chem. 17. 2003, 68, 4039.
- 18. Personal communication with Prof. Dr. De Kimpe, N.

## **Supplementary Material**

Supplementary material associated with this article can be found in the online version at doi:

## **Graphical Abstract**

## Synthesis of four diastereomers and structural revision of tetradenolide

Maki Tokuda, Yuji Kurogome, Rieko Katoh, Yukie Nohara, Yasunao Hattori, and Hidefumi Makabe\*

OH OH ŌH ŌH ÓН tetradenolide deacetylboronolide