Palladium-Catalyzed Insertion of *N*-tosylhydrazones and Trapping with Carbon Nucleophiles

LETTERS XXXX Vol. XX, No. XX 000–000

ORGANIC

Ping-Xin Zhou, Yu-Ying Ye, and Yong-Min Liang*

State Key Laboratory of Applied Organic Chemistry, Lanzhou University, and State Key Laboratory of Solid Lubrication, Lanzhou Institute of Chemical Physics, Chinese Academy of Science, Lanzhou 730000, P. R. China

liangym@lzu.edu.cn

Received August 26, 2013

ABSTRACT



A Pd-catalyzed three-component cross-coupling reaction of vinyl iodide, *N*-tosylhydrazone, and carbon nucleophiles is reported, and a one-pot procedure is also developed. The cross-coupling is proposed to proceed through a palladium—carbene migratory insertion, carbopalladation other than classic palladium—carbene migratory insertion, and β -H elimination. Moreover, the reaction proceeds under mild conditions and with high stereoselectivity.

Transition-metal-catalyzed cross-coupling reactions have been recognized as a powerful and efficient method for C–C bond construction.¹ Despite the wide applicability of these reactions, most of these cross-coupling transformations involve the participation of a stoichiometric organometallic reagent, which are usually expensive, toxic, and moisturesensitive.² Moreover, highly reactive organometallic reagents often suffer from a problem of functional group tolerance. Thus, it is highly desirable to further develop a novel coupling process.

The utility of *N*-tosylhydrazones in organic synthesis has a long history. Bamford and Stevens initially demonstrated that *N*-tosylhydrazones can be employed as a source of diazo compounds.³ The first example of showing their utility in a palladium-catalyzed cross-coupling reaction was reported by Barluenga.⁴ Since then, palladium-catalyzed insertion of *N*-tosylhydrazones has been emerging as a new type of cross-coupling reaction and has attracted increasing attention. In 2009, Wang reported a palladium-catalyzed cross-coupling of benzyl halides with tosylhydrazones to generate a variety of polysubstituted olefins.⁵ Recently, Barluenga reported the first examples of the use of alkenyl halides in cross-coupling reactions with tosylhydrazones to give linear dienes (Scheme 1, eq 1).⁶ However, all of these catalytic cycles involve Pd carbene migratory insertion followed by β -hydride elimination, and only one C(sp²)– C(sp²) bond is formed.⁷ If we can combine Pd-catalyzed insertion of *N*-tosylhydrazones and Pd-catalyzed cascade reactions, it would provide new possibilities for the development of more novel coupling processes. Indeed, a threecomponent coupling of aryl halides, *N*-tosylhydrazones,

 ^{(1) (}a) Negishi, E. Handbook of Organopalladium Chemistry for Organic Synthesis; Wiley: New York, 2002. (b) de Meijere, A.; Diederich, F. Metal-Catalyzed Cross-Coupling Reactions, 2nd ed.; Wiley-VCH: Weinheim, 2004. (c) Krische, M. J. Metal Catalyzed Reductive C-C Bond Formation: A Departure from Preformed Organometallic Reagents; Springer: Berlin, 2007.

^{(2) (}a) Komiya, S. Synthesis of Organometallic Compounds: A Practical Guide; Wiley-VCH: Chichester, 1997. (b) Terao, J.; Kambe, N. Acc. Chem. Res. 2008, 41, 1545–1554. (c) Jana, R.; Pathak, T. P.; Sigman, M. S. Chem. Rev. 2011, 111, 1417–1492.

⁽³⁾ Bamford, W. R.; Stevens, T. S. J. Chem. Soc. 1952, 4735-4740.

⁽⁴⁾ Barluenga, J.; Moriel, P.; Valdés, C.; Aznar, F. Angew. Chem., Int. Ed. 2007, 46, 5587–5590.

⁽⁵⁾ Xiao, Q.; Ma, J.; Yang, Y.; Zhang, Y.; Wang, J. Org. Lett. 2009, 11, 4732–4735.

⁽⁶⁾ Barluenga, J.; Tomás-Gamasa, M.; Aznar, F.; Valdés, C. Adv. Synth. Catal. 2010, 352, 3235–3240.

⁽⁷⁾ For selected palladium-catalyzed insertion of *N*-tosylhydrazones processes based on migratory insertion, followed by β -hydride elimination, see: (a) Barluenga, J.; Escribano, M.; Moriel, P.; Aznar, F.; Valdés, C. *Chem.—Eur. J.* **2009**, *15*, 13291–13294. (b) Zhou, L.; Ye, F.; Ma, J.; Zhang, Y.; Wang, J. *Angew. Chem., Int. Ed.* **2011**, *50*, 3510–3514. (c) Chen, Z.-S.; Duan, X.-H.; Wu, L.-Y.; Ali, S.; Ji, K.-G.; Zhou, P.-X.; Liu, X.-Y.; Liang, Y.; Wang, J. *Org. Lett.* **2011**, *17*, 6918–6921. (d) Zhou, L.; Ye, F.; Zhang, Y.; Wang, J. *Org. Lett.* **2012**, *14*, 922–925. (e) Xiao, Q.; Wang, B.; Tian, L.; Yang, Y.; Ma, J.; Zhang, Y.; Chen, S.; Wang, J. *Angew. Chem., Int. Ed.* **2013**, *52*, 9305–9308. (f) Zeng, X.; Cheng, G.; Shen, J.; Cui, X. *Org. Lett.* **2013**, *15*, 3022–3025.

and terminal alkynes for the synthesis of propargylarenes was developed by Wang, in which two separate C-C bonds (one sp^2-sp^3 bond, one $sp-sp^3$ bond) were formed via a sequential palladium carbene migratory insertion and transmetalation process.8 We also reported palladiumcatalvzed cross-coupling reactions of N-(2-iodobenzyl)anilines with α_{β} -unsaturated tosylhydrazones to afford isoindolines.⁹ In this reaction, a C-C bond and a C-N bond were formed via carbene migratory insertion and C-N bond formation. Recently, we demonstrated that stabilized carbon nucleophiles can also be used as partners in cross-coupling with α . β -unsaturated N-tosylhydrazones (Scheme 1, eq 2).¹⁰ In connection with the common n^3 -allylpalladium intermediates in Barluenga's and our work, we wish to report a palladium-catalyzed crosscoupling of alkenyl halides, tosylhydrazones and carbon nucleophiles involving a carbopalladation other than β -hydride elimination (Scheme 1, eq 3).

Scheme 1. Combine Pd-Catalyzed Insertion of *N*-Tosylhydrazone and Pd-Catalyzed Cascade Reactions



The three-component reaction was started by employing vinyl iodide **1a**, *N*-tosylhydrazone **2a**, and carbon nucleophiles as substrates in the presence of Pd₂(dba)₃·CHCl₃ (2.5 mol %), PPh₃ (15 mol %), and K₂CO₃ (3.0 equiv) in THF (6 mL) at 70 °C for 2 h. To our delight, the cross-coupling product **3a** was delivered in 37% yield (Table 1, entry 1). Encouraged by this preliminary result, we proceeded to optimize the reaction. First, we investigated the effect of the additive. In order to prevent the undesired β -hydride elimination, 2.0 equiv of LiCl was added; however,

(10) Ye, Y.-Y.; Zhou, P.-X.; Luo, J.-Y.; Zhong, M.-J.; Liang, Y.-M. Chem. Commun. 201310.1039/C3CC45583A.

(11) The halide ion inhibiting the β -H elimination of palladium(II) species, see: (a) Wang, Z.; Zhang, Z.; Lu, X. Organometallics **2000**, 19, 775–780. (b) Zhang, Z.; Lu, X.; Xu, Z.; Zhang, Q.; Han, X. Organometallics **2001**, 20, 3724–3728. (c) Liu, G.; Lu, X. Org. Lett. **2001**, 3, 3879–3882. (d) Aurrecoechea, J. M.; Durana, A.; Pérez, E. J. Org. Chem. **2008**, 73, 3650–3653.

3a was afforded in comparable yield (data not shown in Table 1).¹¹ When 2.0 equiv of BTAC was used as an additive to enhance the solubility of the sodium dimethyl malonate, it was found to be ineffective (data not shown in Table 1). Next, we inspected the stoichiometry of solvent, PPh₃, carbon nucleophiles, and the effect of temperature. Decreasing the amount of solvent led to a slight improvement in yield (Table 1, entry 2). The yield could be further improved when the reaction was carried out at a lower temperature (Table 1. entry 3). Notably, when the PPh₃ loading was added, a better vield was obtained (Table 1, entry 4). In a similar study, Van Vranken and co-workers showed that using excess inexpensive sodium malonate could significantly improve the yield.¹² Indeed, by increasing the amount of malonate, the product yield was increased to 79% (Table 1, entry 5). Then, we went on to screen other reaction parameters, such as different solvents, catalysts, and ligands. Other solvents including DMF, MeCN, dioxane, and toluene were examined (Table 1, entries 6-9). However, they were found to be unfavorable. Further inspection of the reaction conditions revealed that ligand was essential for this reaction (Table 1, entries 10-13). No reaction occurred when Phen or $[HP'Bu_3]BF_4$ was used, and a lower yield was observed when

Table 1. Conditions for the Palladium-Catalyzed Reaction of **1a** and $2a^{\alpha}$



entry	catalyst	ligand	solvent	yield ^{b} (%)
1	$Pd_2(dba)_3 \cdot CHCl_3$	PPh_3^c	THF^d	37^e , f
2	Pd ₂ (dba) ₃ ·CHCl ₃	PPh_3^c	THF	$42^{e,f}$
3	Pd ₂ (dba) ₃ ·CHCl ₃	PPh_3^c	THF	46 ^f
4	Pd ₂ (dba) ₃ ·CHCl ₃	PPh_3	THF	52^{f}
5	$Pd_2(dba)_3 \cdot CHCl_3$	PPh_3	THF	79
6	$Pd_2(dba)_3 \cdot CHCl_3$	PPh_3	toluene	NR
7	$Pd_2(dba)_3 \cdot CHCl_3$	PPh_3	dioxane	22
8	$Pd_2(dba)_3 \cdot CHCl_3$	PPh_3	MeCN	66
9	$Pd_2(dba)_3 \cdot CHCl_3$	PPh_3	DMF	NR
10	$Pd_2(dba)_3 \cdot CHCl_3$	Phen	THF	NR
11	Pd ₂ (dba) ₃ ·CHCl ₃	dppb	THF	61
12	Pd ₂ (dba) ₃ ·CHCl ₃	[HP ^t Bu ₃]BF ₄	THF	NR
13	Pd ₂ (dba) ₃ ·CHCl ₃		THF	16
14	$Pd_2(dba)_3$	PPh_3	THF	70
15	$Pd(PPh_3)_4^g$	PPh_3	THF	74
16	$Pd(OAc)_2^g$	PPh_3	THF	74
17	$Pd(PPh_3)_2Cl_2^g$	PPh_3	THF	71
18		PPh_3	THF	NR

^{*a*}**1a** (0.3 mmol, 1.0 equiv), **2a** (0.675 mmol, 2.25 equiv), carbon nucleophiles (3.6 mmol, 12.0 equiv), $Pd_2(dba)_3 \cdot CHCl_3$ (2.5 mol %), ligand (10 mol %), K_2CO_3 (0.9 mmol, 3.0 equiv), solvent (2 mL), 46 °C, 2 h. ^{*b*} Yield of isolated product. ^{*c*} 15 mol % PPh₃ was employed. ^{*d*} 6 mL of THF was employed. ^{*e*} The reaction was carried at 70 °C. ^{*f*} Carbon nucleophiles (1.8 mmol, 6.0 equiv). ^{*g*} Pd (5 mol %).

⁽⁸⁾ Zhou, L.; Ye, F.; Zhang, Y.; Wang, J. J. Am. Chem. Soc. 2010, 132, 13590–13591.

⁽⁹⁾ Zhou, P.-X.; Luo, J.-Y; Zhao, L.-B.; Ye, Y.-Y.; Liang, Y.-M. *Chem. Commun.* **2013**, *49*, 3254–3256.

⁽¹²⁾ Devine, S. K. J.; Van Vranken, D. L. Org. Lett. 2008, 10, 1909–1911.

the reaction was performed in the absence of PPh₃. Employing dppb as a ligand could not further improve the yield. Various palladium catalysts including Pd(II) and Pd(0) were investigated and showed they are less efficient than Pd₂(dba)₃·CHCl₃ (Table 1, entries 14–17). Finally, the reaction could not proceed without palladium catalysis (Table 1, entry 18).

With the optimized conditions, the scope of this reaction was then explored. For meta- or para-substituted N-tosylhydrazones, both electron-donating and electronwithdrawing groups were tolerated and reacted smoothly with 1a to afford the desired products in moderate to good yields (Scheme 2). For ortho-substituted N-tosylhydrazones, the yield was found to decrease with increasing size of the substituent. When \mathbf{R}^1 is a small methyl group, **3j** is given in comparable yield with **3f**. With a sterically bulky Cl group, the product 31 was obtained in lower yield. Additionally, using naphthyl N-tosylhydrazone as substrate, the reaction worked efficiently to afford 3m in 56% yield. Notably, the halogen group was tolerated under the Pd-catalyzed conditions, which allowed a transition-metal-catalyzed coupling reaction. However, no reaction occurred when alkyl tosylhydrazones were utilized as coupling partners.

Scheme 2. Palladium-Catalyzed Reactions of 1a with 2^a



^{*a*} Reaction conditions: vinyl iodide **1a** (0.3 mmol, 1.0 equiv), *N*-tosylhydrazone **2** (0.675 mmol, 2.25 equiv), sodium malonate (3.6 mmol, 12.0 equiv), $Pd_2(dba)_3 \cdot CHCl_3$ (2.5 mol %), PPh_3 (10 mol %), K_2CO_3 (0.9 mmol, 3.0 equiv), THF (2 mL), 46 °C, 2 h. Yields are calculated on isolated products.

Moreover, the reaction of vinyl iodide 1b with *N*-tosylhydrazone 2a was also investigated, and 3n was obtained in lower yield, with an all-carbon quaternary center. A better result was observed when the reaction was carried out at 55 °C (Scheme 3, eq 1). Addition, vinyl iodide 1c worked well with 2a. However, no reaction occurred when (Z)-2-iodo-3-phenylacrylaldehyde was employed.

The reaction also worked with ethyl cyanoacetate (Scheme 3, eq 3). Dimethyl 2-allylmalonate and dimethyl 2-phenylmalonate were also investigated. To our disappointment, none of desired products was obtained, which is consistent with the previous report.¹²



Scheme 3. Scope of the Palladium-Catalyzed Reactions with 2a

We further studied the different malonates and found: Diethyl malonate gave good yield, and diisopropyl malonate was less efficient. No reaction occurred when ditertbutyl malonate was used (Scheme 4). Those phenomenon may be may be due to the steric hindrance.

Scheme 4. Reactions of Different Malonates with 2a



As is known, *N*-tosylhydrazone 2a is readily formed by the condensation of aldehyde with tosylhydrazide (Scheme 5). Thus, we decided to conduct this reaction in a one-pot fashion. Treatment of the benzaldehyde with tosylhydrazide generated 2a in situ, then catalyst, ligand, base, and another cross-coupling partner were added. Indeed, product 3a could be isolated in 76% yield.

A plausible mechanism is proposed as shown in Scheme 6. The cross-coupling starts with oxidative addition of Pd(0)



Scheme 6. Plausible Catalytic Cycle



to the vinyl iodide to afford intermediate **A**, followed by decomposition of the diazo compound **B** (generated in situ from *N*-tosylhydrazone) to give a palladium carbene complex **C**.¹³ Subsequently, migratory insertion of a vinyl group to the carbonic carbon led to the η^1 -allylpalladium complex **D**, which can isomerize to the η^3 -allylpalladium

E. This intermediate is then trapped with sodium malonate to generate product 3^{14}

In conclusion, we have developed a new palladiumcatalyzed three-component coupling of vinyl iodide, *N*-tosylhydrazone, and carbon nucleophiles, in which two separate C–C bonds were formed. This reaction proceeds via Pd carbene migratory insertion, followed by carbopalladation instead of classic β -H elimination. This reaction further demonstrates the concept of combination Pdcatalyzed insertion of diazo compounds and Pd-catalyzed cascade reactions, which opens a new avenue for the development of more novel Pd-catalyzed transformations.

Acknowledgment. We thank the National Science Foundation (NSF 21072080 and 21272101) and National Basic Research Program of China (973 Program) 2010CB833203 and "111" program of MOE and PCSIRT: IRT1138 for financial support.

Supporting Information Available. Detailed experimental procedure and copies of ¹H NMR and ¹³C NMR spectra of all compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

⁽¹³⁾ For early studies on a palladium-catalyzed reaction of *N*-tosylhydrazones, see: (a) Barluenga, J.; Valdés, C. *Angew. Chem., Int. Ed.* **2011**, *50*, 7486–7500. (b) Franssen, N. M. G.; Walters, A. J. C.; Reek, J. N. H.; de Bruin, B. *Catal. Sci. Technol.* **2011**, *1*, 153–165. (c) Shao, Z.; Zhang, H. *Chem. Soc. Rev* **2012**, *41*, 560–572. (d) Xiao, Q.; Zhang, Y.; Wang, J. *Acc. Chem. Res.* **2013**, *46*, 236–247.

⁽¹⁴⁾ The intermolecular substitution could occur on either end of the η 3-allylpalladium and it can generate two regioisomeric products. Three were isolated in complete regioselectivities due to the steric hindrance. For a similar example, see: (a) Deardorff, D. R.; Savin, K. A.; Justman, C. J.; Karanjawala, Z. E.; Sheppeck, J. E., II; Hager, D. C.; Aydin, N. J. *Org. Chem.* **1996**, *61*, 3616–3622. (b) Yamada, Y. M. A.; Watanabe, T.; Torii, K.; Uozumi, Y. *Chem. Commun.* **2009**, 5594–5596. (c) Šebesta, R.; Škvorcová, A.; Horváth, B. *Tetrahedron: Asymmetry* **2010**, *21*, 1910–1915.

The authors declare no competing financial interest.