



Neuroprotective polyhydroxypregnane glycosides from *Cynanchum otophyllum*



Zhi-Min Zhao, Zhang-Hua Sun, Mei-Hui Chen, Qiong Liao, Ming Tan, Xin-Wen Zhang, Han-Dong Zhu, Rong-Biao Pi, Sheng Yin*

School of Pharmaceutical Sciences, Sun Yat-sen University, Guangzhou, Guangdong 510006, China

ARTICLE INFO

Article history:

Received 16 March 2013
Received in revised form 22 May 2013
Accepted 20 June 2013
Available online 4 July 2013

Keywords:

Cynanchum otophyllum
Polyhydroxypregnane glycosides
Structure elucidation
Neuroprotective effect

ABSTRACT

Five new polyhydroxypregnane glycosides, namely cynanotosides A–E (**1–5**), together with two known analogues, deacetylmetaplexigenin (**6**) and cynotophylloside H (**7**), were isolated from the roots of *Cynanchum otophyllum*. Their structures were established by spectroscopic methods and acid hydrolysis. The neuroprotective effects of compounds **1–7** against glutamate-, hydrogen peroxide-, and homocysteic acid (HCA)-induced cell death were tested by MTT assay in a hippocampal neuronal cell line HT22. Compounds **1**, **2**, and **7** exhibited protective activity against HCA-induced cell death in a dose-dependent manner ranging from 1 to 30 μM , which may explain the Traditional Chinese Medicine (TCM) use of this plant for the treatment of epilepsy.

© 2013 Elsevier Inc. All rights reserved.

1. Introduction

Cynanchum otophyllum Schneid (Asclepiadaceae), a perennial weed widely distributed in south-west China, is known as “Qingyangshen” in Traditional Chinese Medicine (TCM) for its treatments of epilepsy, rheumatic pain, kidney weakness, and muscle injuries [1]. Previous chemical investigations of this species have resulted in the isolation of a number of pregnane glucosides with the structural variations usually occurring on the substitutions at C-3 and C-12 of the pregnane core [2–5]. Recently, pregnane glycosides have attracted considerable attention for their broad range of bioactivities, such as anti-epileptic activity [6], multidrug-resistance modulating activity [7], immunological activity [8], and antiviral properties [9]. In our continuing search for structurally and biologically interesting metabolites from medical plant resources [10–12], five new pregnane glycosides together with two known steroids (Fig. 1) have been isolated from the roots of *C. otophyllum*. Their structures were established by spectroscopic analyses combined with chemical methods, and three compounds showed neuroprotective effects on homocysteic acid (HCA)-induced cell death screening in the hippocampal neuronal cell line HT22. We report herein the isolation, structural elucidation, and neuroprotective activity of these compounds.

2. Experimental

2.1. General methods

Optical rotation was recorded on a Perkin-Elmer 341 polarimeter. IR spectra were recorded on a FT-IR Tensor37 spectrometer. NMR spectra were recorded on a Bruker AM-400 and Bruker AM-500 spectrometers at 25 °C. ESIMS and HRESIMS were recorded on a Finnigan LC Q^{DECA} instrument. Silica gel (300–400 mesh, Qingdao Haiyang Chemical Co. Ltd.), C₁₈ reverse-phase silica gel (12 nm, S-50 μm , YMC Co. Ltd), Sephadex LH-20 gel (Amersham Biosciences), and Mitsubishi Chemical Industries (MCI) gel (CHP20P, 75–150 μm , Mitsubishi Chemical Industries Ltd.) were used for column chromatography. All solvents used were of analytical grade (Guangzhou Chemical Reagents Company, Ltd.).

2.2. Plant material

The roots and stems of *C. otophyllum* (2 kg) were collected in October 2011 from Yunnan province, PR China, and were identified by Prof You-Kai Xu of Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences. A voucher specimen (accession number: QYS201110) has been deposited at the School of Pharmaceutical Sciences, Sun Yat-sen University.

2.3. Extraction and isolation

The air-dried powder of the roots and stems of *C. otophyllum* (2.0 kg) was extracted with 95% EtOH (3 \times 10 L) at room temp to

* Corresponding author. Tel./fax: +86 20 39943090.

E-mail address: yinsh2@mail.sysu.edu.cn (S. Yin).




Fig. 1. The structures of compounds 1–7 isolated from *C. otophyllum*.

give 120 g of crude extract, which was suspended in H₂O (1 L) and successively partitioned with petroleum ether (PE, 3 × 1 L), EtOAc (3 × 1 L), and *n*-BuOH (3 × 1 L). The EtOAc extract (22 g) was subjected to MCI gel column chromatography (CC) eluted with a MeOH/H₂O gradient (3:7 → 10:0) to afford three fractions (I–III). Fraction I (1.2 g) was purified by silica gel CC (CHCl₃/MeOH, 10:1 → 1:1) then a C₁₈ reverse-phase CC (MeOH/H₂O 6:4 → 10:0) to give **4** (8 mg) and **5** (6 mg). Fraction II (0.8 g) was subjected to a silica gel CC (CHCl/MeOH, 20:1 → 1:1) then a Sephadex LH-20 column (EtOH) afford **3** (12 mg). Fraction III was separated on a C₁₈ reverse-phase CC (MeOH/H₂O 6:4 → 10:0) to give two fractions (Fr. IIIa and Fr. IIIb). Fr. IIIa was subjected to a silica gel CC (CHCl/MeOH, 30:1 → 5:1) to afford **1** (14 mg) and **2** (7 mg). Fr. IIIb was subjected to a Sephadex LH-20 column (EtOH) then a silica gel CC (CHCl₃/MeOH, 20:1) to give **6** (22 mg) and **7** (31 mg).

2.3.1. Cynanotoside A (**1**)

White, amorphous powder; $[\alpha]_D^{25} + 4.4$ (*c* 0.09, CHCl₃); IR (KBr) ν_{\max} 3440, 1709, 1638, 1457, 1375, 1169, 1087, 991 cm⁻¹; ¹H and ¹³C NMR see Tables 1 and 2; HRESIMS *m/z* 807.3970 [M + Na]⁺ (calcd. for C₄₃H₆₀O₁₃Na, 807.3932).

2.3.2. Cynanotoside B (**2**)

White, amorphous powder; $[\alpha]_D^{25} - 20.6$ (*c* 0.19, CHCl₃); IR (KBr) ν_{\max} 3456, 1710, 1642, 1383, 1224, 1168, 1088, 1017, 988 cm⁻¹; ¹H and ¹³C NMR see Tables 1 and 2; HRESIMS *m/z* 787.4268 [M + Na]⁺ (calcd. for C₄₁H₆₄O₁₃Na, 787.4245).

2.3.3. Cynanotoside C (**3**)

White, amorphous powder; $[\alpha]_D^{20} - 24.0$ (*c* 0.05, CHCl₃); IR (KBr) ν_{\max} 3421, 1637, 1447, 1373, 1167, 1068, 1017, 993, 754 cm⁻¹; ¹H and ¹³C NMR see Tables 1 and 2. HRESIMS *m/z* 679.3685 [M + Na]⁺ (calcd. for C₃₄H₅₆O₁₂Na, 679.3669).

2.3.4. Cynanotoside D (**4**)

White, amorphous powder; $[\alpha]_D^{20} - 25.0$ (*c* 0.36, CHCl₃); IR (KBr) ν_{\max} 3441, 1640, 1375, 1161, 1122, 1064, 996, cm⁻¹; ¹H and ¹³C NMR see Tables 1 and 2. HRESIMS *m/z* 809.4300 [M + Na]⁺ (calcd. for C₄₀H₆₆O₁₅Na, 809.4299).

2.3.5. Cynanotoside E (**5**)

White, amorphous powder; $[\alpha]_D^{20} - 43.4$ (*c* 0.35, CHCl₃); IR (KBr) ν_{\max} 3452, 1641, 1411, 1017, cm⁻¹; ¹H and ¹³C NMR see Tables 1 and 2. HRESIMS *m/z* 849.4240 [M + Na]⁺ (calcd. for C₄₂H₆₆O₁₆Na, 849.4249).

2.3.6. Deacetylmetaplexigenin (**6**)

White, amorphous powder; $[\alpha]_D^{20} + 40.0$ (*c* 0.34, MeOH); The optical rotation of **6** was reported for the first time in the current study. The ¹H and ¹³C NMR data (CD₃OD) agreed well with the literature values [13].

2.3.7. Cynotophylloside H (**7**)

White, amorphous powder; $[\alpha]_D^{20} + 36.1$ (*c* 0.05, MeOH), lit [14] $[\alpha]_D^{20} + 30.0$ (*c* 2.2, MeOH); the ¹H and ¹³C NMR data agreed well with the literature values [14].

2.4. Acid hydrolysis of compounds 1–5 and comparison with standard sugars

To a solution of each compound (2 mg) in MeOH (1 mL), 0.2 M H₂SO₄ (1 mL) was added. The solution was kept at 60 °C for 2 h and then diluted with H₂O (2 mL). The solution was neutralized with satd. aq. Ba(OH)₂ and concentrated under vacuum. The residue (the mixture of aglycone and sugars) was subjected to CC (Sephadex LH-20, MeOH) to give fractions of sugars and aglycone. Constituents of each sugar fraction were identified by co-TLC with authentic sugars: cymarose [*R_f* ca. 0.50 in CHCl₃/MeOH (8:1)], diginose [*R_f* ca. 0.46 in CHCl₃/MeOH (8:1)], and digitoxose (*R_f* ca. 0.40 in CHCl₃/MeOH (8:1)). One of the glycosides **13** (5 mg) was hydrolyzed by the above method to afford digitoxose and diginose. The positive optical rotation of digitoxose $[\alpha]_D^{20} = +46.0$ (*c* = 0.1, H₂O) was indicative of a D-configuration ($[\alpha]_D^{20} = +48.4$), while the negative optical rotation of diginose $[\alpha]_D^{20} = -56.2$ (*c* = 0.1, H₂O) suggested a L-configuration ($[\alpha]_D^{20} = -60.6$) [3]. By the same method, digitoxose obtained from **4** and **5** was determined to be the D-isomer, while the cymarose was determined to be the L-form ($[\alpha]_D^{20} = -48.2$).

2.5. Neuroprotective activity assays

HT22 murine hippocampal neuronal cells were maintained in Dulbecco's modified eagle medium (DMEM) supplemented with 10% (v/v) fetal bovine serum (FBS) and incubated at 37 °C under 5% CO₂. To study the protective effect of compounds on neuronal death induced by inducers, glutamate, H₂O₂ and homocystenic acid (HCA), we seeded cells in 96-well plates (10,000 cells/well) and used 6 wells for each treatment group. HT22 cells were pretreated with compounds at different concentrations for 30 min before exposure to inducers unless stated otherwise. The control group was treated with 0.1% (v/v) DMSO as vehicle control. After 24 h, the cell viability was determined with 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay as previously

Table 1
 1H NMR data for compounds **15** in CDCl₃ (*J* in Hz, δ in ppm).

Position	1 ^a	2 ^a	3 ^b	4 ^b	5 ^b
1	1.10, m 1.92, m	1.10, m 1.92, m	1.07, m 1.86, m	1.06, dt (13.4, 3.0) 1.86, m	1.09, m 1.88, m
2	1.63, m 1.92, m	1.64, m 1.93, m	1.93, m 1.68, m	1.67, m 1.93, m	1.94, m 1.62, m
3	3.56, m	3.57, m	3.56, m	3.56, m	3.57, m
4	2.32, m 2.41, m	2.32, m 2.40, m	2.30, m 2.40, dd (13.0, 3.5)	2.30, m 2.41, dd (13.2, 4.0)	2.46, m 2.32, m
6	5.37, brs	5.36, brs	5.37, brs	5.37, brs	5.35, brs
7	2.18, m 2.24, m	2.19, m 2.19, m	2.10, m 2.15, m	2.10, m 2.14, m	2.18, m 2.18, m
9	1.57, dd (11.5, 5.5)	1.53, m	1.43, dd (13.5, 2.0)	1.43, dd (13.2, 2.7)	1.52, dd (11.1, 6.0)
11	1.89, m 1.92, m	1.85, m 1.82, m	1.65, m 2.02, m	1.64, m 2.03, m	1.82, m 1.76, m
12	4.70, dd (10.0, 6.0)	4.56, t (7.5)	3.57, m	3.57, m	4.51, dd (9.1, 6.6)
15	2.00, m	1.97, m	1.74, m 1.86, m	1.74, m 1.86, m	1.93, m 2.86, m
16	2.88, m 1.90, m	2.87, m 1.86, m	1.73, m 1.85, m	1.76, m 1.84, m	1.82, m 1.82, m
18	1.48, s	1.42, s	1.36, s	1.35, s	1.42, s
19	1.14, s	1.13, s	1.18, s	1.18, s	1.12, s
20	–	–	4.06, q (6.5)	4.05, m	–
21	2.20, s	2.20, s	1.16, d (6.5)	1.18, d (5.4)	2.25, s
2'	6.30, d (15.8)	5.52, brs	–	–	1.95, s
3'	7.62, d (15.8)	–	–	–	–
4'	–	2.37, m	–	–	–
5'	7.51, m	1.06, d (7.0)	–	–	–
6'	7.38, m	1.06, d (7.0)	–	–	–
7'	7.39, m	2.17, s	–	–	–
8'	7.38, m	–	–	–	–
9'	7.51, m	–	–	–	–
1''	D-Digit 4.94, brd (9.5)	D-Digit 4.94, brd (9.0)	D-Digit 4.94, brd (9.0)	D-Digit 4.93, dd (8.6, 1.5)	D-Digit 4.92, brd (8.4)
2''	1.86, m 2.12, m	1.86, m 2.13, m	2.05, m 1.77, m	2.11, m 1.72, m	2.10, m 1.72, m
3''	4.13, d (2.5)	4.12, d (3.0)	4.12, d (2.0)	4.24, m	4.23, m
4''	3.30, dd (10.0, 2.0)	3.30, dd (10.1, 3.0)	3.30, dd (9.0, 2.0)	3.21, m	3.21, m
5''	3.80, m	3.78, m	3.78, m	3.82, m	3.80, m
6''	1.25, d (6.0)	1.25, d (6.0)	1.24, d (6.5)	1.24, d (5.6)	1.24, d (7.0)
1'''	D-Dig 5.07, t (2.0)	D-Dig 5.06, t (2.0)	D-Dig 5.06, t (2.0)	D-Digit 4.89, brd (9.6)	D-Digit 4.89, brd (8.4)
2'''	1.88, m 1.97, m	1.88, m 1.96, m	1.90, m 1.85, m	2.15, m 1.75, m	2.13, m 1.75, m
3'''	3.60, m	3.61, m	3.58, m	4.07, m	4.07, m
4'''	3.80, m	3.80, m	3.80, brs	3.24, m	3.24, m
5'''	3.88, q (6.5)	3.88, q (6.0)	3.87, m	3.78, m	3.79, m
6'''	1.30, d (6.5)	1.30, d (6.5)	1.30, d (6.5)	1.22, d (5.8)	1.22, d (6.4)
OMe	3.39, s	3.40, s	3.39, s	–	–
1''''	–	–	–	L-cym 4.91, brd (3.0)	L-cym 4.90, brd (3.0)
2''''	–	–	–	2.31, m 1.80, m	2.31, m 1.80, m
3''''	–	–	–	3.63, m	3.63, m
4''''	–	–	–	3.27, m	3.26, m
5''''	–	–	–	3.85, m	3.85, m
6''''	–	–	–	1.26, d (6.3)	1.26, d (6.1)
OMe	–	–	–	3.42, s	3.42, s

^a Measured in CDCl₃ at 500 MHz.

^b Measured in CDCl₃ at 400 MHz. Digit = digitoxopyranosyl. Dig = diginopyranosyl. cym = cymaropyranosyl.

described [15]. Optical density was measured using a microplate reader (Bio-Tek, USA) at 570 nm and all data were represented as percent of control.

3. Results and discussion

Compound **1**, a white amorphous powder, has a molecular formula of C₄₃H₆₀O₁₃ as determined by HR-ESI-MS at *m/z* 807.3970 [M + Na]⁺ (calcd. 807.3932). The IR spectrum exhibited the absorption bands for hydroxyl (3440 cm⁻¹), ketone (1709 cm⁻¹), and benzene (1638 and 1457 cm⁻¹) functionalities. Lieberman–Burchard

and Keller–Kiliani tests suggested that **1** was a steroidal glycoside. The ¹H and ¹³C NMR spectra of **1** showed the characteristic signals from a cinnamoyl group [δ_{H} 6.30 (1H, d, *J* = 15.8 Hz), 7.62 (1H, d, *J* = 15.8 Hz), 7.51 (2H, m), 7.38 (2H, m), and 7.39 (1H, m)]; δ_{C} 165.8, 117.7, 145.4, 134.3, 128.2 (C × 2), 128.9 (C × 2), and 130.4], a pregnane core [δ_{H} 1.48 (3H, s), 1.14 (3H, s), and 2.20 (3H, s), together with 21 carbon signals], and two sugar units (δ_{H} 4.94 and 5.07; δ_{C} 95.8 and 99.7). Aforementioned information indicated compound **1** was a pregnane glycoside comprising a cinnamoyl group and two sugar units. The pregnane core in **1** was identified as deacetylmetaplexigenin (**6**) by comparison of its ¹H and ¹³C NMR data with those of **6** [13]. In the ¹H-NMR spectrum

Table 2
 ^{13}C NMR data for compounds **15** in CDCl_3 .

Position	1 ^a	2 ^a	3 ^b	4 ^b	5 ^b
1	38.8	38.8	39.0	38.9	38.7
2	28.9	28.9	29.0	29.0	28.8
3	78.0	78.0	77.9	77.9	77.8
4	38.8	38.8	38.8	38.8	38.7
5	140.7	140.7	139.8	139.7	141.0
6	117.7	117.6	118.4	118.4	117.3
7	34.2	34.3	34.6	34.6	34.0
8	74.4	74.3	73.8	73.8	74.5
9	43.7	43.8	43.7	43.7	43.6
10	37.2	37.2	37.0	37.1	37.2
11	24.2	24.3	28.6	28.5	24.2
12	72.7	71.5	70.9	70.8	72.5
13	58.0	57.9	57.8	57.8	57.6
14	88.0	88.0	87.9	87.9	88.2
15	33.1	33.1	33.5	33.4	32.5
16	32.0	31.9	32.6	32.6	32.1
17	91.5	91.5	88.0	88.0	91.7
18	9.4	9.4	10.1	10.1	9.2
19	18.6	18.6	18.4	18.4	18.7
20	209.1	208.8	72.5	72.5	209.4
21	27.4	27.1	16.9	17.0	27.3
1'	165.8	165.9			170.0
2'	117.7	113.0			20.7
3'	145.4	166.8			
4'	134.3	38.2			
5'	128.2	20.8			
6'	128.9	20.9			
7'	130.4	16.5			
8'	128.9				
9'	128.2				
1''	D-Digit	D-Digit	D-Digit	D-Digit	D-Digit
2''	95.8	95.8	95.7	95.7	95.8
3''	37.5	37.5	37.6	37.0	37.0
4''	67.8	67.8	67.8	66.5	66.5
5''	80.9	80.9	80.9	82.6	82.6
6''	68.1	68.1	68.1	68.6	68.6
1'''	D-Dig	D-Dig	D-Dig	D-Digit	D-Digit
2'''	99.7	99.6	99.6	98.3	98.3
3'''	29.6	29.6	29.6	36.7	36.7
4'''	74.3	74.4	74.4	67.4	67.4
5'''	67.4	67.4	67.4	79.3	79.3
6'''	66.7	66.7	66.7	68.0	68.0
OMe	16.9	16.9	17.0	18.2	18.2
	55.6	55.6	55.6	–	–
1''''				L-cym	L-cym
2''''				97.6	97.5
3''''				30.9	30.9
4''''				75.1	75.1
5''''				71.9	71.9
6''''				65.9	65.9
OMe				17.8	17.8
				56.3	56.3

^a Measured in CDCl_3 at 125 MHz.

^b Measured in CDCl_3 at 100 MHz. Digit = digitoxopyranosyl. Dig = diginopyranosyl. cym = cymaropyranosyl.

the observation of an anomeric H-atom at δ_{H} 4.94 (brd, $J = 9.5$ Hz), a CH at δ_{H} 3.30 (dd, $J = 10.0, 2.0$ Hz) and a secondary Me at δ_{H} 1.25 ($d = 6.0$ Hz) indicated a β -digitoxopyranose sugar unit, while the characteristic carbon signals at δ_{C} 99.7, 29.6, 74.3, 67.4, 66.7, 16.9, and 55.6 suggested that the other sugar unit was an α -diginopyranose [3]. This was further confirmed by TLC comparison of the acidic hydrolyzates of **1** with standard sugar samples. The absolute configurations of β -digitoxopyranose and α -diginopyranose were assigned as D and L, respectively, by comparison of their optical rotation with those of authentic sugars. Detailed 2D analysis fulfilled the connections among cinnamoyl, sugars, and deacetylmetaplexigenin moieties (Fig. 2). HMBC correlation from an oxymethine (4.70, dd, $J = 10.0, 6.0$ Hz, H-12) to a carbonyl at 165.8 (C-1') located the cinnamoyl group at C-12. The digitoxopyr-




Fig. 2. Selected ^1H - ^1H COSY (—) and HMBC (---) correlations of **1**.

anose was linked to C-3 by HMBC correlation of H-1''/C-3, which caused the severely downfield shifted carbon signal of C-3 (δ_{C} 78.0) with respect to the corresponding signal in **6** (δ_{C} 71.6). The sugar sequence was established as α -L-diginopyranosyl-(1 \rightarrow 4)- β -D-digitoxopyranose by HMBC correlations of H-1'''/C-4''. Thus the structure of **1** was determined as depicted and given a name cyanotoside A.

Compound **2** had a molecular formula $\text{C}_{41}\text{H}_{64}\text{O}_{13}$ as revealed by the HR-ESI-MS at m/z 787.4268 [$\text{M} + \text{Na}$]⁺ (calcd. 787.4245). The NMR spectra of **2** was very similar to those of **1**, except for the presence of an ikemaoyl group [δ_{H} 5.52 (1H, brs), 2.37 (1H, m), 1.06 (6H, d, $J = 7.0$ Hz), 2.17 (3H, s); δ_{C} 165.9, 113.0, 166.8, 38.2, 20.8, 20.9, 16.5] in **2** instead of a cinnamoyl group in **1**. HMBC correlation from an oxymethine [δ_{H} 4.56 (t, $J = 7.5$ Hz, H-12)] to the carbonyl at 165.9 (C-1') located the ikemaoyl group at C-12. Comparison of the optical rotation of the sugars obtained from the acid hydrolysate of **2** with those of authentic sugar samples further confirmed the presence of an α -L-diginopyranose and a β -D-digitoxopyranose in **2**. Thus the structure of **2** was determined as depicted and given the trivial name cyanotoside B.

Compound **3** was assigned the molecular formula $\text{C}_{34}\text{H}_{56}\text{O}_{12}$ on the basis of HR-ESI-MS at m/z 679.3685 [$\text{M} + \text{Na}$]⁺ (calcd. 679.3669). The NMR spectra of **3** bore a resemblance to those of **1**, with the notable differences being the absence of the signals for a cinnamoyl and a ketone groups and the presence of a doublet methyl (δ_{H} 1.16, d, $J = 6.5$ Hz) and an additional oxymethine (δ_{H} 4.06, q, $J = 6.5$ Hz; δ_{C} 72.5). This implied that the aglycone of **3** was probably sarcostin, a C-20 reduced derivative of deacetylmetaplexigenin (**6**). Comparison of the 1D NMR data of **3** with those of sarcostin confirmed the presence of sarcostin aglycone [8]. The assignment of ^1H and ^{13}C NMR signals of **3** was achieved by detailed 2D NMR analysis. The absolute configurations of the sugar units in **3** were confirmed as α -L-diginopyranose and a β -D-digitoxopyranose using the same methods as described in **1** and **2**. Compound **3** was given the trivial name cyanotoside C.

The molecular formula of compound **4** was determined to be $\text{C}_{40}\text{H}_{66}\text{O}_{15}$ by the HR-ESI-MS at m/z 809.4300 [$\text{M} + \text{Na}$]⁺ (calcd. 809.4299). The NMR data of **4** showed the presence of three sugar units and a sarcostin aglycone. Characteristic signals of δ_{H} 4.93 (dd, $J = 8.6, 1.5$ Hz), 4.89 (d, $J = 9.6$ Hz); δ_{C} 95.7, 98.3, 37.0 (CH_2), and 36.7 (CH_2) in 1D NMR spectra indicated the presence of two β -digitoxopyranose. The third sugar unit was deduced to be α -cymaropyranose by diagnostic signals at δ_{H} 4.91 (d, $J = 3.0$); δ_{C} 97.6, and 30.9 (CH_2) in 1D NMR spectra [3]. Interpretation of the 2D-NMR data (^1H - ^1H -COSY, HMQC, HMBC, and NOESY) not only confirmed the presence of a three-sugar unit at C-3 but also established the sugar sequence as 3- O - α -cymaropyranosyl-(1 \rightarrow 4)- β -digitoxopyranosyl-(1 \rightarrow 4)- β -digitoxopyranoside. Particularly, the HMBC correlations of H-1''''/C-4''' and H-1'''/C-4'' suggested the connection of the three sugars via two (1 \rightarrow 4) linkages. The absolute configurations of the digitoxopyranoses and cymaropyranose were determined as D and L, respectively, by using the same methods described above. The structure of **4** was thus determined as depicted and given the trivial name cyanotoside D.




Fig. 3. Neuroprotective effects of compounds 1–7 against HCA-induced cell death in mice hippocampal HT22 cells. (A) Compounds 1, 2, and 7 exerted slightly beneficial effects against HCA-induced cell death at 10 μM . (B) Compounds 1, 2, and 7 dose-dependently prevented HCA-induced cell death. * $p < 0.05$, ** $p < 0.01$ vs. control group (CT).

Compound 5 exhibited the molecular formula $\text{C}_{42}\text{H}_{66}\text{O}_{16}$ based on the HR-ESI-MS at m/z 849.4240 ($[\text{M} + \text{Na}]^+$ (calcd. 849.4249). The ^1H - and ^{13}C -NMR spectra of 5 (Tables 1 and 2) indicated that it was a triglycoside. The aglycone moiety of 5 shared a high similarity with those of 6 except for the presence of an additional acetyl group [δ_{H} 1.95 (3H, s); δ_{C} 20.7 and 170.0]. HMBC correlations from an oxymethine (δ_{H} 4.51, H-12) to the carbonyl (δ_{C} 170.0) linked the acetyl group to C-12. Three deoxysugar units in 5 were characterized by NMR signals at δ_{H} 4.92 (brd, $J = 8.4$ Hz), 4.89 (brd, $J = 8.4$ Hz), and 4.90 (brd, $J = 3.0$ Hz); δ_{C} 95.8, 98.3, and 97.5, which were almost identical to those in 4, indicating that 5 possessed a 3- O - α -cymaropyranosyl-(1 \rightarrow 4)- β -digitoxopyranosyl-(1 \rightarrow 4)- β -digitoxopyranoside sugar sequence. The sugar moiety was linked to C-3 by HMBC correlation from H-1" to C-3. The absolute configurations of the digitoxopyranoses and cymaropyranose were determined as D and L, respectively, by using the same methods described above. Detailed 2D analysis allowed the full assignments of 1D NMR data of 5. Thus 5 was determined as depicted and given the trivial name cyanotoside E.

The known compounds deacetylmetaplexigenin (6) [13] and cynotophylloside H (7) [14] were identified by comparison of their NMR data with those in literature. A survey of analogous glycosides from the Asclepiadaceae family suggested that all the β -configured 2,6-dideoxysugars have the D-configuration, while the α -configured sugars are L-sugars. In addition, C-2 of a 2-deoxysugar (cymarose, digitoxose, or diginose) that possesses an α -L-configuration usually appears in the ^{13}C -NMR spectrum at ca. 32.0 ppm or less, while that of a β -D-configured 2-deoxysugar normally resonates at a lower field with a chemical shift larger than 34.0 ppm [3].

C. otophyllum has been widely used as a treatment for epilepsy in Traditional Chinese Medicine [1]. Epilepsy is a highly prevalent serious brain disorder, and oxidative stress is considered as a contributing factor to the onset and evolution of this disease [16,17]. To investigate the potential chemistry related to the anti-epilepsy usage of this plant, we examined compounds 17 in three oxidative stress models induced by glutamate, H_2O_2 , and homocysteic acid (HCA), respectively, using MTT assay in a hippocampal neuronal cell line HT22. Compounds 17 failed to reverse the decrease of cell viability caused by glutamate- and H_2O_2 -induced cell death, while 1, 2, and 7 exhibited slightly beneficial effects on HCA-induced cell death at 10 μM (Fig. 3A). To verify the protection effect of 1, 2, and 7 on HCA model, we increased the maximum concentration of these compounds to 30 μM in a reset testing, in which 1, 2, and 7 showed significant dose-dependent protection to HCA-induced cell death ranging from 1 to 30 μM (Fig. 3B). HCA-induced model leads to the death of neurons by depletion of glutathione, the cells major intracellular antioxidant. Moreover, HCA is also considered to be related to NMDA-independent epilepsy in human being [18] and used to establish epilepsy models in immature rats [19].

Thus, the protective effect of 1, 2, and 7 on this model may explain the TCM use of this plant for the treatment of epilepsy. However, the exact mechanisms and detailed connections between this model and epilepsy require further investigation.

Acknowledgements

We gratefully acknowledge financial support for this project from National Natural Science Foundation of China (Nos. 81102339, 81102782) and Natural Science Foundation of Guangdong Province (No. S2011040002429).

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.steroids.2013.06.007>.

References

- [1] Jiangsu New Medical College. The encyclopedia of traditional Chinese medicine. 2nd ed. Shanghai: Shanghai Science and Technology Press; 1985 [p. 1238].
- [2] Ma XX, Wang D, Zhang YJ, Yang CR. Identification of new qingyangshengenin and caudatin glycosides from the roots of *Cynanchum otophyllum*. Steroids 2011;76:10039.
- [3] Ma LF, Shan WG, Zhan ZJ. Polyhydroxypregnane glycosides from the roots of *Cynanchum otophyllum*. Helv Chim Acta 2011;94:227282.
- [4] Ma XX, Jiang FT, Yang QX, Liu XH, Zhang YJ, Yang CR. New pregnane glycosides from the roots of *Cynanchum otophyllum*. Steroids 2007;72:77886.
- [5] Zhao YB, He HP, Lu CH, Mu QZ, Shen YM, Hao XJ. C21 steroidal glycosides of seven sugar residues from *Cynanchum otophyllum*. Steroids 2006;71:93541.
- [6] Mu Q, Lu J, Zhou Q. Two new antiepilepsy compounds – otophyllosides A and B. Sci Sin Ser B 1986;29:295301.
- [7] Hwang BY, Kim SE, Kim YH, Kim HS, Hong YS, Ro JS, et al. Pregnane glycoside multidrugresistance modulators from *Cynanchum wilfordii*. J Nat Prod 1999;62:6403.
- [8] Li XY, Sun HX, Ye YP, Chen FY, Tu J, Pan YJ. Four new immunomodulating steroidal glycosides from the stems of *Stephanotis mucronata*. Steroids 2006;71:68390.
- [9] Li YM, Wang LH, Li SL, Chen XY, Shen YM, Zhang ZK, et al. Seco-pregnane steroids target the subgenomic RNA of alphavirus-like RNA viruses. PNAS 2007;104:80838.
- [10] Han QH, Liu X, Yao WQ, Cheng ZB, Lin TT, Song C, et al. Unusual 9,19:24,32-dicyclotetracyclic triterpenoids from *Lygodium japonicum*. Planta Med 2012;78:19715.
- [11] Lan WJ, Wang J, Guo YQ, Yin S. Oropheayunnol, an unusual 22,23-epoxy apotirucallane triterpenoid from *Orophea yunnanensis*. Nat Prod Commun 2012;7:4956.
- [12] Han QH, Wang DM, Cheng ZB, Yang X, Xu XJ, Wang J, et al. Chemical constituents from the leaves and twigs of *Syzygium tetragonum* wall. Biochem Sys Ecol 2012;41:3–5.
- [13] Takashi Y, Koji H, Hiroshi M, Mamoru I, Kazuhiro M. Carbon-13 nuclear magnetic resonance spectroscopy of C,D-cis polyoxypregnanes. Tetrahedron Lett 1973;37:352730.
- [14] Shan WG, Liu X, Ma LF, Zhan ZJ. New polyhydroxypregnane glycosides from *Cynanchum otophyllum*. J Chem Res 2012;1:38–40.
- [15] Huang YJ, Jin MH, Pi RB, Zhang JJ, Chen MH, Ouyang Y, et al. Protective effects of caffeic acid and caffeic acid phenethyl ester against acrolein-induced

- neurotoxicity in HT22 mouse hippocampal cells. *Neurosci Lett* 2013;535:14651.
- [16] Aguiar CCT, Almeida AB, Araújo PVP, Abreu RNDC, Chaves EMC, Vale OC, et al. Oxidative stress and epilepsy: literature review. *Oxid Med Cell Longev* 2012;2012:112.
- [17] Sudha K, Rao AV, Rao A. Oxidative stress and antioxidants in epilepsy. *Clin Chim Acta* 2001;303:19–24.
- [18] Turski WA. Homocysteic acid: convulsant action of stereoisomers in mice. *Brain Res* 1989;479:3713.
- [19] Folbergrová J, Druga R, Haugvicová R, Mares P, Otáhal J. Anticonvulsant and neuroprotective effect of (S)-3,4-dicarboxyphenylglycine against seizures induced in immature rats by homocysteic acid. *Neuropharm* 2008;54:66575.