


# Rhodium-Catalyzed Asymmetric Tandem Cyclization for Efficient and Rapid Access to Underexplored Heterocyclic Tertiary Allylic Alcohols Containing a Tetrasubstituted Olefin

Yi Li and Ming-Hua Xu\*

State key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 555 Zuchongzhi Road, Shanghai 201203, China

**S** Supporting Information

**ABSTRACT:** The first Rh-catalyzed asymmetric tandem cyclization of nitrogen- or oxygen-bridged 5-alkynones with arylboronic acids was achieved. The simple catalytic system involving a rhodium(I) complex with readily available chiral BINAP ligand promotes the reaction to proceed in a highly stereocontrolled manner. This protocol provides a very reliable and practical access to a variety of chiral heterocyclic tertiary allylic alcohols possessing a tetrasubstituted carbon stereocenter and an all-carbon tetrasubstituted olefin functionality in good yields with great enantioselectivities up to 99% ee.




The transition-metal-catalyzed tandem cyclization reactions of 5-alkyne-1-carbonyl derivatives provide a powerful, straightforward, and atom-economical approach to diverse five-membered cyclic allyl alcohols with an exocyclic olefin, which are employed widely in organic and medicinal chemistry.<sup>1</sup> Among them, nitrogen- or oxygen-linked 5-alkyne-1-carbonyl compounds are particularly important starting materials for the construction of corresponding heterocyclic building blocks, such as tetrahydrofuran or tetrahydropyrrole derivatives. Despite the great progress made in developing efficient transition-metal catalysts including nickel,<sup>2</sup> rhodium,<sup>3</sup> palladium,<sup>4</sup> titanium,<sup>5</sup> or ruthenium<sup>6</sup> complexes, surprisingly, related asymmetric processes to give more valuable enantiomerically enriched products remain an understudied and elusive topic in the literature.<sup>7–9</sup> Rhodium-catalyzed asymmetric hydrogenation of heteroatom-linked acetylenic aldehydes disclosed by Krische in 2006 represents the first successful example of enantioselective cyclization (Scheme 1a).<sup>7a,b</sup> Furthermore, there are two reports of rhodium-catalyzed asymmetric reductive cyclization of similar 5-alkynals with the chelating heteroatom-substituted acetaldehydes or acyl phosphonates by Tanaka, nicely

complementing that obtained by using hydrogen as reducing agent (Scheme 1a).<sup>7c,d</sup>

As for enantioselective cyclization employing nitrogen- or oxygen-linked 5-alkynyl ketones to afford chiral cyclic tertiary allylic alcohols, it is less explored mainly due to the decreased carbonyl electrophilicity and difficulty of stereocontrol (regio- and enantiocontrol). To our knowledge, only a single example of using trimethylsilyl-substituted NTs-tethered 5-alkynone as substrate was reported under palladium catalysis, giving moderate enantiocontrol (82% ee).<sup>7e,8</sup> Alternatively, Hayashi and Murakami have demonstrated that rhodium-catalyzed asymmetric arylation cyclization of 5-alkyne-1-carbonyls (X = C(CH<sub>2</sub>OBn)<sub>2</sub>, C(CO<sub>2</sub>Me)<sub>2</sub> for substrate in Scheme 1a) with stable, commercially available arylboronic acids is an attractive route to carbocyclic allylic alcohols.<sup>9</sup> However, these processes are not equally applicable to nitrogen- or oxygen-linked substrates,<sup>10</sup> and it was assumed that the malonate tether contributes to the high regioselectivity of initial 1,2-addition across alkyne. Therefore, the development of an efficient catalytic system is desirable. We describe herein a highly enantioselective synthesis of heterocyclic tertiary allylic alcohols with an all-carbon tetrasubstituted olefin<sup>11</sup> functionality via arylation cyclization promoted by a Rh(I)/BINAP complex (Scheme 1b).

Our initial survey commenced with evaluation of representative chiral phosphoramidite ligand L1 and phosphine ligands L2–L3 in arylation cyclization of ether-linked 5-alkynone **1a** with (4-methoxyphenyl)boronic acid **2a** in the presence of 2.5 mol % of [Rh(COE)<sub>2</sub>Cl]<sub>2</sub> and KOH (30 mol %) in toluene/H<sub>2</sub>O (10:1) at 60 °C. Intriguingly, the reactions proceeded smoothly to afford the expected cyclized product **3a** (Table 1, entries 1–3). Of these examined, the commercially available

**Scheme 1.** Tandem Cyclization Protocol To Access Chiral Heterocyclic Allylic Alcohols



Received: April 4, 2014

Published: April 28, 2014

**Table 1. Optimization of Reaction Conditions for Rhodium-Catalyzed Asymmetric Arylative Cyclization of 1a with 2a<sup>a</sup>**

Reaction scheme for Table 1: 1a + 2a  $\xrightarrow[\text{base, solvent, T}]{[\text{Rh}(\text{COE})_2\text{Cl}]_2 (2.5 \text{ mol } \%)}$  3a

Ligands L1–L8:

- L1: (R)-MonoPhos
- L2: (R)-MOP
- L3: (R)-BINAP
- L4: (R)-Segphos
- L5: (R)-Sunphos
- L6: (R)-tolBINAP
- L7: (R)-xyBINAP
- L8: (R)-DTBM-Segphos

Ar = 4-MeC<sub>6</sub>H<sub>4</sub> (L6), Ar = 3,5-Me<sub>2</sub>C<sub>6</sub>H<sub>3</sub> (L7), Ar = 2,4,6-Me<sub>3</sub>C<sub>6</sub>H<sub>2</sub> (L8)

entry	L	base	temp (°C)	solvent	yield <sup>b</sup> (%)	ee <sup>c</sup> (%)
1	L1	KOH	60	toluene	83	58
2	L2	KOH	60	toluene	70	51
3	L3	KOH	60	toluene	86	94
4	L4	KOH	60	toluene	75	92
5	L5	KOH	60	toluene	74	93
6	L6	KOH	60	toluene	61	88
7	L7	KOH	60	toluene	41	11
8	L8	KOH	60	toluene	<5	
9	L3	KOH	40	toluene	87	97
10	L3	KOH	RT	toluene	78	97
11	L3	KOH	40	dioxane	37	88
12	L3	KOH	40	CH <sub>2</sub> Cl <sub>2</sub>	77	91
13	L3	KOH	40	MeOH	60	64
14	L3	K <sub>2</sub> CO <sub>3</sub>	40	toluene	74	97
15	L3	KF	40	toluene	77	96

<sup>a</sup>The reaction was performed with 1a (0.1 mmol), 2a (0.3 mmol), [Rh(COE)<sub>2</sub>Cl]<sub>2</sub> (2.5 mol %), L (10 mol %), and base (30 mol %) in 2 mL of solvent/H<sub>2</sub>O (10:1). COE = cyclooctene. <sup>b</sup>Isolated yields.

<sup>c</sup>Determined by HPLC analysis using a chiral stationary phase.

(R)-BINAP L3 (10 mol %) exhibited the greatest preference both in terms of catalytic activity and enantiocontrol (86% yield and 94% ee) (entry 3). To attain higher reactivity and enantioselectivity, a series of biaryl-type bisphosphine ligands L4–L8 that contain different chiral backbones or sterically hindered aromatic substituents on the phosphorus were subsequently tested (entries 4–8). Unfortunately, no superior results were obtained. In all cases, the bisphosphine ligands were either less effective or altogether inactive. Following up on the encouraging results obtained with (R)-BINAP, effects related to the reaction temperature, solvent, and base were carefully examined. As revealed, it was found that lowering the reaction temperature to 40 °C was beneficial for the improvement of enantioselectivity (97% ee) without loss of reactivity (entry 9). However, at room temperature, a small erosion of yield was observed (entry 10). In addition, the investigation of various solvents including dioxane, CH<sub>2</sub>Cl<sub>2</sub>, and MeOH suggested that the use of toluene was the best choice (entries 9 and 11–13). Further efforts to improve the reaction yield and ee by tuning the effect of base turned out to be unhelpful (entries 14 and 15).

With the optimal reaction conditions in hand, we sought to evaluate the scope and generality of the reaction. As summarized in Table 2, a wide variety of ether-linked 5-

**Table 2. Rh/(R)-BINAP-Catalyzed Asymmetric Arylative Cyclization of 1<sup>a</sup>**

Reaction scheme for Table 2: 1 + 2  $\xrightarrow[\text{toluene/H}_2\text{O} (10/1), 40^\circ\text{C}]{[\text{Rh}(\text{COE})_2\text{Cl}]_2 (2.5 \text{ mol } \%), (\text{R})\text{-BINAP} (\text{L3}, 10 \text{ mol } \%), \text{KOH} (30 \text{ mol } \%)}$  3

1a: R = Me, Ar<sup>1</sup> = C<sub>6</sub>H<sub>5</sub>  
 1b: R = Me, Ar<sup>1</sup> = 4-MeOC<sub>6</sub>H<sub>4</sub>  
 1c: R = Me, Ar<sup>1</sup> = 4-BrC<sub>6</sub>H<sub>4</sub>  
 1d: R = Me, Ar<sup>1</sup> = 4-ClC<sub>6</sub>H<sub>4</sub>  
 1e: R = Me, Ar<sup>1</sup> = 3-ClC<sub>6</sub>H<sub>4</sub>  
 1f: R = Me, Ar<sup>1</sup> = 2-Naphthyl  
 1g: R = C<sub>2</sub>H<sub>5</sub>, Ar<sup>1</sup> = C<sub>6</sub>H<sub>5</sub>  
 1h: R = C<sub>6</sub>H<sub>13</sub>, Ar<sup>1</sup> = C<sub>6</sub>H<sub>5</sub>  
 1i: R = H, Ar<sup>1</sup> = C<sub>6</sub>H<sub>5</sub>

entry	1	Ar <sup>2</sup>	3	yield <sup>b</sup> (%)	ee <sup>c</sup> (%)
1	1a	4-OMeC <sub>6</sub> H <sub>4</sub>	3a	87	97
2 <sup>d</sup>	1a	4-MeC <sub>6</sub> H <sub>4</sub>	3b	73	92
3	1a	4- <sup>t</sup> BuC <sub>6</sub> H <sub>4</sub>	3c	70	97
4 <sup>d</sup>	1a	C <sub>6</sub> H <sub>5</sub>	3d	75	91
5	1a	4-ClC <sub>6</sub> H <sub>4</sub>	3e	72	94
6	1a	4-FC <sub>6</sub> H <sub>4</sub>	3f	76	96
7	1a	4-BrC <sub>6</sub> H <sub>4</sub>	3g	78	94
8	1a	2-naphthyl	3h	88	96
9 <sup>d</sup>	1a	3-MeC <sub>6</sub> H <sub>4</sub>	3i	78	90
10 <sup>d</sup>	1a	3-FC <sub>6</sub> H <sub>4</sub>	3j	70	91
11	1b	4-OMeC <sub>6</sub> H <sub>4</sub>	3k	82	96
12	1b	4-FC <sub>6</sub> H <sub>4</sub>	3l	70	94
13	1c	4-OMeC <sub>6</sub> H <sub>4</sub>	3m	80	99
14	1c	4-FC <sub>6</sub> H <sub>4</sub>	3n	79	97
15	1d	4-OMeC <sub>6</sub> H <sub>4</sub>	3o	85	97
16	1d	4-FC <sub>6</sub> H <sub>4</sub>	3p	83	97
17	1e	4-OMeC <sub>6</sub> H <sub>4</sub>	3q	80	98
18	1e	4-FC <sub>6</sub> H <sub>4</sub>	3r	78	98
19	1f	4-OMeC <sub>6</sub> H <sub>4</sub>	3s	72	98
20	1f	4-FC <sub>6</sub> H <sub>4</sub>	3t	72	98
21	1g	4-OMeC <sub>6</sub> H <sub>4</sub>	3u	84	70
22	1h	4-OMeC <sub>6</sub> H <sub>4</sub>	3v	77	65
23	1i	4-OMeC <sub>6</sub> H <sub>4</sub>	3w	68	96
24	1i	4-FC <sub>6</sub> H <sub>4</sub>	3x	49	94

<sup>a</sup>The reaction was performed with 1 (0.1 mmol), 2 (0.3 mmol), [Rh(COE)<sub>2</sub>Cl]<sub>2</sub> (2.5 mol %), (R)-BINAP (L3, 10 mol %), and KOH (30 mol %) in 2 mL of toluene/H<sub>2</sub>O (10:1) at 40 °C. <sup>b</sup>Isolated yields. <sup>c</sup>Determined by HPLC analysis using a chiral stationary phase; see the Supporting Information for details. <sup>d</sup>The reaction was performed at 60 °C.

alkynones 1 bearing different substituents on the aromatic ring attached to carbonyl group and/or at the alkyne terminus were successfully reacted with various arylboronic acids 2, providing the corresponding chiral tertiary allylic alcohols containing a tri- or tetrasubstituted olefin and tetrahydrofuran skeleton 3a–x mostly in good to high yields and excellent enantioselectivities (up to 99% ee). In general, with respect to *para*-substituted and *meta*-substituted arylboronic acids, the electronic properties of the phenyl ring of boronic acids did not appear to affect the reaction yield and enantiomeric excess values (entries 1–10). However, with the more hindered 2-methylphenylboronic acid or 1-naphthylboronic acid, almost no reaction occurred (data not shown). Moreover, ether-linked 5-alkynones 1b–f with either electron-donating or electron-withdrawing groups on the aromatic ring attached to carbonyl group were found to be suitable substrates, as they exhibited high reactivity (70–85%

yield) and led to the formation of highly optically active products **3k–t** (94–99% ee) (entries 11–20). In the cases of 5-alkynones **1g–h** bearing sterically demanding ethyl or *n*-hexyl, respectively, at the alkyne terminus, a distinct decrease in enantioselectivity was observed (entries 21 and 22). When R = Ph, unfortunately, no cyclized product was obtained. Notably, when more challenging 5-alkynone **1i**, possessing acidic hydrogen at the alkyne terminus, was used as the substrate, the tandem reactions proceeded equally smoothly and generated cyclized products with an exocyclic trisubstituted olefin in very high levels of stereocontrol (94–96% ee) (entries 23 and 24).

Fortunately, the structure of the product **3r** was confirmed unambiguously by single-crystal X-ray analysis, and the newly formed chiral center was determined to have the *S* configuration (Figure 1). Assuming an analogous reaction mechanism, the same absolute configurations of the obtained cyclic tertiary allylic alcohol derivatives were assigned.




Figure 1. X-ray crystal structure of (*S*)-**3r**.

Given the remarkable performance of the present catalyst system in asymmetric arylation cyclization of ether-linked 5-alkynones **1** with arylboronic acids **2**, its further applicability was also examined with tosylamide-linked 5-alkynones **4** (Table 3, entries 1–4). To our delight, under the same reaction conditions, reaction involving tosylamide-linked 5-alkynones **4a,b** with several electronically different *para*-substituted arylboronic acids were all found to be successful to afford the corresponding chiral tertiary allylic alcohols **5a–d** containing tetrahydropyrrole skeleton in up to 71% yield with outstanding enantioselectivity (96–99% ee).


Table 3. Rh/(*R*)-BINAP-Catalyzed Asymmetric Arylation Cyclization of **4**<sup>a</sup>

$\text{TsN-CH}_2\text{-C}\equiv\text{C-Me} + \text{Ar}^2\text{B(OH)}_2 \xrightarrow[\text{toluene/H}_2\text{O (10/1), 40 }^\circ\text{C}]{[\text{Rh}(\text{COE})_2\text{Cl}]_2 (2.5 \text{ mol } \%), (\text{R})\text{-BINAP (L3, 10 mol } \%), \text{KOH (30 mol } \%)}$					
entry	Ar <sup>1</sup>	Ar <sup>2</sup>	5	yield <sup>b</sup> (%)	ee <sup>c</sup> (%)
1	C <sub>6</sub> H <sub>5</sub>	4-OMeC <sub>6</sub> H <sub>4</sub>	<b>5a</b>	61	96
2	C <sub>6</sub> H <sub>5</sub>	4-FC <sub>6</sub> H <sub>4</sub>	<b>5b</b>	71	98
3	4-ClC <sub>6</sub> H <sub>4</sub>	4-OMeC <sub>6</sub> H <sub>4</sub>	<b>5c</b>	62	99
4	4-ClC <sub>6</sub> H <sub>4</sub>	4-FC <sub>6</sub> H <sub>4</sub>	<b>5d</b>	70	97

<sup>a</sup>The reaction was performed with **4** (0.1 mmol), **2** (0.3 mmol), [Rh(COE)<sub>2</sub>Cl]<sub>2</sub> (2.5 mol %), (*R*)-BINAP (**L3**, 10 mol %), and KOH (30 mol %) in 2 mL of toluene/H<sub>2</sub>O (10:1) at 40 °C. <sup>b</sup>Isolated yields. <sup>c</sup>Determined by HPLC analysis using a chiral stationary phase; see the Supporting Information for details.

Considering that glycidol containing a tetrahydrofuran skeleton is a unique motif existing in some biologically active compounds, such as peribysin B<sup>12</sup> and milbemycin derivatives,<sup>13</sup> epoxidation of the above obtained chiral tertiary allylic alcohol was attempted. As illustrated in Scheme 2, treatment of

Scheme 2. Synthesis of the Spirocyclic Compound **6**



**3a** with *m*-CPBA in CH<sub>2</sub>Cl<sub>2</sub> at 0 °C for 12 h led to efficient access to the desired product **6** in 74% yield with >20:1 diastereomeric ratio while maintaining high enantioselectivity. The molecular structure as well as the absolute configuration of the oxirane stereocenters were confirmed by single-crystal X-ray crystallography. Thus, a particularly interesting spirocyclic architecture bearing three consecutive quaternary stereocenters that is otherwise difficult-to-access was established in a highly stereoselective fashion.

On the basis of the reaction stereochemical outcome, a plausible catalytic cycle and enantio-differentiation models are proposed in Figure 2. Initiation of the reaction through the transmetalation of hydroxyrhodium(I) **A** with arylboronic acid **2** generates the arylrhodium(I) **B**, which subsequently




Figure 2. Proposed catalytic cycle and models for reaction enantioselectivity.

undergoes regioselective *syn* addition to the carbon–carbon triple bond in **1** to afford the vinylrhodium(I) intermediate **C**. The observed high regioselectivity of addition across alkyne might be attributed to chelation control<sup>14</sup> through the coordination of rhodium to the carbonyl group.<sup>3b,9b</sup> The resulting vinylrhodium(I) intermediate **C** undergoes intramolecular enantioselective nucleophilic addition to the carbonyl group in a *5-exo* mode, forming the rhodium(I) alkoxide intermediate **D**, which is readily hydrolyzed under protic conditions to regenerate **A** with liberation of product **3**. In the cyclization process, the enantiodifferentiating carboration<sup>15</sup> takes place preferentially from the *si* face of the ketone carbonyl through **C II** to avoid the steric congestion with bisphosphine ligand, thereby affording the *S* product.

In summary, we have developed a highly efficient rhodium-catalyzed asymmetric tandem arylation/cyclization of nitrogen- or oxygen-bridged 5-alkynones with arylboronic acids by using a structurally simple, commercially available chiral BINAP as ligand. The reaction showed remarkably broad substrate scope and excellent enantiocontrol (up to 99% ee). Importantly, this method offers a reliable and practical access to a variety of highly enantioenriched tertiary allylic alcohols containing a heterocyclic skeleton, such as tetrahydrofuran or tetrahydropyrrole, which are important units that have been found in numerous natural products and pharmaceutical compounds. Further applications of the methodology are currently underway in our laboratory.

## ■ ASSOCIATED CONTENT

### Supporting Information

Experimental methods and characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [xumh@sim.ac.cn](mailto:xumh@sim.ac.cn).

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We thank the National Natural Science Foundation of China (21325209) and the State key Laboratory of Drug Research for financial support.

## ■ REFERENCES

- (1) For reviews, see: (a) Ojima, I.; Tzamarioudaki, M.; Li, Z.; Donovan, R. J. *Chem. Rev.* **1996**, 96, 635. (b) Montgomery, J. *Acc. Chem. Res.* **2000**, 33, 467. (c) Montgomery, J. *Angew. Chem., Int. Ed.* **2004**, 43, 3890. (d) Moslin, R. M.; Moslin, K. M.; Jamison, T. F. *Chem. Commun.* **2007**, 4441. (e) Miura, T.; Murakami, M. *Chem. Commun.* **2007**, 217.
- (2) Ni-catalyzed reactions: (a) Oblinger, E.; Montgomery, J. J. *Am. Chem. Soc.* **1997**, 119, 9065. (b) Tang, X.-Q.; Montgomery, J. J. *Am. Chem. Soc.* **1999**, 121, 6098. (c) Ni, Y.; Amarasinghe, K. K. D.; Montgomery, J. *Org. Lett.* **2002**, 4, 1743. (d) Saito, N.; Sugimura, Y.; Sato, Y. *Org. Lett.* **2010**, 12, 3494. (e) Baxter, R. D.; Montgomery, J. J. *Am. Chem. Soc.* **2011**, 133, 5728.
- (3) Rh-catalyzed reactions: (a) Ojima, I.; Tzamarioudaki, M.; Tsai, C.-Y. *J. Am. Chem. Soc.* **1994**, 116, 3643. (b) Matsuda, T.; Makino, M.; Murakami, M. *Angew. Chem., Int. Ed.* **2005**, 44, 4608. (c) Miura, T.; Shimada, M.; Murakami, M. *Tetrahedron* **2007**, 63, 6131.
- (4) Pd-catalyzed reactions: (a) Zhao, L.; Lu, X. *Angew. Chem., Int. Ed.* **2002**, 41, 4343. (b) Tsukamoto, H.; Ueno, T.; Kondo, Y. *J. Am. Chem. Soc.* **2006**, 128, 1406. (c) Wang, H.; Han, X.; Lu, X. *Tetrahedron* **2010**, 66, 9129.
- (5) Ti-catalyzed reaction: Crowe, W. E.; Rachita, M. J. *J. Am. Chem. Soc.* **1995**, 117, 6787.
- (6) Ru-catalyzed reaction: Patman, R. L.; Chaulagain, M. R.; Williams, V. M.; Krische, M. J. *J. Am. Chem. Soc.* **2009**, 131, 2066.
- (7) For asymmetric reductive cyclization, see: (a) Rhee, J. U.; Krische, M. J. *J. Am. Chem. Soc.* **2006**, 128, 10674. (b) Rhee, J. U.; Jones, R. A.; Krische, M. J. *Synthesis* **2007**, 3427. (c) Tanaka, R.; Noguchi, K.; Tanaka, K. *J. Am. Chem. Soc.* **2010**, 132, 1238. (d) Masuda, K.; Sakiyama, N.; Tanaka, R.; Noguchi, K.; Tanaka, K. *J. Am. Chem. Soc.* **2011**, 133, 6918. (e) Shen, K.; Han, X.; Lu, X. *Org. Lett.* **2013**, 15, 1732.
- (8) For a report of cationic Pd(II)-catalyzed asymmetric cyclization of arylmethyl 2-alkynoates bearing an electron-deficient alkyne to produce optically active hydroxy lactone, see: Song, J.; Shen, Q.; Xu, F.; Lu, X. *Org. Lett.* **2007**, 9, 2947.
- (9) (a) Shitani, R.; Okamoto, K.; Otomaru, Y.; Ueyama, K.; Hayashi, T. *J. Am. Chem. Soc.* **2005**, 127, 54. (b) Miura, T.; Shimada, M.; Murakami, M. *Synlett* **2005**, 4, 667.
- (10) A single example of O-tethered 5-alkynone cyclization was reported, giving the corresponding racemic tetrahydrofuran product in only 21% yield due to the poor regioselectivity control; see ref 9b.
- (11) Flynn, A. B.; Ogilvie, W. W. *Chem. Rev.* **2007**, 107, 4698.
- (12) Yamada, T.; Iritani, M.; Minoura, K.; Kawai, K.; Numata, A. *Org. Biomol. Chem.* **2004**, 2, 2131.
- (13) Naito, S.; Saito, A.; Furukawa, Y.; Hata, T.; Nakada, Y.; Muramatsu, S.; Ide, J. *J. Antibiot.* **1994**, 47, 812.
- (14) For examples of the chelation-controlled regioselective *syn* addition of arylrhodium(I) to the carbon–carbon triple bond, see: (a) Lautens, M.; Yoshida, M. *Org. Lett.* **2002**, 4, 123 and references cited therein. (b) Lautens, M.; Yoshida, M. *J. Org. Chem.* **2003**, 68, 762 and references cited therein.
- (15) (a) Fagnou, K.; Lautens, M. *Chem. Rev.* **2003**, 103, 169. (b) Hayashi, T.; Yamasaki, K. *Chem. Rev.* **2003**, 103, 2894.