

Completely Regioselective Synthesis of 5- and 6- Amino and Fluoro-hexofuranoses via cyclic sulphates.

José Fuentes*, Manuel Angulo, and M. Angeles Pradera

Dpto de Química Orgánica, Facultad de Química, Universidad de Sevilla, Apartado de Correos No 553, E-41071 Sevilla, Spain.

Received 24 April 1998; accepted 20 July 1998

Abstract: The nucleophilic opening of new and previously described 5,6- and 3,5-glucohexofuranose cyclic sulphates is a regioselective and efficient way to prepare 6- and 5-azido(amino)- and 6- and 5-fluoro-aldofuranose derivatives (D-gluco and L-ido configurations). © 1998 Published by Elsevier Science Ltd. All rights reserved.


The cyclic sulphates of *vic*-diols are compounds which have been known for a long time, but until recently, they have not been widely used in organic synthesis, probably due to the lack of a good method for their preparation. Ten years ago, the catalytic oxidation of cyclic sulphites using ruthenium trichloride with sodium periodate was reported¹ as a very general method for the synthesis of cyclic sulphates. Since then, the scope of the method has been studied, and the interest for the synthetic applications of these compounds has grown continuously. Most of the synthetic uses of the cyclic sulphates described until now are based on nucleophilic displacement reactions. These sulphates are synthetic equivalents of epoxides and cyclic sulphites, with the advantages of a greater reactivity and versatility than cyclic sulphites and, in many cases, than oxiranes.² In spite of this interest, the applications in carbohydrate chemistry are limited,²⁻⁷ especially in the case of fluoroderivatives, about which we have found data on the preparation only of 4-fluoro-D-xylulose⁸ derivatives and 2-deoxy-2-fluoro-D-glucose.⁹

The introduction of an azido function into a sugar molecule is an interesting target because azidosugars play important roles in the synthesis of oligosaccharides,¹⁰ in the treatment of infectious diseases¹¹ and because they can easily be reduced to amino sugars. Recently, 5- and 6-aminosugars are being applied to the preparation of azasugars¹² and to the stereocontrolled syntheses of C-nucleosides.¹³ The introduction of an amino function into the position 5 of aldohexofuranoses is a non-trivial problem as the described methods, based on the reduction of 5-oxime derivatives¹⁴ or on the inversion of the configuration of C-5 derivatives,¹⁵ have a high number of steps, substitution competes with elimination, and/or mixtures of diastereomers, which are difficult to resolve, are formed. These difficulties are especially important in the case of minor configurations, such as L-ido.

At the same time the carbohydrate fluoro derivatives exhibit interesting antileukaemia¹⁶ and anti-retroviral¹⁷ activities. These biological properties are probably due to a modification of the protein-carbohydrate interactions. Substitution of a hydroxyl group by a fluorine atom can provide insight into whether the hydroxyl group removed was involved in critical hydrogen-bond interactions with the protein-binding site. The fluorine atom can act as hydrogen bond acceptor but not as hydrogen bond donor, and in this way can inform about the

role of the removed hydroxyl group. Consequently the synthesis of fluorosugars is a major challenge in carbohydrate chemistry. The described methods to introduce a fluorine atom in the position 6 of glycopyranoses^{18,19} and glycofuranoses¹⁹ have low-medium yields, especially in the second case, where the data are very scarce. Some attempts²⁰ to introduce a fluorine atom in the position 5 of hexofuranoses (only 6-deoxy derivatives) have been carried out, but elimination overwhelmingly predominated on substitution.

We now communicate a new way to prepare 5- and 6- azido, amino, and fluoro sugars starting from previously described³ (**1**, **2**) and new (**3**, **4**) glucofuranose 5,6- and 3,5-cyclic sulphates which were prepared from suitably protected glucofuranose derivatives. Compounds **3** and **4**²¹ were easily prepared in practically quantitative yields through the corresponding cyclic sulphites, which were obtained as diastereomeric mixtures and directly oxidised with $\text{NaIO}_4\text{-RuCl}_3\cdot 3\text{H}_2\text{O}$ following the method described by Sharpless.¹



The nucleophilic ring opening of the cyclic sulphates **1-3** with sodium azide in DMF at 50 °C gave, after elimination of the solvent and treatment with a catalytic amount of 1:10 $\text{H}_2\text{SO}_4\text{:H}_2\text{O}$ in THF, the corresponding 6-azido-6-deoxy-glucofuranose derivative (**5**, $\text{R}^1 = \text{Ac, Bn, Ms}$; $\text{R}^2 = \text{N}_3$) in quantitative yield. As expected, the nucleophilic attack took place on the less hindered primary position in a completely regioselective manner. In the case of **3** no displacement of the mesyloxy group on C-3 was observed either during the described conditions for the opening of the sulphate ring, or on treatment with sodium azide at 105 °C in DMF. Catalytic (10% Pd/C) hydrogenation (r.t, 1 atm of pressure for 50 min) of the 6-deoxy-6-azidoglucofuranose derivatives yielded quantitatively the corresponding 6-amino glucose derivative (**5**, $\text{R}^1 = \text{Ac, Bn, Ms}$; $\text{R}^2 = \text{NH}_2$). In the case of the 3-*O*-benzyl compound, debenzylation took place when a reaction time of 12 h was used, obtaining the 3-*O*-unprotected 6-amino-1,2-*O*-isopropylidene-6-deoxy- α -D-glucofuranose (**5**, $\text{R}^1 = \text{H}$, $\text{R}^2 = \text{NH}_2$).

The nucleophilic opening of the non-vicinal cyclic sulphate **4** with sodium azide, using the same experimental conditions and work-up described above, produced quantitatively, and with 100% of regioselectivity, the 5-azido-5-deoxy-L-ido derivative (**6**, $\text{R}^1 = \text{N}_3$, $\text{R}^2 = \text{Ac}$). When this compound was reduced, simultaneous O-N transacetylation took place, and the 5-acetamido-5-deoxy derivative (**6**, $\text{R}^1 = \text{NHAc}$, $\text{R}^2 = \text{H}$) was isolated. The 5-amino-1,2-*O*-isopropylidene-5-deoxy- β -L-idofuranose (**6**, $\text{R}^1 = \text{NH}_2$, $\text{R}^2 = \text{H}$) could be prepared by deacetylation (90%) of the 6-*O*-acetyl azide (**6**, $\text{R}^1 = \text{N}_3$, $\text{R}^2 = \text{Ac}$) and after reduction (quantitative).

The 6-fluoro-6-deoxy glucofuranoses (**5**, $\text{R}^1 = \text{Ac, Bn, Ms}$; $\text{R}^2 = \text{F}$) were prepared by opening of the sulphate ring of **1-3** using tetraethylammonium fluoride dihydrate as nucleophile in acetone at r.t (2 h). The

yields were 85-90% and the reaction was completely regioselective. The yields became virtually quantitative when anhydrous tetrabutylammonium fluoride was used as fluorinating reagent.

To introduce a fluorine atom in the position 5 of a glucofuranose derivative, we have started from the cyclic sulphate **4**. The reaction has been carried out with several nucleophiles and conditions (see table I) and, although the elimination, with formation of the 4,5 unsaturated derivative, competed with the substitution to obtain the 5-fluorocompound (entries a, b), the amount of the latter was significant. The best yield of the fluoroderivative **6** ($R^1 = F$, $R^2 = H$) was using tris(dimethylamino)sulphur (trimethylsilyl)difluoride (TAS-F) as nucleophile and the conditions of entry c, in this case there was practically no formation of elimination product.

Table 1			
Fluorination reactions on compound 4 .			
Entry	Nucleophile/4 ratio	Conditions	Elimination: Substitution
a	Bu ₄ NF 4:1	DMF 65 °C 15 min	3:1
b	Bu ₄ NF 4:1	DMF 0°C 40 h	3:1.8
c	TAS-F 3:1	CH ₂ Cl ₂ -10°C 24h, and after 35°C 72h	1:100

The azido-, amino-, and fluorosugars²² **5** and **6** were stable compounds, whose analytical and spectroscopic data²³ (IR, ¹H-, and ¹³C-NMR) confirmed the structure. Thus, the IR spectra of the azido derivatives showed a band at $\approx 2104\text{ cm}^{-1}$ for the azido group, and the chemical shift for the resonance of C-6 was $\approx 54.0\text{ ppm}$ for the 6-azido compounds, $\approx 45.0\text{ ppm}$ for the 6-amino compounds, and $\approx 84.5\text{ ppm}$ for the 6-fluoro derivatives. The ¹J_{C-F} coupling constants for **5** ($R_2 = F$) and **6** ($R_1 = F$) were $\approx 170\text{ Hz}$.

In conclusion, we describe a new and efficient method to introduce the azido (amino) and fluoro functions into the positions 5 and 6 of glucofuranose derivatives. As the 1,2-O-isopropylidene group is easy to remove²⁴, the method can give access to pyranoid and furanoid D-glucos and L-idos derivatives. The reactions are experimentally easy and the overall yields are better than in other described methods.^{14,15,18,19}

The scope and limitations of this method are currently under study in our laboratory.

ACKNOWLEDGMENTS

We thank the Dirección General de Investigación Científica y Técnica of Spain for the financial support (grant number PB 94/1440-C02-01) and for the award of a grant to M. A.

REFERENCES AND NOTES

- Gao, Y.; Sharpless, K.B. *J. Am. Chem. Soc.* **1988**, *110*, 7538-7539.
- For reviews see Kolb, H.C.; VanNieuwenhze, M.S.; Sharpless, K.B. *Chem. Rev.* **1994**, *94*, 2483-2547 and Lohray, B.B. *Synthesis* **1992**, 1035-1052.
- Calvo-Flores, F.G.; García-Mendoza, P.; Hernández-Mateo, F.; Isac-García, J.; Santoyo-González, F. *J. Org. Chem.* **1997**, *62*, 3944-3961.
- Vanhessche, P.M.; Sharpless, K.B. *Chem. Eur. J.* **1997**, *3*, 517-522.
- Berridge, M.S.; Franceschini, M.P.; Rosenfeld, E.; Tewson, T.J. *J. Org. Chem.* **1990**, *55*, 1211-1217.
- Oi, R.; Sharpless, K.B. *Tetrahedron Lett.* **1991**, *32*, 999-1002.
- Application to the synthesis of 2,3, and 4-azidoglycopyranoses are reported in Van der Klein, P.A.M.; Filemon, W.; Veeman, G.H.; Van der Marel, G.A.; Van Boom, J.H. *J. Carbohydr. Chem.* **1992**, *11*,

- 837-848. The synthesis of 6-deoxy-D-manno-heptopyranose derivatives is described in Van der Klein, P.A.M.; Van Boom, J.H. *Carbohydr. Res.* **1992**, 224, 193-200. For preparation of episulphides, olefins, and thiosugars see 3. For use as alkylating agent see Gómez, A.M.; Valverde, S.; Fraser Reid, B. *J. Chem. Soc. Chem. Commun.* **1991**, 1207-1208.
8. Vanhessche, K.; Van der Eycken, E.; Vandewalle, M.; Roper, H. *Tetrahedron Lett.* **1990**, 31, 2337-2340.
 9. Tewson, T.J. *J. Org. Chem.* **1983**, 48, 3507-3510.
 10. For a review see Banoub, J.; Boullanger, P.; Lafont, D. *Chem. Rev.* **1992**, 92, 1167-1195.
 11. For a review see Hury, D.M.; Okabe, M. *Chem. Rev.* **1992**, 92, 1745-1768.
 12. Jiménez-Blanco, J.L.; Díaz Pérez, V.M.; Ortiz Mellet, C.; Fuentes, J.; García Fernández, J.M.; Díaz Arribas, J.C.; Cañada, F.J. *Chem. Commun.* **1997**, 1969-1970.
 13. García Fernández, J.M.; Ortiz Mellet, C.; Fuentes, J. *J. Org. Chem.* **1993**, 58, 5192-5199.
 14. Legler, G.; Julich, E. *Carbohydr. Res.* **1984**, 128, 61-72.
 15. Dox, K.; Gaigg, B.; Grassberger, V.; Koblinger, B.; Stutz, A.E. *J. Carbohydr. Chem.* **1990**, 9, 479-499.
 16. Sharma, M.; Bernacki, R.J.; Hillman, J.; Korytnyk, W. *Carbohydr. Res.* **1993**, 240, 85-93.
 17. Raifeld, Y.E.; Niktenko, A.A.; Arshava, B.M.; Mikerin, I.E.; Zilberg, L.L.; Vid, G.Y.; Lang, S.A.; Lee, V.J. *Tetrahedron* **1994**, 50, 8603-8616, and references cited therein.
 18. Mulard, L.A.; Kovac, P.; Glaudemans, C.P.J. *Carbohydr. Res.* **1994**, 259, 157-129 and **1994**, 259, 21-34.
 19. Evelyn, L.; Hall, L.D. *Carbohydr. Res.* **1976**, 47, 285-297.
 20. a) Mori, Y.; Harada, K.; Morishima, N.; Zen, S. *Chem. Pharm. Bull.* **1993**, 41, 755-757. b) Mori, Y.; Morishima, N. *Bull. Chem. Soc. Jpn.* **1994**, 67, 236-241.
 21. Selected data for **4**: $[\alpha]_D^{24} +38^\circ$ (c 1.0, dichloromethane); mp 116-118°C dec; IR ν_{\max} 1753, 1398, 1254, 1204, 1094, 1042, 866 cm^{-1} ; $^1\text{H-NMR}$ (500 MHz, CDCl_3) δ 6.06 (d, 1 H, $J_{1,2}=3.7$, H-1), 5.19 (d, 1 H, $J_{3,4}=2.5$, H-3), 5.00 (t d, 1 H, $J_{4,5}=2.5$, $J_{5,6a}=J_{5,6b}=6.4$, H-5), 4.79 (d, 1 H, H-2), 4.57 (m, 1 H, H-6a), 4.54 (m, 1 H, H-6b), 4.35 (t, 1 H, H-4), 2.13(s, 3 H, COCH_3), 1.53, 1.36 (each s, each 3 H, $\text{C}(\text{CH}_3)_2$); $^{13}\text{C-NMR}$ (125.7 MHz; CDCl_3) δ 169.9 (CO), 113.2 ($\text{C}(\text{CH}_3)_2$), 104.8 (C-1), 85.1 (C-3), 82.9 (C-5), 82.5 (C-2), 71.7 (C-4), 61.5 (C-6), 26.5, 26.0 ($\text{C}(\text{CH}_3)_2$), 20.4 (COCH_3); FABMS m/z 347 (95, $[\text{M}+\text{Na}]^+$). Compound **4** gave satisfactory microanalysis (C,H,N,S).
 22. All are new compounds except 6-azido-3-O-benzyl-6-deoxy-1,2-O-isopropylidene- α -D-glucofuranose which was previously described. [Saeki, H.; Ohki, E. *Chem. Pharm. Bull.* **1968**, 16, 2471].
 23. As examples the structural data of the following compounds are given:
 5-azido-5-deoxy-1,2-O-isopropylidene- β -L-idofuranose had $[\alpha]_D^{24} -57^\circ$ (c 1.0, dichloromethane); mp 72-74°C; IR ν_{\max} 3443, 2104, 1740, 1373, 1260, 1082, 1020 cm^{-1} ; $^1\text{H-NMR}$ (500 MHz, CDCl_3) δ 5.98 (d, 1 H, $J_{1,2}=3.7$, H-1), 4.53 (d, 1 H, H-2), 4.32 (dd, 1 H, $J_{6a,6b}=11.7$, H-6a), 4.22 (m, 1 H, H-3), 4.14 (dd, 1 H, H-6b), 4.13 (dd, 1 H, $J_{3,4}=2.5$, H-4), 3.94 (t d, 1 H, $J_{4,5}=7.6$, $J_{5,6a}=3.4$, $J_{5,6b}=7.6$, H-5), 2.66 (d, 1 H, OH), 2.13(s, 3 H, COCH_3), 1.50, 1.32 (each s, each 3 H, $\text{C}(\text{CH}_3)_2$); $^{13}\text{C-NMR}$ (125.7 MHz; CDCl_3) δ 170.9 (CO), 112.0 ($\text{C}(\text{CH}_3)_2$), 104.5 (C-1), 85.2 (C-2), 79.6 (C-4), 75.2 (C-3), 63.8 (C-6), 59.8 (C-5), 26.7, 26.1 ($\text{C}(\text{CH}_3)_2$), 20.6 (COCH_3); FABMS m/z 310 (100, $[\text{M}+\text{Na}]^+$).
 5-deoxy-5-fluoro-1,2-O-isopropylidene- β -L-idofuranose had $[\alpha]_D^{24} -10^\circ$ (c 0.8, dichloromethane); IR ν_{\max} 3455, 2965, 1742, 1377, 1260, 1084, 1020 cm^{-1} ; $^1\text{H-NMR}$ (500 MHz, CDCl_3) δ 5.94 (d, 1 H, $J_{1,2}=3.7$, H-1), 4.90 (dtd, 1 H, $J_{4,5}=6.3$, $J_{5,6a}=2.6$, $J_{5,6b}=6.3$, $J_{5,F}=49.4$, H-5), 4.46 (m, 1 H, H-2), 4.37 (ddd, 1 H, $J_{6a,F}=27.0$, $J_{6a,6b}=12.7$, H-6a), 4.29 (ddd, 1 H, $J_{6b,F}=23.6$, H-6b), 4.22 (ddd, 1 H, $J_{3,4}=2.9$, $J_{4,F}=17.1$, H-4), 4.21 (m, 1 H, H-3), 2.06(s, 3 H, COCH_3), 1.43, 1.26 (each s, each 3 H, $\text{C}(\text{CH}_3)_2$); $^{13}\text{C-NMR}$ (125.7 MHz; CDCl_3) δ 170.9 (CO), 112.1 ($\text{C}(\text{CH}_3)_2$), 104.8 (C-1), 90.5 (d, $^1J_{5,F}=176$, C-5), 85.3 (C-2), 78.6 (d, $^2J_{4,F}=17.6$, C-4), 75.5 (d, $^3J_{1,F}=5.7$, C-3), 63.6 (d, $^2J_{6,F}=22.6$, C-6), 26.7, 26.2 ($\text{C}(\text{CH}_3)_2$), 20.6 (COCH_3); FABMS m/z 265 (85, $[\text{M}+1]^+$).
 24. See for example Wolfrom, M.L.; Thomas, G.H.S. *Methods Carbohydr. Chem.* **1963**, 11, 32-34 and see also reference 13.