

Tetrahedron Letters 40 (1999) 3705-3708

TETRAHEDRON LETTERS

Synthesis of C2-Symmetric Bis(Cyclic Isothioureas)

As Potent Inhibitors of Glycosidases

Laurence Gauzy, Yves Le Merrer,* Jean-Claude Depezay,

Université René Descartes, Laboratoire de Chimie et Biochimie Pharmacologiques et Toxicologiques, UMR 8601 CNRS, 45, rue des Saints-Pères, 75270 Paris cedex 06, France.

Dominique Damour-Barbalat and Serge Mignani

Rhône-Poulenc Rorer S.A. Centre de Recherche de Vitry-Alfortville, 13, quai Jules Guesde, BP14, 94403 Vitry sur Seine cedex, France.

Received 9 February 1999; accepted 18 March 1999

Abstract : Enantiopure C₂-symmetric *bis*(cyclic isothioureas), considered as potent inhibitors of glycosidases, have been synthesized from D-mannitol. The key step involved a mercuric-catalyzed transformation of a cyclic 1,3-thiazolidine-2-thione into a 2-*N*-tert-butylamino-1,3-thiazoline. © 1999 Elsevier Science Ltd. All rights reserved.

Keywords : Glycosidase, Isothiourea, Thioformamide, Thiazolidinethione, Mercuric chloride.

Glycosidases are involved in many biological processes, such as glycoprotein trimming, catabolism of glycoconjugates and degradation of polysaccharides. As a consequence, specific inhibitors of these enzymes are of great interest as possible therapeutic agents to treat many diseases such as diabete,¹ cancer² or viral infections.³

Most inhibitors of glycosidases described to date are mimics of the supposed transition state oxocarbenium.⁴ Among sugar mimics,⁵ azasugars are structures that are the most widely used. However, we focused our attention to other structures, namely isoureas⁶ and isothioureas⁷ which were already combined with an aminocyclitol or a sugar and revealed to be active against glycosidases. Our goal was to examine the potent inhibitory activity of such compounds in the absence of the sugar-like moiety.

As a matter of fact, we decided to synthesize some cyclized C_2 -symmetric *bis*(isothioureas) from Dmannitol. We report here preliminary results concerning the syntheses of the enantiopure *bis*(2-amino-1,3thiazolines) 1 and 2, as potent inhibitors of glycosidases :



Three different methods for the synthesis of 2-amino-1,3-thiazoline from β -aminoalcohol are depicted in Scheme 1 : 1) reaction with thiocyanate ion, and subsequent cyclization; 2) reaction with an alkyl-, aryl- or

acyl-isothiocyanate, and cyclization; or 3) reaction with carbon disulfide to intermediately form a cyclic 1,3thiazolidine-2-thione, followed by S-methylation and subsequent displacement of the resulting methylthio group with either ammonia or a primary amine. Among those three pathways, only a procedure derived from the last one was successful with our compounds derived from D-mannitol.



Scheme 1 : A = OH or leaving group, R = H or alkyl.

The synthesis of 1 began with the 1,6-diazido compound 5 (Scheme 2), obtained from the commercially available 3,4-O-methylethylidene-D-mannitol 3 by selective tosylation of primary hydroxyl groups with p-toluenesulfonyl chloride (86%),⁸ followed by nucleophilic substitution of the resulting tosylates with sodium azide (75%). Heterogeneous catalytic reduction of 5 in the presence of di-*tert*-butyl dicarbonate⁹ led to the protected 1,6-diamine 6 (83%). Mesylation of the secondary hydroxyl groups followed by deprotection of the amines with trifluoroacetic acid led to the 1,6-diammonium-2,5-dimesylate *bis*-trifluoroacetate 8 in quantitative yield.



Scheme 2 : (a) TsCl, pyridine, 0° C ; (b) NaN₃, DMF, Δ ; (c) H₂, 10% Pd/C, (Boc)₂O, EtOAc, rt, 2h ; (d) MsCl , NEt₃, CH₂Cl₂ ; (e) TFA / CH₂Cl₂, rt, 1h ; (f) CS₂, NEt₃, CH₂Cl₂, rt, 15h ; (g) ^tBuNH₂, NEt₃, HgCl₂, DMF, 80°C, 15h ; (h) concentrated HCl, reflux, 15h

Direct transformation of **8** into the *bis*(1,3-thiazolidine-2-thione) **9** with inversion of configuration at C₂ and C₅ cleanly occurred in presence of carbon disulfide and slow addition of triethylamine (90%). Under these conditions, involving the good electrophile CS₂,¹⁰ nucleophilic addition of the primary amines and

subsequent formation of the thiazolidine rings was much faster than the intramolecular formation of aziridine rings. Although methylation at the exocyclic sulfur atoms of 9 with methyl iodide was easily achieved, all attempts to substitute the thiomethyl groups with ammonia were unsuccessful. To overcome this problem, by analogy with transformation of thiourea or thioamide into guanidine¹¹ or amidine,¹² respectively, direct substitution of the thioxo groups of 9 was examined. As expected, *tert*-butylamine treatment of 9 in DMF in the presence of mercuric dichloride and triethylamine led to the *bis*(2-*N*-*tert*-butylamino-1,3-thiazoline) 10 in quantitative yield. The final acetonide hydrolysis and *N*-*tert*-butyl cleavage were carried out in refluxing concentrated aqueous HCl and led to compound 1 (70%), isolated as its *bis*-chlorhydrate salt.¹³

Furthermore, from the N-Boc protected 2,5-diamine 15 the synthesis of 2, which is a N,S-isomer of position of 1, was efficiently carried out (Scheme 3). The diamine 15 was easily obtained from the 3,4-O-methylethylidene-D-mannitol 3 as follows : selective benzylation of both primary hydroxyl groups through stannylidene activation with Bu₂SnO and treatment with benzyl bromide (86%), then mesylation of the secondary hydroxyl groups and nucleophilic substitution with sodium azide (57%). Heterogeneous catalytic reduction of the diazido compound 13 in the presence of di-*tert*-butyl dicarbonate⁹ led to the protected 2,5-diamine 14 (76%) which was subsequently debenzylated with sodium in liquid ammonia in quantitative yield to afford 15. This *bis*(N-Boc-aminoalcohol) was successively subjected, as above, to mesylation, N-Boc deprotection and cyclization into *bis*(1,3-thiazolidine-2-thione) with carbone disulfide, then HgCl₂-catalyzed transformation into *bis*(2-*N*-tert-butylamino-1,3-thiazoline), prior to the final removal of all protecting groups.¹⁴



Scheme 3: (a) *i*- Bu₂SnO, toluene, Δ , 15h; *ii*- BnBr, ⁿBu₄NI, toluene, 70°C, 15h; (b) MsCl, NEt₃, CH₂Cl₂; (c) NaN₃, DMF, 120°C, 15h; (d) H₂, 10% Pd/C, (Boc)₂O, EtOAc, rt, 2h30; (e) Na / NH₃ liq., 1h; (f) MsCl, NEt₃, CH₂Cl₂; (g) TFA / CH₂Cl₂, rt, 1h; (h) CS₂, NEt₃, CH₂Cl₂, rt, 15h; (i) ^tBuNH₂, NEt₃, HgCl₂, DMF, 80°C, 15h; (j) concentrated HCl, reflux, 15h.

In summary, this study presents an efficient synthetic way to construct enantiopure C_2 -symmetrical bis(2-amino-1,3-thiazolines) from D-mannitol. The key step involves the HgCl₂-catalyzed transformation of a

cyclic 1,3-thiazolidine-2-thione into a 2-*N-tert*-butylamino-1,3-thiazoline. The biological activity of the new compounds described here has been studied; full results will be reported in due course.

Acknowledgements. One of us (LG) thanks Rhône-Poulenc Rorer for a grant.

References and Notes

- (a) Junge, B.; Heiker, F-R.; Kurz, J.; Müller, L.; Schmidt, D.D.; Wünsche, C. Carbohydr. Res. 1984, 128, 235-238. (b) Horii, S.; Fukase, H.; Matsuo, T.; Kameda, Y.; Asano, N.; Matsui, K. J. Med. Chem. 1986, 29, 1038-1046. (c) Truscheit, E.; Frommer, W.; Junge, B.; Müller, L.; Schmidt, D.D.; Wingender, W. Angew. Chem. Int. Ed. Engl. 1981, 20, 744-761.
- (a) Denis, J.W. Cancer Res. 1986, 46, 5131-5136. (b) Humphries, K.J.; Matsumoto, K.; White, S.; Olden, K. Cancer Res. 1986, 46, 5215-5222. (c) Ostrander, G.K.; Scriber, N.K.; Rohrschneider, L.R. Cancer Res. 1988, 48, 1091-1094.
- (a) Gruters, R.A.; Neefjes, J.J.; Tersmette, M.; de Goede, R.E.Y.; Tulp, A.; Huisman, H.G.; Miedema, F.; Ploegh, H.L. *Nature* 1987, 330, 74-77. (b) Walker, B.D.; Kowalski, M.; Goh, W.C.; Kozarsky, K.; Sodroski, J. Proc. Natl. Acad. Sci. USA 1987, 84, 8120-8124.
- 4. Sinnot, M.L. Chem. Rev. 1990, 90, 1171-1202.
- (a) Blériot, Y.; Tellier, C. Regard sur la Biochimie 1997, 1, 41-51. (b) Bols, M. Acc. Chem. Res. 1998, 31, 1-8. (c) Carbohydrate Mimics, Chapleur, Y. Eds., Wiley-VCH, Weinheim, 1998.
- 6. Uchida, C.; Kimura, H.; Ogawa, S. Bioorg. Chem. Lett. 1997, 5, 921-939.
- (a) Knapp, S.; Kirk, B.A.; Vocadlo, D.; Withers, S.G. Synlett 1997, 5, 435-436. (b) Avalos Gonzalez, M.; Fuentes Mota, J.; Gomez Monterrey, I.M.; Jimenez Requejo, J.L.; Palacios Albarran, J.C.; Ortiz Mellet, M.C. Carbohydr. Res. 1986, 154, 49-62.
- 8. Duréault, A.; Greck, C.; Depezay, J.C. Tetrahedron Lett. 1986, 27, 4157-4160.
- 9. Saito, S.; Nakajima, H.; Inaba, M.; Moriwake, T. Tetrahedron Lett. 1989, 30, 837-838.
- 10. Delaunay, D.; Toupet, L.; Le Corre, M. J. Org. Chem. 1995, 60, 6604-6607.
- (a) Levallet, C.; Lerpiniere, J.; Ko, S-Y. Tetrahedron, 1997, 53, 5291-5304. (b) Kim, K-S.; Qian, L. Tetrahedron Lett. 1993, 34, 7677-80. (c) Atwall, K.S.; Ahmed, S.Z.; O'Reilly, B.C. Tetrahedron Lett. 1989, 30, 7313-7316. (d) Poos, M.A.; Iwanowicz, E.; Reide, J.A.; Lin, J.; Gu, Z. Tetrahedron Lett. 1992, 33, 5933-5936. (e) Gauzy, L.; Le Merrer, Y.; Depezay, J.C. Synlett 1998, 402-404.
- 12. Suzuki, K.; Fujii, T.; Sato, K-I.; Hashimoto, H. Tetrahedron Lett. 1996, 37, 5921-24.
- 13. Selected physical data of 1·2 HCl: [α]²⁰_D +23 (c 0.9, CH₃OH); ¹H NMR (250 MHz, D₂O) δ: 3.86-3.99 (m, 4H, CH₂N), 4.09 (m, 2H, CHO), 4.47 (m, 2H, CHS); ¹³C NMR (63 MHz, D₂O) δ: 53.4 (CH₂N), 54.8 (CHS), 73.7 (CHO), 176.0 (C=N).
- Selected physical data of 2·2 HCl: [α]²⁰_D +13 (c 1.2, CH₃OH) ; ¹H NMR (250 MHz, CD₃OD) δ: 3.54-3.60 (m, 2H, CH₂S), 3.63-3.73 (m, 2H, CH₂S), 3.82 (m, 2H, CHN), 4.46-4.57 (m, 2H, CHO) ; ¹³C NMR (63 MHz, CD₃OD) δ: 35.2 (CH₂S), 67.6 (CHN), 74.6 (CHO), 177.3 (C=N).