This article was downloaded by: [University of Chicago Library] On: 12 November 2014, At: 16:37 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

High Throughput Screening Under Zinc-Database and Synthesis a Dialkylphosphinic Acid as a Potential Kari Inhibitor

Xing-Hai Liu $^{\rm a}$, Pei-Quan Chen $^{\rm a}$, Wan-Cheng Guo $^{\rm a}$, Su-Hua Wang $^{\rm a}$ & Zheng-Ming Li $^{\rm a}$

^a The Research Institute of Elemento-Organic Chemistry, The State-Key Laboratory of Elemento-Organic Chemistry, National Pesticide Engineering Research Center, Nankai University, Tianjin, China Published online: 20 Jun 2008.

To cite this article: Xing-Hai Liu , Pei-Quan Chen , Wan-Cheng Guo , Su-Hua Wang & Zheng-Ming Li (2008) High Throughput Screening Under Zinc-Database and Synthesis a Dialkylphosphinic Acid as a Potential Kari Inhibitor, Phosphorus, Sulfur, and Silicon and the Related Elements, 183:2-3, 775-778, DOI: <u>10.1080/10426500701807897</u>

To link to this article: <u>http://dx.doi.org/10.1080/10426500701807897</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the

Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



High Throughput Screening Under Zinc-Database and Synthesis a Dialkylphosphinic Acid as a Potential Kari Inhibitor

Xing-Hai Liu, Pei-Quan Chen, Wan-Cheng Guo, Su-Hua Wang, and Zheng-Ming Li

The Research Institute of Elemento-Organic Chemistry, The State-Key Laboratory of Elemento-Organic Chemistry, National Pesticide Engineering Research Center, Nankai University, Tianjin, China

A novel phosphorus derivative was synthesized through HTVS. The title compounds were confirmed by MS, 1H NMR, ^{31}P NMR. The DOCK was also studied.

Keywords DOCK; KARI; phosphorus derivatives; synthesis

1. INTRODUCTION

Ketol-acid reductoisomerase (KARI; EC 1.1.1.86)¹ is an attractive target for argochemical discovery because it catalyzes the second important step in the biosynthesis of the branched chain amino acids that exist in higher plants only and not in animals. Thus, it is an ideal target from which to design nontoxic KARI-inhibitors as potential novel herbicides. The reaction catalyzed by KARI is shown in Figure 1.

In particular, as high throughput screening considerably increased the numbers of new chemical entities to be studied, the expectations to find new bio-active molecules among these compounds were high at the beginning. However, high throughput screenings (HTS) yielded so far the expected success, and therefore virtual screening approaches emerged and largely evolved. Today, high throughput virtual screening (HTVS) is useful in argochemical innovation.

This work was funded by the National Basic Research Program of China (No. 2003CB114406), National Natural Science Foundation Key Project of China (No. 20432010), and The High Performance Computing Project of Tianjin Ministry of Science and Technology of China (No. 043185111-5).

Address correspondence to Zheng-ming Li, The Research Institute of Elemento-Organic Chemistry, The State-Key Laboratory of Elemento-Organic Chemistry, National Pesticide Engineering Research Center, Nankai University, Tianjin 300071, China. E-mail:nkzml@vip.163.com; xhliu@mail.nankai.edu.cn



FIGURE 1 Reaction catalyzed by KARI.

2. METHODS AND EXPERIMENTS

2.1. HTVS

Based on the reported crystal structure of complexes of the enzyme KARI, 1000 new molecules were predicted with high affinity for KARI from ZINC-database (Drug-Like) searching, using program eHiTS.⁵ The computational flow chart for this study was shown as in Figure 2. Among them, many compounds contain phosphorus structures. A dialkylphosphinic acid was noticed which is analogy of HOE 704 (Figure 1).



FIGURE 2 Computational and experimental flow chart for this study.

2.2. Synthesis

The dialkylphosphinic acid was synthesized according the reference⁶ (Scheme 1). The physical chemistry data were accordance with the reference.



SCHEME 1 The synthesis route of the title compound.

3. RESULTS AND DISCUSSION

3.1. HTVS

A dialkylphosphinic acid was docked with KARI and estimated the pKi = -5.235. The interaction pattern (Figures 3 and 4) between the titled compound and KARI was same as IpOHA (Figure 1). The Mg2+ ions and the crystal water molecules, inhibitor in the active site are necessary for constructing and maintaining the conformation of the active



FIGURE 3 Spinage KARI active site with title compound.



FIGURE 4 Superimposition of the docked modeled into the binding pocket conformation.

site and thus the binding affinity of ligand or the enzymatic activity of title compound and IpOHA in complex with spinage KARI.

3.2. Synthesis and Biological Activity

The synthesis route of the title compound is shown in the Scheme 1. The method was not modified. The dialkylphosphinic acid was synthesized for further bioassay experiments.

4. CONCLUSION

A new potential lead compound of KARI inhibitor was found by virtual and laboratory screening for further optimization.

REFERENCES

- R. Dumas, V. Biou, F. Halgand, R. Douce, and R. G. Duggleby, Acc. Chem. Res., 34, 399 (2001).
- [2] A. Schulz, P. Spönemann, H. Köcher, and F. Wengenmayer FEBS, Lett., 238, 375 (1988).
- [3] A. Aulabaugh and J. V. Schloss Biochemistry, 29, 2824 (1990).
- [4] Y. T. Lee, H. T. Ta, and R.G. Duggleby, Plant Science, 168, 1035 (2005).
- [5] SimBioSys Inc., http://www.simbiosys.ca/ehits/.
- [6] P. Majewski, Phosphorus, Sulfur, Silicon, 45, 151 (1989).