

The Production of Acridines by a Nitrene Insertion Reaction

By G. R. CLIFF and GURNOS JONES*

(*University of Keele, Keele, Staffordshire ST5 5BG*)

Summary Thermal decomposition of 2-azido-2'-methoxydiphenylmethane gives predominantly acridans and acridines, as well as the expected azepinoindoles, while 2-azido-4'-methoxydiphenylmethane and 2-azido-2'-methyldiphenylmethane give azepinoindoles with no trace of acridans or acridines.

NITRENE insertion reactions leading to the production of five- or seven-membered heterocyclic rings have often been reported.¹ The only aromatic azide decomposition reactions proceeding through a nitrene intermediate and leading to six-membered heterocycles are those in which the nitrene

inserts into the C-H bond of a suitably placed methyl or methylene group (giving dihydrophenanthridines² or tetrahydroquinolines³) and those leading to phenothiazines.^{4,5} When an alkyl nitrene substitutes in an aromatic ring to give a dihydrophenanthridine,⁶ the yield is low. We now report a case where nitrene insertion leads predominantly to acridine derivatives, as well as to azepinoindoles.

The product from thermal decomposition[†] of the parent azide (**1**) was the azepinoindole (**2**).⁷ Similarly, 2-azido-4'-methoxydiphenylmethane (**3**) gave the methoxyazepinoindole (**4**).[‡] However, decomposition of 2-azido-2'-methoxydiphenylmethane (**5**) gave a mixture of ten components.


[†] All decompositions were performed in trichlorobenzene under a nitrogen atmosphere, at 180–200°.

[‡] All compounds mentioned were fully characterised and gave satisfactory analyses unless otherwise stated.

The principal products and their percentage yields estimated by g.c. were as follows: the methoxyazepinoindole (6) and the azepinoindolone (7) (7.7%); acridan (8) (10%) and acridine (9) (2.8%); 1-methoxyacridan (10) (26%) and 1-methoxyacridine (11) (7%).§ Acridine derivatives make up approximately 46% of the total products, and over 80% of those separable by g.c.;¶ most of the compounds can be derived from the two azanorcaradiene intermediates shown in the Scheme. Comparison with the reaction producing methoxylated phenothiazines would have suggested the isomeric 4-methoxyacridan as the most likely product.⁵

The methoxyacridan (10) is extremely readily oxidised; rigid exclusion of oxygen during decomposition and work-up suggests that an internal oxidation-reduction system is

responsible for the production of 1-methoxyacridine (11). The new insertion process is not due to an *ortho* (steric) effect, since decomposition of 2-azido-2'-methyldiphenylmethane (14) gave as the main characterised product the azepinoindole (15). Examination of the crude mixture of products by n.m.r. spectroscopy showed no trace of acridan or acridine products.



We thank the S.R.C. for a grant (to G.R.C.).

(Received, November 9th, 1970; Com. 1941.)

§ Identical with a synthetic sample supplied by Professor E. Ritchie.

¶ Also shown to be present by g.l.c.-m.s. examination were 2-amino-2'-methoxydiphenylmethane, and the methoxyazepinoindole (16).

¹ See, for example, R. K. Smalley and H. Suschitzky, *Chem. and Ind.*, 1970, 1338.

² G. Smolinsky, *J. Amer. Chem. Soc.*, 1960, **82**, 4717; *ibid.*, 1961, **83**, 2489.

³ G. Smolinsky and B. I. Feuer, *J. Org. Chem.*, 1964, **29**, 3097.

⁴ P. A. S. Smith, B. B. Brown, R. K. Putney, and R. F. Reinisch, *J. Amer. Chem. Soc.*, 1953, **75**, 6335.

⁵ J. I. G. Cadogan and S. Kulik, *Chem. Comm.*, 1970, 792 and references therein.

⁶ R. A. Abramovitch and E. P. Kyba, *Chem. Comm.*, 1969, 265.

⁷ G. R. Cliff, E. W. Collington, and G. Jones, *J. Chem. Soc. (C)*, 1970, 1490.