

Gold(III)-Catalyzed Direct Acetoxylation of Arenes with Iodobenzene Diacetate


Di Qiu, Zhitong Zheng, Fanyang Mo, Qing Xiao, Yu Tian, Yan Zhang, and Jianbo Wang*

Beijing National Laboratory of Molecular Sciences (BNLMS) and Key Laboratory of Bioorganic Chemistry and Molecular Engineering of Ministry of Education, College of Chemistry, Peking University, Beijing 100871, China

wangjb@pku.edu.cn

Received June 1, 2011

ABSTRACT



AuCl₃-catalyzed direct acetoxylation of electron-rich aromatic compounds has been achieved with iodobenzene diacetate as the acetoxylation reagent.

Direct acetoxylation of arenes with transition metal catalysis has remained a challenging task over the past decades.¹ In 1996, Crabtree and co-workers reported the direct palladium(II)-catalyzed acetoxylation of arenes with iodobenzene diacetate.² The reaction achieved moderately high efficiency; however, the regioselectivity of the reaction is far from satisfactory. More recently, Sanford and co-workers have developed highly regioselective acetoxylation by employing a directing group strategy.^{1,3} Detailed mechanistic investigation indicated that the Pd(IV)/Pd(II) catalytic cycle was involved in the reaction. With a similar Pd(II)-catalyzed system, regioselective acetoxylation of indole derivatives has been reported by Lei,^{4a} Pelcman,^{4b} Zhang,^{4c} and Kwong^{4d} recently. Moreover, Yu and co-

workers have reported a Cu(II)-catalyzed acetoxylation with O₂ as the oxidant.⁵ Although remarkable achievements have been made in this area,⁶ it is still highly desirable to develop an alternative approach toward the goal of direct catalytic acetoxylation of aromatics with high efficiency and generality.

In recent years, gold catalysts have been extensively studied for their unique feature in mediating various transformations in organic chemistry.⁷ In particular, gold-catalyzed direct C–H bond functionalization has attracted great attention as an effective approach toward aromatic C–H bond activation.^{7d} It has been well-established that gold(III) can electrophilically metalate an

(1) For a recent review, see: Lyons, T. W.; Sanford, M. S. *Chem. Rev.* **2010**, *110*, 1147.

(2) Yoneyama, T.; Crabtree, R. H. *J. Mol. Catal. A* **1996**, *108*, 35.

(3) (a) Dick, A. R.; Hull, K. L.; Sanford, M. S. *J. Am. Chem. Soc.* **2004**, *126*, 2300. (b) Desai, L. V.; Hull, K. L.; Sanford, M. S. *J. Am. Chem. Soc.* **2004**, *126*, 9542. (c) Dick, A. R.; Kampf, J. W.; Sanford, M. S. *J. Am. Chem. Soc.* **2005**, *127*, 12790. (d) Dick, A. R.; Kampf, J. W.; Sanford, M. S. *Organometallics* **2005**, *24*, 482. (e) Kalyani, D.; Sanford, M. S. *Org. Lett.* **2005**, *7*, 4149. (f) Racowski, J. M.; Dick, A. R.; Sanford, M. S. *J. Am. Chem. Soc.* **2009**, *131*, 10974. (g) Stowers, K. J.; Sanford, M. S. *Org. Lett.* **2009**, *11*, 4584. (h) Neufeldt, S. R.; Sanford, M. S. *Org. Lett.* **2010**, *12*, 532.

(4) For palladium-catalyzed acetoxylation of indoles, see: (a) Liu, Q.; Li, G.; Yi, H.; Wu, P.; Liu, J.; Lei, A. *Chem.—Eur. J.* **2011**, *17*, 2353. (b) Mutule, I.; Suna, E.; Olofsson, K.; Palcman, B. *J. Org. Chem.* **2009**, *74*, 7195. (c) Liang, Z.; Zhao, J.; Zhang, Y. *J. Org. Chem.* **2010**, *75*, 170. (d) Choy, P. Y.; Lau, C. P.; Kwong, F. Y. *J. Org. Chem.* **2011**, *76*, 80.

(5) Chen, X.; Hao, X.; Goodhue, C. E.; Yu, J. *J. Am. Chem. Soc.* **2006**, *128*, 6790.

(6) For other examples of palladium-catalyzed acetoxylation of arenes, see: (a) Okamoto, M.; Yamaji, T. *Chem. Lett.* **2001**, *30*, 212. (b) Wang, G.; Yuan, T.; Wu, X. *J. Org. Chem.* **2008**, *73*, 4717.


(7) For selected recent reviews on Au-catalyzed transformations, see: (a) Li, Z.; Brouwer, C.; He, C. *Chem. Rev.* **2008**, *108*, 3239. (b) Arcadi, A. *Chem. Rev.* **2008**, *108*, 3266. (c) Jiménez-Núñez, E.; Echavarren, A. M. *Chem. Rev.* **2008**, *108*, 3326. (d) Gorin, D. J.; Sherry, B. B. D.; Toste, F. D. *Chem. Rev.* **2008**, *108*, 3351. (e) Boorman, T. C.; Larrosa, I. *Chem. Soc. Rev.* **2011**, *40*, 1910. (f) Bandini, M. *Chem. Soc. Rev.* **2011**, *40*, 1358. (g) Hopkinson, M. N.; Gee, A. D.; Gouverneur, V. *Chem.—Eur. J.* **2011**, *17*, 8248.

(8) (a) Kharasch, M. S.; Isbell, H. S. *J. Am. Chem. Soc.* **1931**, *53*, 3053. (b) Liddle, K. S.; Perkin, C. *J. Chem. Soc., Chem. Commun.* **1972**, 26. (c) de Graaf, P. W.; Boersma, J.; van der Kerk, G. J. M. *J. Organomet. Chem.* **1976**, *105*, 399. (d) Fuchita, Y.; Utsunomiya, Y.; Yasutake, M. *J. Chem. Soc., Dalton Trans.* **2001**, 2330.

aromatic C–H bond to generate aryl gold(III) species.⁸ Several AuCl₃-catalyzed C–C bond-forming reactions have been suggested to follow the direct auration pathway.⁹ We have recently reported a highly efficient AuCl₃-catalyzed halogenation of aromatics by *N*-halo-succinimide.¹⁰ To account for the extraordinary high catalytic efficiency of this reaction, a dual activation mode has been proposed. Herein we further demonstrate that AuCl₃ is also a highly efficient catalyst in the direct acetoxylation of arenes with iodobenzene diacetate.^{11,12} A dual activation process may also operate with direct auration of aromatic substrates and activation of PhI(OAc)₂ through complexation.

At the outset of the investigation, mesitylene **1a** was employed as the substrate to react with PhI(OAc)₂ in the presence of AuCl₃ (2 mol %) in 1,2-dichloroethane at different temperatures. We observed that the reaction was highly sensitive to temperature (Table 1, entries 1, 2). At 60 °C, only a trace amount of product could be detected, while, at 80 °C, we could isolate the acetoxylation product in moderately high yield (entry 2). However, di- and triacetoxylation products were also observed. The over-acetoxylation problem could be easily circumvented by employing an excess amount of aromatic substrates (entries 4–7). It was also observed that the reaction time could be significantly shortened by carrying it out at 110 °C.

Table 1. AuCl₃-Catalyzed Reaction of Mesitylene **1a** and PhI(OAc)₂^a



entry	ratio	catalyst (mol %)	solvent	temp (°C)	<i>t</i> (h)	yield (%) ^b
1	1:1.2	AuCl ₃ (2)	DCE	60	24	trace
2	1:1.2	AuCl ₃ (2)	DCE	80	24	56 ^c
3	1:1.5	AuCl ₃ (2)	DCE	80	24	68 ^c
4	1.5:1	AuCl ₃ (2)	DCE	110	12	61 ^c
5	2.0:1	AuCl ₃ (2)	DCE	110	12	72
6	2.5:1	AuCl ₃ (2)	DCE	110	12	75
7	3.0:1	AuCl ₃ (2)	DCE	110	12	69
8	2.5:1	AuCl ₃ (1)	DCE	110	24	61
9	2.5:1	AuCl ₃ (5)	DMF	110	24	0
10	2.5:1	AuCl ₃ (5)	CH ₃ CN	110	12	<5
11	2.5:1	AuCl ₃ (5)	AcOH	110	24	53
12	1.5:1	FeCl ₃ (20)	DCE	80	24	0
13	1.5:1	FeBr ₃ (20)	DCE	80	24	0
14	2.5:1	Pd(OAc) ₂ (5)	DCE	110	24	trace
15	2.5:1	AcOH(200)	DCE	110	24	0
16	2.5:1	H ₂ SO ₄ (200)	DCE	110	24	trace
17	2.5:1	– ^d	AcOH	110	24	trace

^a Reaction conditions: mesitylene **1a**, PhI(OAc)₂ (1 mmol), solvent (2 mL). The reaction was monitored by GC-MS. The ratio refers to **1a**/PhI(OAc)₂. ^b Isolated yield based on PhI(OAc)₂. ^c Di- and triacetoxylation products were observed. ^d No catalyst was used. DCE = 1,2-dichloroethane, DMF = *N,N*-dimethylformamide.

Reducing the Au(III) catalyst loading to 1 mol % resulted in a longer reaction time and slightly diminished yield (entry 8). The solvents were found to significantly affect the reaction, and the initially used DCE provided the optimal results (entries 9–11). Finally, for comparison we also carried out the reaction under Fe(III)- and Pd(II)-catalyzed conditions, and under strong acidic conditions as well. None of them afforded the expected acetoxylation products (entries 12–17). The control experiment indicates that no reaction occurs in the absence of catalyst (entry 17).

With the optimal reaction conditions in hand, we proceeded to extend the scope of the reaction, and the results are summarized in Table 2. The reaction is general for electron-rich aromatics. For the substrates bearing three or more methyl groups, the acetoxylation all worked well with 2 mol % AuCl₃ catalyst, even for the substrates bearing halogen substituents or an acetyl group (entries 4–6). When the strong electron-withdrawing group CN was present, a slightly higher catalyst loading was required (entry 7). The aromatic substrates bearing only two methyl substituents require a higher catalyst loading and the reaction took a longer time (entry 8). The diminished yield in this case was due to the formation of homocoupling of *p*-xylene.^{13,14} Interestingly, the acetoxylation also worked with toluene derivatives bearing electron-withdrawing substituents (entries 11, 12). However, an acetoxylation product could not be identified with toluene itself as substrate.

The AuCl₃-catalyzed conditions could also be applied to the aromatic substrates bearing methoxy groups, although a slightly higher catalyst loading was required. It was also observed that the reaction showed excellent regioselectivities. Thus, the reaction with anisole provided the product with an acetoxy group introduced to the *para* position. For all other substituted anisole substrates, the regioselectivity of the acetoxylation is dominated by the methoxy substituent.

To gain insight into the reaction mechanism, kinetic isotope effect (KIE) experiments have been carried out in both an inter- and intramolecular manner. For the intermolecular experiment, an inverse KIE of 0.92 was observed (eq 1), while the corresponding intramolecular experiment afforded an inverse KIE of 0.83 (eq 2). For

(9) (a) Li, Z.; Capretto, D. A.; Rahaman, R. O.; He, C. *J. Am. Chem. Soc.* **2007**, *129*, 12058. (b) Shi, Z.; He, C. *J. Org. Chem.* **2004**, *69*, 3669. (c) Shi, Z.; He, C. *J. Am. Chem. Soc.* **2004**, *126*, 5964. (d) Shi, Z.; He, C. *J. Am. Chem. Soc.* **2004**, *126*, 13596. (e) Reetz, M. T.; Sommer, K. *Eur. J. Org. Chem.* **2003**, 3485. (f) Dyker, G.; Muth, E.; Hashmi, A. S. K.; Ding, L. *Adv. Synth. Catal.* **2003**, *345*, 1247. (g) Hashmi, A. S. K.; Schwarz, L.; Choi, J. H.; Frost, T. M. *Angew. Chem., Int. Ed.* **2000**, *39*, 2285. (h) Luo, Y.; Li, C.-J. *Chem. Commun.* **2004**, 1930. (i) Sun, X.; Sun, W.; Fan, R.; Wu, J. *Adv. Synth. Catal.* **2007**, *349*, 2151.

(10) (a) Mo, F.; Yan, J. M.; Qiu, D.; Li, F.; Zhang, Y.; Wang, J. *Angew. Chem., Int. Ed.* **2010**, *49*, 2028. (b) Qiu, D.; Mo, F.; Zheng, Z.; Zhang, Y.; Wang, J. *Org. Lett.* **2010**, *12*, 5474.

(11) For a recent report on AuCl₃-catalyzed direct amination of arenes, see: Gu, L.; Neo, B. S.; Zhang, Y. *Org. Lett.* **2011**, *13*, 1872.

(12) For a review on hypervalent iodine, see: Zhdkankin, V. V.; Stang, P. J. *Chem. Rev.* **2008**, *108*, 5299.

(13) Kar, A.; Mangu, N.; Kaiser, H. M.; Tse, M. K. *J. Organomet. Chem.* **2009**, *694*, 524.

(14) For detailed discussion on the acetoxylation and homocoupling, see Supporting Information.

Table 2. AuCl₃-Catalyzed Acetoxylation of Arenes^a

entry	substrate 1	AuCl ₃ (mol %)	t(h)	product 2	yield (%) ^b	entry	substrate 1	AuCl ₃ (mol %)	t(h)	product 2	yield (%) ^b
1		2	12		75	12		5	24		66
2		2	8		70	13		5	12		60
3		2	8		73	14		5	12		45
4		2	12		77	15		5	12		66
5		2	8		82	16		5	24		56
6		2	12		60	17		5	24		37
7		3	12		62	18		5	24		36
8		5	24		40	19		5	24		60
9		5	24		76 ^c	20		5	24		53
10		5	24		48	21		5	24		35 ^d
11		5	24		50	22		5	24		49

^a Reaction condition: a solution of arene **1** (2.5 mmol), PhI(OAc)₂ (1 mmol), and AuCl₃ (2 or 5 mol %) in DCE (2 mL) was heated at 110 °C. The reaction was monitored by GC-MS. ^b Isolated yield by silica gel column chromatography based on PhI(OAc)₂. ^c The ratio of **2i** to **2i'** was 2:1, as determined by the integral of ¹H NMR. **2i**: 4-bromo-2,6-dimethylphenyl acetate. **2i'**: 2-bromo-4,6-dimethylphenyl acetate. ^d The ratio of **2u** to **2u'** was 1:4, as determined by the integral of ¹H NMR. **2u**: 2,6-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl acetate. **2u'**: 2,4-dimethyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl acetate.

comparison, the intramolecular KIE for our previously reported AuCl₃-catalyzed bromination with NBS was also measured (eq 3).¹⁰ It gave a similar inverse KIE value of 0.82. The inverse KIE value is consistent with electrophilic aromatic substitution, which shows similar inverse KIE

data in many cases.^{15,16} On the contrary, it has been suggested that electrophilic metalations are characterized by significant primary isotope effects. For example, the electrophilic mercuration of chlorobenzene by Hg(O₂CCF₃)₂ shows a primary isotope effect of 3.9.¹⁷ Recently, a large primary KIE of 5 has been observed for




(15) Tunge, J. A.; Foresee, L. N. *Organometallics* **2005**, *24*, 6440 and references cited therein.

(16) Recently, Zhang and coworkers have observed a normal KIE value of 1.1 in Au(III)-catalyzed aromatic auration; see: Zhang, G.; Luo, Y.; Wang, Y.; Zhang, L. *Angew. Chem., Int. Ed.* **2011**, *50*, 4450.

(17) Lau, W.; Kochi, J. K. *J. Am. Chem. Soc.* **1986**, *108*, 6720.

(18) Lu, P.; Boorman, T. C.; Slawin, A. M. Z.; Larrosa, I. *J. Am. Chem. Soc.* **2010**, *132*, 5580.

direct C–H activation of electron-poor arenes with Au(I) complexes.¹⁸




The formation of the arylgold(III) species has been well-documented in the literature, and it has been demonstrated that the aromatic auration proceeds in an electrophilic manner.⁸ The dual activation mode provides a probable explanation for the extraordinarily high catalytic efficiency of AuCl_3 as compared with other Lewis acid or protonic acid catalysts (Table 1, entries 12–16).^{10,11,19} To accommodate the observed inverse KIE data, we assume that the formation of σ -complex **A** (Wheland intermediate) is the rate-limiting step, in which an inverse secondary KIE is expected (Scheme 1). The subsequent deprotonation to

(19) However, the possibility cannot be ruled out that acetoxylation proceeds through intermediate **B** without activation of $\text{PhI}(\text{OAc})_2$.

(20) Other mechanistic possibilities cannot be ruled out. For example, in the first step of the catalytic cycle, an intermediate iodonium salt of the electron-rich aromatic substrate may be formed, which undergoes a subsequent Au-mediated process leading to the final product. See: (a) Lubriks, D.; Sokolovs, I.; Suna, E. *Org. Lett.* **2011**, *13*, 4324. Such a mechanism is possible because it is known that nonsymmetrical aryl-(phenyl)iodonium acetates can be formed from electron-rich arenes under Lewis acidic conditions. See: (b) Shah, A.; Pike, V. W.; Widdowson, D. A. *J. Chem. Soc., Perkin Trans. 1* **1997**, 2463. We thank the referee who called our attention to such mechanistic possibility.

form mesitylgold(III) species **B** is a fast step. Finally, the mesitylgold(III) species **B** attacks the Au-activated iodobenzene diacetate to afford the acetoxylation product. The rate-limiting step assumption is consistent with the observed similar data of intermolecular KIE. However, further studies are needed to unambiguously establish the reaction mechanism.²⁰

Scheme 1. Plausible Reaction Pathway



In conclusion, we have developed an efficient gold(III)-catalyzed acetoxylation of arenes with iodobenzene diacetate.²¹ This catalytic system is suitable for the electron-rich aromatics, which affords the corresponding acetoxylation products in moderately high yields. Further studies to enhance the catalytic activity and to design a new acetoxylation reagent with higher reactivity are currently underway in our laboratory.

Acknowledgment. The project is supported by National Basic Research Program of China (973 Program, No. 2009CB825300), Natural Science Foundation of China (Grant Nos. 20902005, 20832002, 20772003, 20821062).

Supporting Information Available. Experiment procedure, characterization data, ^1H and ^{13}C NMR spectra, KIE measurement experiments. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(21) For a recent report of direct acetoxylation of anilide promoted by stoichiometric $\text{BF}_3 \cdot \text{OEt}_2$, see: Liu, H.; Wang, X.; Gu, Y. *Org. Biomol. Chem.* **2011**, *9*, 1614.