FACILE REDUCTION OF BENZENETHIOL ESTER UNDER MILD CONDITIONS WITH ZINC BOROHYDRIDE

Hiyoshizo KOTSUKI,* Naka YOSHIMURA, Yasuyuki USHIO, Tatsuya OHTSUKA, and Masamitsu OCHI Department of Chemistry, Faculty of Science, Kochi University, Akebono-cho, Kochi 780

Benzenethiol esters are efficiently reduced to alcohols under mild conditions by treatment with zinc borohydride prepared from zinc chloride and sodium borohydride.

Reductive synthesis of alcohols from esters is one of the most fundamental methods in organic chemistry and several kinds of reducing agents have been developed in the past years.¹⁾ For the reduction of thiol ester group Raney Ni has been commonly used.²⁾ Recently, however, it has been demonstrated that NaBH₄ could serve as a useful reagent for this type of reduction.³⁾

In our separate work we required a method for selective reduction of thiol ester in the presence of other functional groups. The use of NaBH_4 is not enough for our purpose because of its rather basic nature. On the other hand zinc borohydride prepared from ZnCl_2 and NaBH_4 is known as a neutral reducing agent and conventionally employed for the reduction of ketonic carbonyl compounds.⁴⁾ However, the feasibility of extension to the esters was not apparent. Herein we wish to report the facile method for reduction of benzenethiol esters with this reagent.

Benzenethiol esters (1 mmol), which are easily available from acyl chloride and benzenethiol, were dissolved in dry Et_2 O or THF (2-3 mL) and treated with ca. 1 mmol of $\text{Zn}(\text{BH}_4)_2$ (ca. 0.15 M ether solution, 6.7 mL) at room temperature or at 40 °C under N₂ for 2 days. After usual work-up, flash column or preparative layer chromatography on SiO₂ gave the corresponding alcohols in good yields (Eq. 1 and

$$\operatorname{RCOSPh} \xrightarrow{\operatorname{Zn}(\operatorname{BH}_4)_2} \operatorname{RCH}_2OH \qquad (1)$$

Table 1). Under the same conditions methyl benzoate or S-ethyl thiobenzoate could not be reduced. Thus, phthalic acid derivative was reduced only on the benzenethiol ester side (Table 1). We believe that the present method has a synthetic utility for selective reduction of benzenethiol ester group under neutral conditions.

Substrate	Solvent	Temp	Product	Yield/% ^{b)}
O-cosph	Et ₂ 0	RT	О сн ₂ он	81 (84)
C1 COSPh	Et ₂ 0	RT	сц сн ² он	90
MeOOC - COSPh	THF/Et ₂ 0	40 °C	меоос — СР-сн ₂ он	92
	THF/Et ₂ 0	40 °C	С сн ₂ он	81, 84 ^{C)}
COSPh	THF/Et ₂ 0	RT	сн ₂ он	77 ^{d)}
COSPh	Et20	RT	—— сн ₂ он	(89)
COSPh	Et ₂ 0	RT	С-сн2он	99

Table 1. Reduction of benzenethiol ester with zinc borohydride^{a)}

a) Reactions were carried out for 2 days unless otherwise noted. All the products were identified by comparing with authentic samples.

b) Yields represent pure isolated products, values in parentheses are yields determined by GLC.

- c) At room temperature for 4 days.
- d) Reduced yield is due to the volatile product.

References

- W. Carruthers, "Some Modern Methods of Organic Synthesis," 2nd ed, Cambridge University Press, Cambridge (1978), Chap. 7; J. March, "Advanced Organic Chemistry," 3rd ed, Wiley, New York (1985), Chap. 19.
- C. A. Buehler and D. E. Pearson, "Survey of Organic Syntheses," Wiley, New York (1970), p. 209.
- Y. Nagao, K. Kawabata, and E. Fujita, J. Chem. Soc., Chem. Commun., <u>1978</u>, 330;
 H.-J. Liu, R. R. Bukownik, and P. R. Pednekar, Syn. Commun., 11, 599(1981).
- 4) M. Fieser and L. F. Fieser, "Reagents for Organic Synthesis," Wiley, New York, Vol. 11(1984), p. 599; Vol. 10(1982), p. 460; Vol. 5(1975), p. 761; Vol. 3 (1972), p. 337. See also T. Oishi and T. Nakata, Acc. Chem. Res., <u>17</u>, 338 (1984).

(Received April 12, 1986)