

Published on Web 04/29/2006

## Kinetic Resolution of 2-Oxazolidinones via Catalytic, Enantioselective N-Acylation

Vladimir B. Birman,\* Hui Jiang, Ximin Li, Lei Guo, and Eric W. Uffman

Department of Chemistry, Washington University, One Brookings Drive, St. Louis, Missouri 63130

Received March 6, 2006; E-mail: birman@wustl.edu

Kinetic resolution (KR)<sup>1</sup> of racemic alcohols<sup>2</sup> and amines<sup>3</sup> via enzyme-catalyzed enantioselective acyl transfer is well-known. Many nonenzymatic catalysts, which have been developed over the past decade, allow catalytic, enantioselective O-acylation of alcohols.<sup>4</sup> In contrast, there is to date only one report describing the analogous N-acylation of amines.<sup>5</sup> To the best of our knowledge, catalytic, enantioselective N-acylation of another common class of nitrogen nucleophiles—chiral secondary amides—has not been achieved previously using either enzymes or synthetic catalysts. Unlike amines, which can often be resolved via crystallization of their diastereomeric salts with chiral acids, amides are not basic enough to form stable salts. The development of enantioselective acylation of racemic amides would thus provide a convenient means of achieving their resolution.<sup>6</sup> In this communication, we report the first successful examples of this process.

We were encouraged by reports on *achiral* DMAP-catalyzed N-acylation of 4-substituted oxazolidinones with carboxylic acid anhydrides.<sup>7</sup> In addition, various amides, both cyclic and acyclic, are known to be acylated with Boc<sub>2</sub>O in the presence of DMAP.<sup>8</sup> Thus, we decided to employ our recently developed DHIP-based enantioselective acylation catalyst  $1^{9a,c}$  to test the feasibility of the asymmetric version of this reaction (Figure 1). Since our earlier experimental observations in KR of benzylic alcohols indicated the importance of  $\pi - \pi$  and cation $-\pi$  interactions in the chiral recognition of benzylic alcohols, we chose 4-phenyl-2-oxazolidinone **4a** as the starting point of our study (Figure 2).

In the first, proof-of-principle experiment, substrate **4a** was acylated with propionic anhydride in the presence of CF<sub>3</sub>-PIP (1), under conditions analogous to those previously employed for benzylic alcohol substrates. After 24 h, the conversion reached 39%, and the selectivity factor<sup>10</sup> was determined to be 7.8. As anticipated on the basis of our model, the *S*-enantiomer of the substrate (configurationally analogous to *R*-alcohols) was acylated preferentially (Scheme 1).

The second-generation catalyst, Cl-PIQ (2), developed at that time in our laboratory,<sup>9b</sup> proved to be more effective than 1 and thus was selected for further optimization studies. Lower concentrations were chosen to provide a standard set of conditions for all oxazolidinone substrates 4-6, many of which have only limited solubilities. The selectivity was improved when isobutyric anhydride was substituted for propionic, whereas other anhydrides tested were far less successful (Table 1, entries 1-5). *tert*-Amyl alcohol<sup>11</sup> performed better than chloroform as a reaction media, and both were considerably superior to other solvents screened (cf. entries 3 and 6-10).

The influence of the substrate structure was investigated next. As expected, replacement of the phenyl group with 1- or 2-naphthyl led to increased selectivities (entries 11, 12). Comparable results were obtained with the 2-furyl analogue **4d** (entry 13), while the thiophene derivative **4e** was resolved with lower selectivity than





Figure 2. Oxazolidinone substrates.

Scheme 1



4a (entry 14). The importance of  $\pi$ -interactions was confirmed by the absence of any appreciable reaction with the isopropylsubstituted oxazolidinone 4f even at room temperature (entry 15). gem-Dimethyl substituted substrates 5a-c reacted with much higher selectivities than their unsubstituted counterparts 4a-c (entries 16-18 vs 6, 11, 12), whereas in the case of the heteroaryl substrates the differences in selectivity were not significant (entries 19, 20 vs 13, 14). In the case of 5a, increased catalyst loading was needed to achieve respectable reaction rates. Both cis- and trans-4,5diphenyl-2-oxazolidinones (6a and b, entries 21, 22) were resolved with selectivities and rates comparable to those of substrate 4a. It should be noted that the limited solubility of some of the oxazolidinones (4c, 5b, c and 6a) in tert-amyl alcohol prevented reliable selectivity measurements.<sup>12</sup> Substrate 6c was practically insoluble in this solvent, which rendered its resolution inefficient (s < 2).

In a parallel development in our laboratory, catalyst **3** was synthesized and shown to possess markedly higher enantioselectivity than **2** in the KR of benzylic alcohols.<sup>9d</sup> Naturally, these findings prompted us to test the new catalyst in the KR of 2-oxazolidinones. The results far surpassed our expectations. Acylation of substrates **4a–e** and **6a, b** catalyzed by BTM (Table 2, entries 1–7) proceeded

entry	substrate	R′	solvent	time (h)	% conv	S
1	4a	Me	CHCl <sub>3</sub>	3	43	1.3
2	4a	Et	CHCl <sub>3</sub>	3.5	45	10
3	4a	<i>i</i> -Pr	CHCl <sub>3</sub>	24	42	17
4	4a	Ph	CHCl <sub>3</sub>	24	16	-5.5
$5^b$	4a	t-BuO	CHCl <sub>3</sub>	24	<3	ND
6	4a	<i>i</i> -Pr	EtMe <sub>2</sub> COH	19	44	24
7	4a	<i>i</i> -Pr	THF	24	17	12
8	4a	<i>i</i> -Pr	MeCN	24	25	3
9	4a	<i>i</i> -Pr	EtOAc	24	12	8
$10^{b,c}$	4a	<i>i</i> -Pr	PhMe	24	7	9
11	4b	<i>i</i> -Pr	EtMe <sub>2</sub> COH	17	46	38
$12^{c}$	4c	<i>i</i> -Pr	EtMe <sub>2</sub> COH	19	44	44
13	<b>4d</b>	<i>i</i> -Pr	EtMe <sub>2</sub> COH	17	50	25
14	4e	<i>i</i> -Pr	EtMe <sub>2</sub> COH	15	47	16
$15^{b}$	<b>4f</b>	<i>i</i> -Pr	CDCl <sub>3</sub>	16	0	ND
$16^{d}$	5a	<i>i</i> -Pr	EtMe <sub>2</sub> COH	21.5	43	55
$17^{c}$	5b	<i>i</i> -Pr	EtMe <sub>2</sub> COH	10	40	$1.1 \times 10^{2}$
$18^{e}$	5c	<i>i</i> -Pr	EtMe <sub>2</sub> COH	10	49	70
19	5d	<i>i</i> -Pr	EtMe <sub>2</sub> COH	28	50	28
20	5e	<i>i</i> -Pr	EtMe <sub>2</sub> COH	19	49	18
$21^c$	6a	<i>i</i> -Pr	EtMe <sub>2</sub> COH	24	43	26
22	6b	<i>i</i> -Pr	EtMe <sub>2</sub> COH	15	48	19

<sup>a</sup> Unless specified otherwise, the following conditions were used: 0.2 M substrate, 0.75 equiv (R'CO)<sub>2</sub>O, 0.75 equiv *i*-Pr<sub>2</sub>NEt, 4 mol % (R)-2, 0 °C. <sup>b</sup> RT. <sup>c</sup> Incompletely dissolved substrate at 0.2 M. <sup>d</sup> 8 mol % (R)-2 was used. e Incompletely dissolved substrate at 0.1 M.

Table 2. KR of 2-oxazolidinones Using BTM (3)

entry	substrate	ee <sub>SM</sub> <sup>a</sup>	ee <sub>P</sub> <sup>a</sup>	time (h)	%conv <sup>a</sup>	S <sup>a</sup>
$1^b$	4a	54.1	98.3	8.5	36	$2.0 \times 10^2$
$2^b$	4b	71.6	99.1	21	42	$4.5 \times 10^{2}$
$3^b$	<b>4</b> c	87.6	97.8	14	47	$2.6 \times 10^{2}$
$4^b$	<b>4d</b>	70.5	95.7	8.5	42	96
$5^b$	<b>4e</b>	96.0	98.2	6	49	$4.3 \times 10^{2}$
$6^b$	6a	71.7	98.5	6	43	$3.0 \times 10^{2}$
$7^b$	6b	72.3	91.8	6	44	50
$8^b$	6c	96.2	93.2	5.75	51	$1.1 \times 10^{2}$
$9^b$	5a	17.0	98.3	8.5	15	$1.3 \times 10^{2}$
$10^{c}$	<b>4</b> a	89.6	96.4	5	48	$1.7 \times 10^{2}$
$11^{c}$	5a	48.4	99.1	12	33	$3.4 \times 10^{2}$
$12^{c}$	5b	80.4	99.1	7	45	$5.2 \times 10^{2}$
$13^{c}$	5c	56.6	98.2	8.5	37	$2.0 \times 10^{2}$
$14^c$	5d	70.5	95.2	7	43	88
$15^c$	5e	98.7	97.4	6.5	50	$3.9 \times 10^2$

<sup>a</sup> Averages of duplicate runs. <sup>b</sup> General conditions: 0.2 M substrate, 0.75 equiv (i-PrCO)<sub>2</sub>O, 0.75 equiv i-Pr<sub>2</sub>NEt, 4 mol % (R)-3, Na<sub>2</sub>SO<sub>4</sub>, CHCl<sub>3</sub>, rt. General conditions: 0.2 M substrate, 1.5 equiv (i-PrCO)<sub>2</sub>O, 0.75 equiv *i*-Pr<sub>2</sub>NEt, 8 mol % (*R*)-3, Na<sub>2</sub>SO<sub>4</sub>, CHCl<sub>3</sub>, rt.

at room temperature in chloroform with selectivity factors 2.5 to 27(!) times higher than those obtained with  $2^{.13}$ 

We were finally able to resolve substrate 6c with high enantioselectivity, taking advantage of its improved solubility under these conditions (entry 8). This compound has previously served as an intermediate in a concise racemic synthesis<sup>13</sup> of cytoxazone (6d), a naturally occurring oxazolidinone possessing cytokine modulating activity.14 BTM-catalyzed KR of gem-dimethyl substituted substrates, 5a-e, proceeded more slowly than that of their unsubstituted counterparts 4a - e (cf., e.g., entries 1 and 9). In most of these cases, high conversions were obtained on a convenient time scale by doubling the catalyst and the anhydride loadings (entries 10-15). Efficacy of both Cl-PIQ and BTM catalysts on preparative scale was confirmed experimentally.15

In conclusion, we have achieved for the first time kinetic resolution of chiral 2-oxazolidinones via catalytic, highly enantioselective N-acylation. Besides providing a convenient route to these compounds in enantiopure form, these findings pave the way for further exploration of enantioselective N-acylation of other classes of chiral secondary amides. Studies in this direction are currently underway in our laboratory.

Acknowledgment. We thank NIGMS (R01 GM072682) and Washington University for financial support. Mass spectrometry was provided by the Wash. U. Mass Spectrometry Resource, an NIH Research Resource (Grant No. P41RR0954).

Supporting Information Available: Experimental procedures and NMR spectra. These materials are available free of charge via the Internet at http://pubs.acs.org.

## References

- (1) Kagan, H. B.; Fiaud, J. C. Top. Stereochem. 1988, 18, 249
- Enzymatic enantioselective acylation of alcohols: (a) Sih, C. J.; Wu, S.-H. *Top. Stereochem.* **1989**, *19*, 63. (b) Klibanov, A. M. *Acc. Chem. Res.* **1990**, *23*, 114. (c) Schoffers, E.; Golebiowski, A.; Johnson, C. R. Tetrahedron 1996, 52, 3769.
- (3) Enzymatic enantioselective acylation of amines: van Rantwijk, F.; Sheldon, R. A. Tetrahedron 2004, 60, 501.
- (4) For recent reviews, see: (a) Vedejs, E.; Jure, M. Angew. Chem., Int. Ed. 2005, 44, 3974. (b) Dalko, P. I.; Moisan, L. Angew. Chem., Int. Ed. 2004, 43, 5138. (c) Jarvo, E. R.; Miller, S. J. Asymmetric Acylation. In Comprehensive Asymmetric Catalysis, Supplement 1; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer-Verlag: Berlin, Heidelberg, 2004; Chapter 43. (d) France, S.; Guerin, D. J.; Miller, S. J.; Lectka, T. Chem. Rev. 2003, 103, 2985.
- (5) Arai, S.; Bellemin-Laponnaz, S.; Fu, G. C. Angew. Chem., Int. Ed. 2001, 40, 234.
- (6) Resolution of chiral oxazolidinones has been achieved by (a) chromatographic separation of their diastereomeric *N*-acyl derivatives: Ishizuka, T.; Kimura, K.; Ishibuchi, S.; Kunieda, T. *Chem. Lett.* **1992**, 991; and (b) enantioselective deacylation of N-acyl-oxazolidinones via catalytic, asymmetric borane reduction: Hashimoto, N.; Ishizuka, T.; Kunieda, T. Tetrahedron Lett. 1998, 39, 6317.
- 2424. (b) Grehn, L.; Ragnarsson, U. Angew. Chem., Int. Ed. Engl. **1985**, 24, 510. (c) Hansen, M. M.; Harkness, A. R.; Coffey, D. S.; Bordwell, F. G.; Zhao, Y. Tetrahedron Lett. **1995**, 36, 8949.
- (9) (a) Birman, V. B.; Uffman, E. W.; Jiang, H.; Li, X.; Kilbane, C. J. J. Am. Chem. Soc. 2004, 126, 12226. (b) Birman, V. B.; Jiang, H. Org. Lett.
   2005, 7, 3445. (c) Birman, V. B.; Li, X.; Jiang, H.; Uffman, E. W.
   Tetrahedron 2006, 62, 285. (d) Birman, V. B.; Li, X. Org. Lett. 2006, 8, 1351.
- (10) Selectivity factor is defined as s = k(fast-reacting enantiomer)/k(slowreacting enantiomer). Both % conversion and selectivity factor are calculated from ee's of the product and the recovered starting material. Average values of duplicate runs are given for the % ee, % C and s. (11) Ruble, J. C.; Tweddell, J.; Fu, G. C. J. Org. Chem. **1998**, 63, 2794.
- (12) Gradual dissolution of a racemic substrate in the course of KR will lead to a lower apparent selectivity compared with that in a homogeneous reaction.
- (13) Hamersak, Z.; Ljubovic, E.; Mercep, M.; Mesic, M.; Sunjic, V. Synthesis 2001, 1989. In the original synthetic route, (-)-cytoxazone 6d was obtained by KR of its racemate via enzymatic O-acylation with s = 39.
- (14) Isolation, structure elucidation, and biological activity: (a) Kakeya, H.; Morishita, M.; Kobinata, K.; Osono, M.; Ishizuka, M.; Osada, H. J. Antibiot. 1998, 51, 1126. (b) Kakeya, H.; Morishita, M.; Koshino, H.; Morita, T.-i.; Kobayashi, K.; Osada, H. J. Org. Chem. 1999, 64, 1052.
- For a list of leading references to synthesis of 6d, see: (c) Kim, J. D.; Kim, I. S.; Jin, C. H.; Zee, O. P.; Jung, Y. H. Org. Lett. 2005, 7, 4025.
  (15) (a) KR of 0.814 g of (±)-5c with Cl-PIQ produced 40% isolated yield of the N-acylated product with 92.1% ee, 48% recovered SM with 75.6% ee ( $C_{HPLC} = 45\%$ , s = 56) and 79% recovered catalyst. (b) KR of 1.00 g of  $(\pm)$ -4e with BTM gave 51% isolated yield of the product with 96.7% ee, 48% recovered SM with 99.4% ee (C<sub>HPLC</sub> = 51%, s = 3.5 × 10<sup>2</sup>) and 60% recovered catalyst. (c) KR of 0.753 g of  $(\pm)$ -**6c** with BTM gave 48% isolated yield of the product with 91.3% ee, 44% recovered SM with 97.3% ee (C<sub>HPLC</sub> = 52%, s = 94) and 62% recovered catalyst. See experimental details in Supporting Information.

JA061560M