

A Novel Intramolecular Photocyclization of *N*-(2-Bromoalkanoyl) Derivatives of 2-Acylanilines *via* 1,8-Hydrogen Abstraction

by Takehiko Nishio^{*a)}, Hiroyuki Koyama^{a)}, Daigo Sasaki^{a)}, and Masami Sakamoto^{b)}

^{a)} Department of Chemistry, Graduate School of Pure and Applied Sciences, University of Tsukuba, Tsukuba-shi, Ibaraki, 305-8571 Japan

^{b)} Department of Applied Chemistry and Biotechnology, Faculty of Engineering, Chiba University, Yayoi-cho, Inage-ku, Chiba, 263-8522, Japan

The photochemical reactions of different *N*-(2-acylphenyl)-2-bromo-2-methylpropanamides have been investigated. Irradiation of the *N*-unsubstituted anilides **1a**–**1c** gave the corresponding dehydrobromination, cyclization, and bromo-migration products **2**, **3**, and **4**, respectively (Table 1). Irradiation of the *N*-alkyl anilides **1e**–**1g** afforded the corresponding deacylation and cyclization products **5** and **6**, respectively, whereas irradiation of the *N*-alkyl anilides **1i**–**1k**, carrying 2-benzoyl groups on the aromatic rings, afforded the unexpected tricyclic lactams **7** (besides **2**, **5**, and **6**). The formation of the cyclization products **6** could be rationalized in terms of an electrocyclic ring closure of the 6 π -electron-conjugated enamides **2** produced by dehydrobromination of **1**, followed by thermal 1,5-acyl migration (Path B in the Scheme). The formation of the bridged lactams **7** probably follows a mechanism involving the 1,7-diradical **8** generated by ζ -H-abstraction (1,8-H transfer) by an excited acyl O-atom (Path A).

1. Introduction. – Intramolecular H-abstraction reactions by an excited C=O group have been extensively investigated from synthetic and mechanistic points of view [1–3]. Generally, H-atoms in γ -position are abstracted most rapidly through a six-membered cyclic transition state (1,5-H transfer), as, *e.g.*, in *Norrish* type II reactions. This type of H-abstraction is greatly facilitated by favorable stereoelectronic and geometric conditions. However, carbonyl compounds that lack suitably aligned γ -H-atoms by reason of conformation or substitution can still undergo intramolecular reactions, *e.g.*, by abstraction of δ - or ε -H-atoms [2][3]. Abstraction from remote positions is a very attractive issue in the photochemistry of carbonyl groups [3–9], but these reactions are generally disfavored for medium- and large-sized cyclic transition states, both statistically and energetically. Elegant examples of remote H-abstractions induced by photoexcitation of aromatic carbonyl compounds have been reported by *Breslow* [10]. Long-distance *Norrish* type II abstractions *via* electron transfer have been observed in the photochemistry of imides [11], amino ketones [12], and sulfide-containing glyoxylates [13].

In this paper, we report a novel example of a photochemical ζ -H-abstraction (1,8-H transfer) effected by photochemical irradiation of different *N*-(2-acylphenyl)-2-bromo-2-methylpropanamides (anilides), and related compounds.

2. Results and Discussion. – Irradiation of *N*-(2-acetylphenyl)-2-bromo-2-methylpropanamide (**1a**) in MeCN with a high-pressure Hg lamp (with Pyrex filter) under Ar atmosphere at ambient temperature afforded the dehydrobromination product **2a**, the quinoline based cyclization product **3a**, and the bromo-migration product **4a** (Table 1).

Table 1. Photoproducts of Compounds **1a–1l**. Unless noted otherwise, irradiations were carried out for 1 h with a Hg lamp.

Entry	X	R ¹	R ²	Solvent	Isolated yield [%]						
					2	3	4	5	6	7	
1	1a	H	Me	H	MeCN	42	18	8	–	–	–
2 ^{a)}	1a				MeCN	33	39	15	–	–	–
3 ^{b)}	1a				MeCN	19	51	19	–	–	–
4	1b	H	Ph	H	MeCN	34	trace	–	–	–	–
5 ^{b)}	1b				MeCN	71	7	6	–	–	–
6 ^{b)}	1c	Cl	Ph	H	MeCN	48	24	trace	–	–	–
7 ^{b)}	1d	H	EtO	H	MeCN	11	56	–	trace	–	–
8	1e	H	Me	Me	MeCN	trace	–	–	17	65	–
9	1e				Benzene	8	–	–	17	38	–
10	1e				MeOH	trace	–	–	6	24	–
11	1f	H	Me	Et	MeCN	trace	–	–	13	42	–
12	1g	H	Me	Bn	MeCN	trace	–	–	3	9	–
13	1h	H	EtO	Me	MeCN	6	–	–	13	41	–
14	1i	H	Ph	Me	MeCN	7	–	–	9	30	32
15 ^{c)}	1i				MeCN	34	–	–	trace	10	19
16	1i				Benzene	16	–	–	14	28	28
17	1i				MeOH	8	–	–	5	17	30
18	1j	Cl	Ph	Me	MeCN	15	–	–	19	30	35
19	1k	H	Ph	Et	MeCN	14	–	–	18	23	18
20	1l	H	Ph	Bn	MeCN	13	–	–	10	–	–

^{a)} Irradiation time: 3 h. ^{b)} Irradiation time: 5 h. ^{c)} Irradiation time: 0.5 h.

We noticed that, with increasing irradiation time, the yields of both the 3,4-dihydroquinolin-2(1*H*)-one **3a** and the migration product **4a** increased on the expense of the dehydrobromination product **2a** (Entries 1–3 in Table 1). Similar results were obtained when compounds **1b** and **1c** were irradiated under the same conditions. Irradiation of the EtOCO-substituted anilide **1d** basically gave only the elimination product **2d** and the cyclization product **3d**.

The *N*-alkylated anilides **1e–1g**, when irradiated under similar conditions, yielded the deacylation products **5e–5g**¹⁾ and the 3-acetyl-3,4-dihydroquinolin-2(1*H*)-ones **6e–6g** (Entries 8–12). When benzene instead of MeCN was used as solvent, the dehydrobromination product **2e** was obtained in low yield, together with **5e** and **6e**. Irradiation of **1h**, which carries an ester function on the benzene ring, yielded the products **2h**, **5h**, and **6h**. In contrast, irradiation of the *N*-substituted anilides **1i–1k** carrying benzoyl substituents on the aniline ring afforded the unexpected tricyclic lactams **7i–7k** (besides **2i–2k**, **5i–5k**, and **6i–6k**; Entries 14–19). Irradiation of the *N*-

¹⁾ A similar photodeacylation was observed in the photolysis of 2-(*N*-acylamino)benzophenones [14].

benzyl (Bn) analog **11** led to a complex product mixture, from which **21** and **51** were isolated by flash chromatography. The structures of the above photoproducts were assigned on the basis of spectral and analytical evidence. In the case of **7i**, the assignment was further confirmed by X-ray crystal-structure analysis (*Figure*).





Figure. X-Ray crystal structure of compound **7i** (ORTEP view)

The formation of the cyclization products **3** can be rationalized in terms of an electrocyclic ring closure of **2** produced by C–Br bond homolysis, followed by dehydrobromination (*Path B* in the *Scheme*) [15e]. Thereby, for $R^2 = H$ (compounds **a–d**), cyclization occurred in *meta*-position to the acyl group due to strong H-bonding between the aniline NH and the acyl C = O groups. The formation of **6e–6k** can also be rationalized by ring closure of **2** (after change in enamide conformation), followed by a 1,5-acyl migration. Analogous electrocyclic reactions have been observed in the photochemistry of enamides [15][16]. The migration products **4** are probably formed by an elimination/addition sequence of HBr, compounds **2** acting as ‘intermediates’.

The formation of the tricyclic lactams **7** can be rationalized by *Path A* in the *Scheme*. After amide bond isomerization, ζ -H-abstraction from the 2-methylpropanoyl group by the excited carbonyl O-atom of the acyl group on the aniline ring may take place *via*

Scheme



a nine-membered-ring transition state. This would result in formation of the hypothetical 1,7-biradical **8**. Subsequent ring closure would then yield the spirocyclic lactam **9**, which may undergo two kinds of ring opening, either to the amide **10** or to the aziridinone **11**. Finally, intramolecular ring closure affords **7**.

The photochemical behavior of the *N*-substituted anilides **1e–1g** and **1i–1k** reveals that the product distribution strongly depends on the acyl group on the aromatic ring, probably due to conformational and steric effects between the neighboring acyl and 2-bromo-2-methylpropanoyl groups. In the room-temperature $^1\text{H-NMR}$ (CDCl_3) spectrum of **1i**, the signals for the two α -Me groups and that of the NMe group at $\delta(\text{H})$ 1.75 (6 H) and 3.51 (3 H), respectively, were broadened. In CD_3CN , these signals were still broad, but became sharp on raising the temperature to *ca.* 60°. This strongly indicates that the conformation of **1i** is highly restricted at room temperature.

Irradiation of **1i** in Me_3CN at 0° or at 60° led, in both cases, to the same products **2i**, **5i**, **6i**, and **7i**. However, their distribution depended on the temperature (*Table 2*). The yield of the H-abstraction product **7i** increased at lower temperature, the ratio (**2i** + **6i**)/**7i** changing from 1:2 to 1:0.37 upon raising the temperature from 0° to 60°.

Table 2. Temperature-Dependent Distribution of Photoproducts in the Reaction of **1i** in MeCN Solution

Time	<i>T</i>	Isolated yield [%]				Ratio (2 + 6)/ 7
		2	5	6	7	
1 h	25°	7	9	30	32	1:0.86
1 h	0°	10	13	22	49	1:1.53
5 h	0°	trace	12	28	56	1:2.00
1 h	60°	31	22	18	18	1:0.37

Finally, we investigated analogous anilides with 4-acyl substituents on the benzene ring. However, irradiation of compounds **12a–12f** in MeCN gave only the corresponding dehydrobromination and cyclization products **13** and **14**, respectively (*Table 3*). Indolones were not detected [15e].

Table 3. Yield of Photoproducts upon Irradiation of Substrates **12a–12f**. Conditions: irradiation for 5 h in MeCN at 25°.

	R^1	R^2	Isolated yield [%]		
			13	14	
12a	Me	H	19	33	
12b	Ph	H	9	43	
12c	EtO	H	40	5	
12d	Me	Me	21	65	
12e	Ph	Me	7	65	
12f	EtO	Me	25	56	

Conclusions. – Hydrogen-abstraction reactions by excited C=O O-atoms typically require favorable stereoelectronic and geometric conditions [2], long-range H-abstractions being rare [3]. Many of the known cases involve amino ketones, amino imides, or sulfide-containing imides, and proceed *via* electron-transfer reactions [11–13]. Our results concerning the photochemical reactions of 2-acyl-substituted anilides with an α -Br substituent on the alkanoyl part demonstrate that long-range H-abstractions are feasible, given the molecules are present in favorable conformations.

Experimental Part

General. Flash chromatography (FC): *Wakogel C-300* silica gel. M.p.: *Yanaco MP-J3* micro-melting-point apparatus; uncorrected. B.p.: *Shibata GTO-350-RD* glass-tube-oven distillation apparatus. IR Spectra: *Jasco FT/IR-300* spectrophotometer; in cm^{-1} . ^1H - and ^{13}C -NMR Spectra: *Jeol JNM-EX 270* (270/67.5 MHz) or *Varian Gemini-200* (200/50 MHz) spectrometers; in CDCl_3 , with Me_4Si as internal standard; δ in ppm, J in Hz.

General Procedure for the Photochemical Reactions of Anilides 1 or 12. A soln. of the anilide (1 mmol) in MeCN (70 ml), unless noted otherwise, was irradiated in a Pyrex tube with a high-pressure 500-W Hg lamp (*Halos EHP; Eikosha*) under Ar atmosphere for 0.5–5 h at r.t. After removal of the solvent, the residue was subjected to FC (SiO_2 ; toluene/AcOEt 19 : 1 $\rightarrow$ 4 : 1) to afford the photoproducts **2–7**, or **13** and **14** (Tables 1 and 3). The structures of the photoproducts **3a**, **3b**, **3d**, **5e–5l**, **6e–6k**, and **14a–14f** were confirmed by comparison of their spectral data with those given in the literature [16].

N-(2-Acetylphenyl)-2-methylprop-2-enamide (**2a**). B.p. 155°/3 Torr. IR (film): 3225, 1685, 1630. ^1H -NMR: 2.12 (s, 3 H); 2.68 (s, 3 H); 5.54 (br. s, 1 H); 6.03 (s, 1 H); 7.09–7.16 (m, 1 H); 7.54–7.60 (m, 1 H); 7.89–7.94 (m, 1 H); 8.83–8.87 (m, 1 H); 12.21 (br. s, 1 H). ^{13}C -NMR: 19.0; 29.0; 121.1; 122.3; 122.9; 133.2; 135.7; 141.0; 141.6; 167.6; 203.4. Anal. calc. for $\text{C}_{12}\text{H}_{13}\text{NO}_2$: C 70.91, H 6.45, N 6.89; found: C 70.95, H 6.24, N 6.60.

N-(2-Acetylphenyl)-3-bromo-2-methylpropanamide (**4a**). M.p. 56.5–58.0°. IR (KBr): 3204, 1696, 1651. ^1H -NMR: 1.42 (d, $J = 6.9$, 3 H); 2.68 (s, 3 H); 2.87–2.96 (m, 1 H); 2.91 (dd, $J = 7.3$, 13.2, 1 H); 3.71 (dd, $J = 7.3$, 9.9, 1 H); 7.51 (dt, $J = 1.0$, 8.6, 1 H); 7.57 (dt, $J = 1.3$, 7.3, 1 H); 7.92 (dd, $J = 1.7$, 7.9, 1 H); 8.78 (dd, $J = 1.0$, 8.6, 1 H); 11.92 (br. s, 1 H). Anal. calc. for $\text{C}_{12}\text{H}_{13}\text{BrNO}_2$: C 50.72, H 4.97, N 4.93; found: C 50.96, H 4.99, N 4.87.

N-(2-Benzoylphenyl)-2-methylprop-2-enamide (**2b**). Compounds **2b** and **4b** could not be completely separated by FC. Thus, for spectral analysis, **2b** was prepared independently by reaction of 2-amino-benzophenone and methacryloyl chloride in the presence of Et_3N . B.p. 225°/3 Torr. IR (film): 3301, 1681, 1632. ^1H -NMR: 2.13 (s, 3 H); 5.54 (d, $J = 0.7$, 1 H); 6.04 (s, 1 H); 7.06–7.13 (m, 1 H); 7.45–7.71 (m, 7 H); 8.74–8.79 (m, 1 H); 11.47 (br. s, 1 H). ^{13}C -NMR: 18.5; 121.3; 121.3; 122.0; 123.0; 128.2; 129.8; 132.3; 133.8; 134.3; 138.7; 140.3; 140.8; 166.8; 199.9. Anal. calc. for $\text{C}_{17}\text{H}_{15}\text{NO}_2$: C 76.96, H 5.70, N 5.28; found: C 76.61, H 5.85, N 5.12.

N-(2-Benzoylphenyl)-3-bromo-2-methylpropanamide (**4b**). ^1H -NMR: 1.42 (d, $J = 6.9$, 3 H); 2.84–2.95 (m, 1 H); 3.51 (dd, $J = 5.6$, 9.9, 1 H); 3.71 (dd, $J = 7.6$, 9.9, 1 H); 7.06–7.14 (m, 1 H); 7.45–7.72 (m, 7 H); 8.75 (d, $J = 1.3$, 1 H); 11.09 (br. s, 1 H). ^{13}C -NMR (non-aromatic signals only): 17.0; 34.4; 45.4; 176.2; 199.2.

N-(2-Benzoyl-4-chlorophenyl)-2-methylprop-2-enamide (**2c**). The products **2c** and **3c** could not be completely separated by FC. Thus, **2c** was prepared independently by reaction of 2-amino-4-chlorobenzophenone and methacryloyl chloride in the presence of Et_3N . M.p. 92–93°. IR (KBr): 3277, 1693, 1629. ^1H -NMR: 2.11 (s, 3 H); 5.55 (s, 1 H); 6.02 (s, 1 H); 7.15–7.28 (m, 2 H); 7.39–7.72 (m, 5 H); 8.74 (d, $J = 8.9$, 1 H); 11.29 (br. s, 1 H). ^{13}C -NMR (non-aromatic and non-olefinic signals only): 18.8; 167.0; 199.0. Anal. calc. for $\text{C}_{17}\text{H}_{14}\text{ClNO}_2$: C 68.11, H 4.67, N 4.67; found: 68.00, H 4.43, N 4.35.

8-Benzoyl-6-chloro-3,4-dihydro-3-methylquinolin-2(1H)-one (**3c**). ^1H -NMR: 1.40 (d, $J = 6.9$, 3 H); 2.89 (dd, $J = 7.3$, 12.9, 1 H); 3.46–3.53 (m, 1 H); 3.65–3.72 (m, 1 H); 7.15–7.28 (m, 2 H); 7.49–7.72 (m, 4 H); 8.65 (d, $J = 9.5$, 1 H); 10.88 (br. s, 1 H). ^{13}C -NMR (non-aromatic signals only): 17.4; 34.6; 45.7; 172.6; 198.8. Anal. calc. for $\text{C}_{17}\text{H}_{14}\text{ClNO}_2$ (mixture of **2c** and **3c**): C 68.11, H 4.67, N 4.67; found: 68.32, H 4.62, N 4.92.

Ethyl 2-[(2-Methyl-1-oxoprop-2-enyl)amino]benzoate (**2d**). M.p. 43.0–44.5°. IR (KBr): 3254, 1684. ^1H -NMR: 1.42 (t, $J = 7.1$, 3 H); 2.17 (s, 3 H); 4.39 (q, $J = 7.1$, 2 H); 5.53 (s, 1 H); 6.00 (s, 1 H); 7.05–7.12 (m, 1 H); 7.52–7.59 (m, 1 H); 8.04–8.09 (m, 1 H); 8.79–8.87 (m, 1 H); 11.62 (br. s, 1 H). ^{13}C -NMR: 14.1; 18.5; 61.3; 115.3; 120.3; 121.0; 122.4; 130.8; 134.5; 130.7; 141.6; 144.7; 166.7; 168.4. Anal. calc. for $\text{C}_{13}\text{H}_{15}\text{NO}_3$: C 66.93, H 6.48, N 5.84; found: C 66.62, H 6.53, N 5.84.

N-(2-Acetylphenyl)-N-methyl-2-methylprop-2-enamide (**2e**). B.p. 165°/3 Torr. IR (film): 1691, 1649. ^1H -NMR: 1.69 (br. s, 3 H); 2.53 (s, 3 H); 3.31 (br. s, 3 H); 4.90 (br. s, 1 H); 4.99 (br. s, 1 H); 7.22–7.30 (m,

1 H); 7.35–7.41 (*m*, 1 H); 7.49–7.52 (*m*, 1 H); 7.55–7.69 (*m*, 1 H). ^{13}C -NMR: 19.3; 28.4; 37.2; 115.3; 119.3; 127.1; 128.8; 129.5; 132.2; 135.1; 139.7; 170.7; 198.3. Anal. calc. for $\text{C}_{13}\text{H}_{15}\text{NO}_2$: C 71.86, H 6.96, N 6.45; found: C 71.61, H 6.81, N 6.15.

Ethyl 2-[Methyl(2-methyl-1-oxoprop-2-enyl)amino]benzoate (2h). M.p. 86–87°. IR (KBr): 1715, 1620. ^1H -NMR: 1.38 (*t*, *J* = 6.9, 3 H); 1.70 (br. *s*, 3 H); 3.30 (br. *s*, 3 H); 4.30–4.39 (*m*, 2 H); 4.90 (br. *s*, 1 H); 4.96 (br. *s*, 1 H); 7.21–7.27 (*m*, 1 H); 7.34–7.41 (*m*, 1 H); 7.50–7.57 (*m*, 1 H); 7.93–7.97 (*m*, 1 H). ^{13}C -NMR: 14.1; 20.0; 37.6; 61.5; 119.3; 127.6; 128.6; 129.5; 131.8; 133.0; 140.3; 144.5; 165.5; 171.5. Anal. calc. for $\text{C}_{14}\text{H}_{17}\text{NO}_3$: C 67.99, H 6.93, N 5.66; found: C 67.92, H 7.01, N 5.60.

N-(2-Benzoylphenyl)-N-methyl-2-methylprop-2-enamide (2i). M.p. 74–75°. IR (KBr): 1664, 1621. ^1H -NMR: 1.61 (br. *s*, 3 H); 3.23 (br. *s*, 3 H); 4.96 (br. *s*, 1 H); 5.03 (br. *s*, 1 H); 7.27–7.62 (*m*, 7 H); 7.77–7.80 (*m*, 2 H). Anal. calc. for $\text{C}_{18}\text{H}_{17}\text{NO}_2$: C 77.32, H 6.39, N 5.01; found: C 77.59, H 6.31, N 4.91.

1,2,4,5-Tetrahydro-2,4-dimethyl-1-phenyl-1,4-epoxy-2-benzazepin-3(3H)-one (7i). M.p. 122.5–124.0°. IR (KBr): 1713. ^1H -NMR: 1.63 (*s*, 3 H); 2.65 (*s*, 3 H); 2.92 (*d*, *J* = 17.2, 1 H); 3.12 (*d*, *J* = 17.2, 1 H); 7.01 (*dd*, *J* = 1.0, 7.6, 1 H); 7.12–7.19 (*m*, 2 H); 7.25–7.33 (*m*, 1 H); 7.46–7.57 (*m*, 5 H). ^{13}C -NMR: 21.1; 28.0; 35.8; 80.2; 95.8; 125.1; 126.1; 128.6; 128.7; 129.6; 130.3; 133.6; 134.5; 176.8. Anal. calc. for $\text{C}_{18}\text{H}_{17}\text{NO}_2$: C 77.39, H 6.13, N 5.01; found: C 77.40, H 6.27, N 4.95.

N-(2-Benzoyl-4-chlorophenyl)-N-methyl-2-methylprop-2-enamide (2j). B.p. 210°/3 Torr. IR (film): 1667, 1650, 1630. ^1H -NMR: 1.70 (br. *s*, 3 H); 3.23 (br. *s*, 3 H); 4.92 (br. *s*, 1 H); 5.07 (*s*, 1 H); 7.25–7.59 (*m*, 6 H); 7.77–7.81 (*m*, 2 H). Anal. calc. for $\text{C}_{18}\text{H}_{16}\text{ClNO}_2$: C 68.90, H 5.10, N 4.47; found: C 69.00, H 5.34, N 4.35.

8-Chloro-1,2,4,5-tetrahydro-2,4-dimethyl-1-phenyl-1,4-epoxy-2-benzazepin-3(3H)-one (7j). M.p. 153–154°. IR: 1720. ^1H -NMR: 1.62 (*s*, 3 H); 2.68 (*s*, 3 H); 2.88 (*d*, *J* = 17.2, 1 H); 3.06 (*d*, *J* = 17.2, 1 H); 7.01 (*d*, *J* = 2.0, 1 H); 7.18–7.23 (*m*, 1 H); 7.25–7.31 (*m*, 1 H); 7.51 (*s*, 5 H). ^{13}C -NMR: 21.3; 29.4; 35.6; 80.3; 126.4; 128.8; 129.1; 129.3; 130.2; 132.0; 132.4; 134.2; 174.0. Anal. calc. for $\text{C}_{18}\text{H}_{16}\text{ClNO}_2$: C 68.90, H 5.10, N 4.47; found: C 68.76, H 5.20, N 4.39.

N-(2-Benzoylphenyl)-N-ethyl-2-methylprop-2-enamide (2k). M.p. 106–107°. IR: 1660, 1622. ^1H -NMR: 1.42 (*t*, *J* = 7.3, 3 H); 1.72 (br. *s*, 3 H); 3.24 (br. *s*, 1 H); 4.20 (br. *s*, 1 H); 7.14–7.57 (*m*, 7 H); 7.78 (br. *d*, *J* = 7.3, 2 H). Anal. calc. for $\text{C}_{19}\text{H}_{19}\text{NO}_2$: C 77.79, H 6.53, N 4.77; found: C 77.65, H 6.60, N 4.60.

1,2,4,5-Tetrahydro-2-ethyl-4-methyl-1-phenyl-1,4-epoxy-2-benzazepin-3(3H)-one (7k). M.p. 126.5–127.0°. IR (KBr): 1712. ^1H -NMR: 0.98 (*t*, *J* = 7.3, 3 H); 1.61 (*s*, 3 H); 2.92 (*d*, *J* = 17.1, 1 H); 3.10 (*d*, *J* = 17.2, 1 H); 3.18 (*q*, *J* = 7.3, 2 H); 7.01–7.35 (*m*, 4 H); 7.46–7.61 (*m*, 5 H). ^{13}C -NMR: 12.0; 21.0; 35.7; 38.3; 79.8; 95.9; 125.3; 125.6; 128.6; 129.6; 130.1; 133.6; 134.9; 136.4; 177.0. Anal. calc. for $\text{C}_{19}\text{H}_{19}\text{NO}_2$: C 77.79, H 6.53, N 4.77; found: C, 77.61, H 6.52, N 4.97.

N-(2-Benzoylphenyl)-2-methyl-N-phenylprop-2-enamide (2l). M.p. 135–136°. IR (KBr): 1667. ^1H -NMR: 1.73 (*s*, 3 H); 4.13 (br. *s*, 1 H); 5.03 (br. *s*, 2 H); 5.56 (br. *s*, 1 H); 6.95 (*d*, *J* = 7.6, 1 H); 7.21–7.60 (*m*, 12 H); 7.77 (*d*, *J* = 7.6, 1 H). Anal. calc. for $\text{C}_{24}\text{H}_{21}\text{NO}_2$: C 81.10, H 5.96, N 3.13; found: C 80.87, H 5.87, N 4.10.

N-(4-Acetylphenyl)-2-methylprop-2-enamide (13a). M.p. 124–126°. IR (KBr): 3326, 1672, 1627. ^1H -NMR: 2.07 (*s*, 3 H); 2.58 (*s*, 3 H); 5.51 (*s*, 1 H); 5.85 (*s*, 1 H); 6.73 (*d*, *J* = 8.6, 2 H); 7.94 (*d*, *J* = 8.6, 1 H); 8.08 (br. *s*, 1 H). ^{13}C -NMR: 18.1; 36.1; 118.7; 120.1; 129.1; 132.3; 149.0; 141.6; 166.3; 196.3. Anal. calc. for $\text{C}_{12}\text{H}_{13}\text{NO}_2$: C 70.91, H 6.45, N 6.89; found: C 70.71, H 6.55, N 7.12.

N-(4-Benzoylphenyl)-2-methylprop-2-enamide (13b). M.p. 111.5–113°. IR (KBr): 3323, 1684, 1643. ^1H -NMR: 2.06 (*s*, 3 H); 5.49 (*s*, 1 H); 5.83 (*s*, 1 H); 7.11–7.29 (*m*, 2 H); 7.39–7.62 (*m*, 3 H); 7.66–7.84 (*m*, 4 H); 8.09 (br. *s*, 1 H). ^{13}C -NMR: 18.1; 113.0; 118.5; 120.0; 127.7; 129.3; 131.7; 132.4; 137.2; 140.2; 141.3; 166.3; 195.2. Anal. calc. for $\text{C}_{17}\text{H}_{15}\text{NO}_2$: C 76.96, H 5.70, N 5.28; found: C 76.74, H 5.85, N 5.21.

Ethyl 4-[(2-Methyl-1-oxoprop-2-enyl)amino]benzoate (13c). M.p. 116–118°. IR (KBr): 1719, 1671. ^1H -NMR: 1.38 (*t*, *J* = 7.3, 3 H); 2.05 (*s*, 3 H); 4.29–4.39 (*m*, 2 H); 5.48 (*s*, 1 H); 5.81 (*s*, 1 H); 7.67 (*d*, *J* = 7.9, 2 H); 8.01 (*d*, *J* = 7.79, 2 H). ^{13}C -NMR: 13.9; 18.2; 60.4; 113.3; 118.7; 120.0; 127.9; 130.3; 131.1; 140.2; 141.6; 165.7; 166.3. Anal. calc. for $\text{C}_{13}\text{H}_{15}\text{NO}_3$: C 66.93, H 6.48, N 6.01; found: C 67.01, H 6.62, N 5.93.

N-(4-Acetylphenyl)-N-methyl-2-methylprop-2-enamide (13d). B.p. 165°/3 Torr. IR (film): 1681, 1657, 1630. ^1H -NMR: 1.82 (*s*, 3 H); 2.61 (*s*, 3 H); 3.39 (*s*, 3 H); 5.01 (*s*, 1 H); 5.10 (*s*, 1 H); 7.24 (*d*, *J* = 6.8, 3 H); 7.51 (*d*, *J* = 6.8, 2 H). ^{13}C -NMR: 19.5; 26.0; 36.8; 119.6; 125.4; 128.8; 134.4; 139.7; 148.3; 171.2; 196.3. Anal. calc. for $\text{C}_{13}\text{H}_{15}\text{NO}_2$: C 71.86, H 6.96, N 6.45; found: C 72.00, H 7.17, N 6.41.

N-(4-Benzoylphenyl)-N-methyl-2-methylprop-2-enamide (13e). M.p. 85–86°. IR (KBr): 1653. ^1H -NMR: 1.84 (*s*, 3 H); 3.42 (*s*, 3 H); 5.04 (*s*, 1 H); 5.13 (*s*, 1 H); 7.26 (*d*, *J* = 8.4, 2 H); 7.45–7.66 (*m*, 3 H); 7.77–7.85 (*m*, 4 H). ^{13}C -NMR: 19.6; 36.9; 119.7; 125.2; 127.8; 129.3; 130.6; 132.9; 134.9; 136.7; 139.7; 147.8; 171.3; 195.0. Anal. calc. for $\text{C}_{18}\text{H}_{17}\text{NO}_2$: C 77.39, H 6.13, N 5.01; found: C 77.50, H 6.24, N 4.90.

Ethyl 4-[Methyl(2-methyl-1-oxoprop-2-enyl)amino]benzoate (13f). B.p 175°/3 Torr. IR (film): 1714, 1658, 1631. ¹H-NMR: 1.40 (t, *J* = 7.3, 3 H); 1.81 (d, *J* = 1.0, 3 H); 3.38 (s, 3 H); 4.28 (q, *J* = 7.3, 2 H); 4.99 (br. s, 1 H); 5.00 (br. s, 1 H); 7.21 (d, *J* = 8.6, 2 H); 8.03 (d, *J* = 8.6, 2 H). ¹³C-NMR: 14.2; 20.0; 37.3; 61.0; 120.0; 125.7; 128.5; 130.5; 140.2; 148.6; 165.6; 171.7. Anal. calc. for C₁₄H₁₇NO₃: C 67.99, H 6.93, N 5.66; found: C 68.22, H 6.93, N 5.63.

X-Ray Crystal-Structure Analysis. A crystal of **7i** was grown from CH₂Cl₂/hexane. The intensity data were collected on a *Mac Science MXC-18* diffractometer, with graphite-monochromated CuK_α radiation ($\lambda = 1.54178 \text{ \AA}$), in the $\omega - 2\theta$ scan mode ($2\theta < 69.99^\circ$). Out of 3041 total reflections, 2005 reflections with intensities greater than $3\sigma(I)$ were used. No absorption corrections were made. The structure was solved by direct methods with the *maXus* program. Least-squares refinements were performed, including anisotropic thermal parameters for non-H-atoms, and isotropic refinement of H-atoms located in difference *Fourier* synthesis. Crystal data for **7i**: formula, C₁₈H₁₇O₂N; *M_r* 279.339; *V* = 2960(2) Å³; *Z* = 8; *D_x* = 1.254 g cm⁻³; orthorhombic system, space group *Pbca*; *a* = 15.962(4), *b* = 21.813(8), *c* = 8.501(2) Å, $\alpha = 90.00^\circ$, $\beta = 90.00^\circ$, $\gamma = 90.00^\circ$; *V* = 2960(2) Å³; *R* = 0.092, *R_w* = 0.089.

The crystallographic data (excluding structure factors) for **7i** have been deposited with the *Cambridge Crystallographic Data Centre (CCDC)* as supplementary publication number CCDC-264914. Copies of the data can be obtained, free charge, by application to CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK (fax: +44-1223-336033; e-mail: data_request@ccdc.cam.ac.uk), or via the internet (<http://www.ccdc.cam.ac.uk/products/csd/request>).

REFERENCES

- [1] P. J. Wagner, *Acc. Chem. Res.* **1971**, *4*, 168; P. J. Wagner, *Acc. Chem. Res.* **1983**, *16*, 461; P. J. Wagner, *Acc. Chem. Res.* **1989**, *22*, 83; P. Wagner, in 'CRC Handbook of Organic Photochemistry and Photobiology', Eds. W. M. Horspool, P.-S. Song, CRC Press, New York, 1995, p. 449.
- [2] P. J. Wagner, B.-S. Park, in 'Organic Photochemistry', Ed. A. Padwa, Marcel Dekker, New York, 1991, Vol. 11, p. 227; P. Wagner, P. Klan, in 'CRC Handbook of Organic Photochemistry and Photobiology', 2nd edn., Eds. W. M. Horspool, F. Lenci, CRC Press, Boca Raton, 2004, p. 52-1.
- [3] G. L. Descotes, in 'CRC Handbook of Organic Photochemistry and Photobiology', Eds. W. M. Horspool, P.-S. Song, CRC Press, New York, 1995, p. 501; T. Nishio, *Koukagaku* [Photochemistry] **2003**, *34*, 2.
- [4] H. A. J. Carless, S. Mwesigye-Kibense, *J. Chem. Soc., Chem. Commun.* **1987**, 1673.
- [5] G. Adam, A. Preiss, P. D. Hung, *Tetrahedron* **1987**, *43*, 5815.
- [6] F. Cottet, L. Cottier, G. Descotes, *Can. J. Chem.* **1990**, *68*, 1251.
- [7] G. A. Kraus, Y. Wu, *J. Am. Chem. Soc.* **1992**, *114*, 8705.
- [8] G. A. Kraus, W. Zhang, Y. Wu, *Chem. Commun.* **1996**, 1439.
- [9] K. Mizuno, S. Konishi, Y. Yoshimi, A. Sugimoto, *Chem. Commun.* **1998**, 1659.
- [10] R. Breslow, *Acc. Chem. Res.* **1980**, *13*, 170.
- [11] Y. Kanaoka, *Acc. Chem. Res.* **1978**, *11*, 407; P. H. Mazzocchi, in 'Organic Photochemistry', Ed. A. Padwa, Marcel Dekker, New York, 1981, p. 421; J. D. Coyle, in 'Synthetic Organic Photochemistry', Ed. W. M. Horspool, Plenum Press, New York, 1984, p. 259; H. Mauder, A. G. Griesbeck, in 'CRC Handbook of Organic Photochemistry and Photobiology', Eds. W. M. Horspool, P.-S. Song, CRC Press, New York, 1995, p. 513.
- [12] T. Hasegawa, T. Ogawa, K. Miyata, A. Karakizawa, M. Komiyama, K. Nishizawa, M. Yoshioka, *J. Chem. Soc., Perkin Trans. 1* **1990**, 901; T. Hasegawa, Y. Yamazaki, M. Yoshioka, *Trends Photochem. Photobiol.* **1997**, *4*, 27.
- [13] S. Hu, D. C. Neckers, *Tetrahedron* **1997**, *53*, 7165; S. Hu, D. C. Neckers, *J. Org. Chem.* **1997**, *62*, 7827.
- [14] T. Hasegawa, S. Kosaku, J. Ueno, S. Oguchi, *Bull. Chem. Soc. Jpn.* **1982**, *55*, 3659.
- [15] a) Y. Ogata, K. Takagi, I. Isibe, *J. Org. Chem.* **1971**, *36*, 3945; b) G. R. Lenz, *Synthesis* **1978**, 489; c) I. Ninomiya, T. Naito, *Heterocycles* **1981**, *15*, 1433; d) I. Ninomiya, T. Naito, *Yuki Gosei Kagaku Kyokaiishi* **1989**, *47*, 916, and refs. cit. therein; e) T. Nishio, H. Asai, T. Miyazaki, *Helv. Chim. Acta* **2000**, *83*, 1475.
- [16] T. Nishio, M. Tabata, H. Koyama, M. Sakamoto, *Helv. Chim. Acta* **2005**, *88*, 78.

Received March 8, 2005