

Preliminary communication

SYNTHESIS AND REACTIVITY OF TRIMETHYLPHOSPHITE COMPLEXES OF COBALT, RHODIUM AND IRIIDIUM. CRYSTAL AND MOLECULAR STRUCTURE OF $[\text{IrH}(\text{P}(\text{OMe})_3)_4(\text{P}(\text{OMe})_2\text{OSnMe}_2\text{Cl}_2)][\text{SnCl}_3\text{Me}_2]$

PETER B. HITCHCOCK, STANLEY I. KLEIN and JOHN F. NIXON*

School of Chemistry and Molecular Sciences, University of Sussex, Brighton, BN1 9QJ, East Sussex (Great Britain)

(Received September 16th, 1982)

Summary

Syntheses of $[\text{M}(\text{P}(\text{O})(\text{OMe})_2)(\text{P}(\text{OMe})_3)_4]$ ($\text{M} = \text{Co}, \text{Rh}, \text{Ir}$) are reported and variable temperature ^{31}P NMR studies on the iridium complex are described. Hydrogen reacts differently with the Ir and Rh complexes giving $[\text{IrH}_2(\text{P}(\text{O})(\text{OMe})_2)(\text{P}(\text{OMe})_3)_3]$ and $[\text{RhH}(\text{P}(\text{OMe})_3)_4]$, respectively. The crystal and molecular structure of the novel compound $[\text{IrH}(\text{P}(\text{OMe})_3)_4(\text{P}(\text{OMe})_2\text{OSnMe}_2\text{Cl}_2)][\text{SnCl}_3\text{Me}_2]$ is described.

The recent report [1] of the synthesis and chemistry of certain binary zerovalent transition metal-phosphite complexes prompts us to report related studies on Co^{I} , Rh^{I} and Ir^{I} compounds. Treatment of $[\text{Ir}(\eta^5\text{-C}_9\text{H}_7\text{-}(\text{C}_2\text{H}_4)_2)]$ [2] with an excess of $\text{P}(\text{OMe})_3$ affords the iridium(I) complex $[\text{Ir}(\text{P}(\text{O})(\text{OMe})_2)(\text{P}(\text{OMe})_3)_4]$ (I), which was also obtained by treatment of $[\text{IrCl}(\text{P}(\text{OMe})_3)_5]$ with: (a) Na in ether, (b) Na or K/Hg in ether or (c) with warm acetone. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of I at room temperature indicated that the complex is fluxional, exhibiting an A_4M spectrum ($\delta(\text{P}_\text{A}) - 32.1$, $\delta(\text{P}_\text{M}) - 96.6$ ppm, (relative to $\text{P}(\text{OMe})_3$; $J(\text{P}_\text{A}\text{P}_\text{M})$ 101.1 Hz). At -77°C (Fig. 1) the spectrum changes to an A_3BM spin system $\delta(\text{P}_\text{A}) - 28.5$, $\delta(\text{P}_\text{B}) - 31.0$, $\delta(\text{P}_\text{M}) - 93.1$ ppm; $J(\text{P}_\text{A}\text{P}_\text{B})$ 55.8 Hz, $J(\text{P}_\text{A}\text{P}_\text{M})$ 57.8 Hz, $J(\text{P}_\text{B}\text{P}_\text{M})$ 569.1 Hz. Interestingly the appearance of the $\text{P}(\text{O})(\text{OMe})_2$ resonance at intermediate temperatures [2], is identical to that of the hydride resonance in $[\text{IrH}(\text{P}(\text{OCH}_2)_3\text{CPr}^n)_4]$, in which it has been proposed that the hydrogen is on a tetrahedral face and the lowest energy path for interchanging the phosphorus environments involves a tetrahedral-jump rearrangement mechanism [3,4]. We consider such a process unlikely for the bulkier $\text{P}(\text{O})(\text{OMe})_2$ group in I.





Fig. 1. $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of I, (a) at room temperature and (b) at -77°C in toluene.


Complex I adds hydrogen slowly at room temperature to yield the iridium(III) dihydride complex $[\text{IrH}_2\text{P}(\text{O})(\text{OMe})_2(\text{P}(\text{OMe})_3)_3]$ (II), whose *cis*-stereochemistry was unambiguously established by ^{31}P and ^1H NMR studies.



These results led us to reinvestigate the reaction of $[\text{Rh}(\eta^5\text{-C}_5\text{H}_5)\text{-(C}_2\text{H}_4)_2]$ and $\text{P}(\text{OMe})_3$ which was reported [5] to form $[\text{Rh}_2(\text{P}(\text{OMe})_3)_8]$ largely based on ^{31}P NMR data. We now report that the product from this reaction is $[\text{Rh}(\text{P}(\text{O})(\text{OMe})_2(\text{P}(\text{OMe})_3)_4)]$ (III) and that it exhibits a similar temperature dependent ^{31}P NMR spectrum to I.

Both I and III react quickly with MeI to form $\text{M}(\text{P}(\text{OMe})_3)_5^+ \text{I}^-$ but in contrast to the behaviour of I, complex III reacts rapidly with hydrogen at room temperature to afford quantitative yields of $\text{RhH}(\text{P}(\text{OMe})_3)_4$ and $\text{PH}(\text{O})(\text{OMe})_2$ as evidenced by carefully monitoring the reaction by ^{31}P NMR spectroscopy.

The basic nature of the metals in I and III is reflected by their ready reaction with HX ($X = \text{BF}_4$, $p\text{-CH}_3\text{C}_6\text{H}_4\text{SO}_3$) the products being $[\text{IrH}(\text{P}(\text{O})(\text{OMe})_2)(\text{P}(\text{OMe})_3)_4] \text{X}$ (IV) and $[\text{Rh}(\text{P}(\text{OMe})_3)_4 \text{X}]$ plus $\text{PH}(\text{O})(\text{OMe})_2$. The ^{31}P NMR spectrum of the cation in IV established the *cis*-stereochemistry and it bears a close similarity to that of the neutral complex



$[\text{RuMe}(\text{P}(\text{O})(\text{OMe})_2)(\text{P}(\text{OMe})_3)_4]$ (V), formed [6] via a thermally induced methyl migration reaction of the zerovalent complex $[\text{Ru}(\text{P}(\text{OMe})_3)_5]$.

We have confirmed independently the very recent report by Muetterties et al. [7] that $[\text{Co}(\text{P}(\text{O})(\text{OMe})_2)(\text{P}(\text{OMe})_3)_4]$ is formed in the reaction between $[\text{Co}(\text{P}(\text{OMe})_3)_n \text{Cl}]$ ($n = 3, 4, 5$) with sodium or potassium amalgam and the existence of a paramagnetic monomer $[\text{Co}(\text{P}(\text{OMe})_3)_4]$ and dinuclear $[\text{Co}_2(\text{P}(\text{OMe})_3)_8]$ indicates the complexity of this type of system. By careful control of the phosphite concentration we have recently obtained NMR spectroscopic evidence for the complex $[\text{Rh}_2(\text{P}(\text{OMe})_3)_8]$ and mercury contain-




Fig. 2. Molecular Structure of the cation $[\text{IrH}(\text{P}(\text{OMe})_3)_4(\text{P}(\text{OMe})_2\text{OSnCl}_2\text{Me}_2)]^+$. Ir—P(1) 2.270(10); Ir—P(2) 2.382(13), Ir—P(3) 2.322(10); Ir—P(4) 2.331(11); Ir—P(5) 2.280(11); P(3)—O(1) 1.51(2); P(3)—O(2) 1.52(2); O(1)—Sn 2.30(2) Å.

ing products from the reaction between $[M(\text{olefin})_2\text{Cl}]_2$ ($M = \text{Rh, Ir}$), $\text{P}(\text{OMe})_3$ and Na/Hg [8]. The crystal structure of $\text{Co}_2\text{Hg}(\text{P}(\text{OMe})_3)_8$ has recently been described [7].

The reaction of I with Me_2SnCl_2 in dry THF gave a single product whose ^1H and ^{31}P NMR spectra showed features similar to IV and elemental analysis and a single crystal X-ray structure determination confirmed the formation of the hydrido-iridium(III) complex $[\text{IrH}(\text{P}(\text{OMe})_3)_4(\text{P}(\text{OMe})_2\text{OSnCl}_2\text{Me}_2)]-[\text{SnCl}_3\text{Me}_2]$ (VI), in which the oxygen of the coordinated $\text{P}(\text{O})(\text{OMe})_2$ ligand is the donor site towards tin (see Fig. 2). Formation of the hydride $\text{CoH}(\text{P}(\text{OMe})_3)_4$ has been briefly noted [7] in reactions of Na/Hg with CoCl_2 in THF in the presence of excess $\text{P}(\text{OMe})_3$ and from $\text{NaCo}(\text{P}(\text{OMe})_3)_4$ with Hg in THF.

Crystal data: $\text{C}_{18}\text{H}_{55}\text{Cl}_5\text{IrO}_{15}\text{P}_5\text{Sn}_2$, orthorhombic, space group $Pnn2$, a 31.857(4), b 13.230(2), c 10.560(2) Å, $Z = 4$. The structure was determined by routine heavy atom methods and refined by least squares to $R = 0.077$ for 1972 reflections with $I > \sigma(I)$ measured on a CAD4 diffractometer. The cation structure is shown in Fig. 2 with some selected bond length data. The anion $[\text{SnCl}_3\text{Me}_2]^-$ has a trigonal bipyramidal structure with the methyl groups occupying equatorial positions.

Further studies on these systems will be reported separately.

Acknowledgement. We thank the Brazilian Government (CNPq) for a Scholarship (to S.I.K.).

References

- 1 H.W. Choi and E.L. Muetterties, *J. Amer. Chem. Soc.*, **104** (1982) 153 and refs. therein.
- 2 S.I. Klein and J.F. Nixon, unpublished results.
- 3 E.M. Hyde, J.R. Swain, J.G. Verkade and P. Meakin, *J. Chem. Soc. Dalton*, (1976) 1169.
- 4 P. Meakin, E.L. Muetterties and J.P. Jesson, *J. Amer. Chem. Soc.*, **94** (1972) 5271.
- 5 R. Mathieu and J.F. Nixon, *J. Chem. Soc. Chem. Commun.*, (1974) 147.
- 6 R.K. Pomeroy and R.F. Alex, *J. Chem. Soc. Chem. Commun.*, (1980) 1114.
- 7 E.L. Muetterties personal communication and E.L. Muetterties, J.R. Bleeke, Z.Y. Yang and V.W. Day, *J. Amer. Chem. Soc.*, **104** (1982) 2940.
- 8 S.I. Klein and J.F. Nixon, in preparation.