

Tetrahedron Letters 40 (1999) 6077-6078

A GENERAL METHOD FOR THE ALKALINE CLEAVAGE OF ENOLISABLE KETONES

Alenka Žabjek and Andrej Petrič*

Faculty of Chemistry and Chemical Technology, University of Ljubljana, Aškerčeva 5, SI-1000 Ljubljana, Slovenia

Received 27 May 1999; accepted 29 June 1999

Abstract: An efficient method is described for the cleavage of enolisable aryl methyl and aryl ethyl ketones using an excess of KOH in DMF at an elevated temperature. It presents a general hydrolytic method yielding aromatic carboxylic acids, and is complementary to the widely used oxidative methods for ketone cleavage. © 1999 Elsevier Science Ltd. All rights reserved.

Besides the Haller–Bauer reaction, a method widely used for the alkaline cleavage of carbon-carbon bonds of non–enolisable ketones,¹⁻³ there is practically no data on the cleavage of enolisable ketones in the literature. This is because the reaction proceeds with either poor or no conversion of the starting ketones, even with sodium amide.^{1.4} Moreover, with bis- and polychloroacetophenones the aromatic ring is cleaved, giving reasonable yields of halogenated benzenes.⁵ Other enolisable ketones require heating for prolonged periods at high temperatures with either neat KOH, KOH in water or in paraffin oil, to undergo cleavage.⁶ For cyclobutafuranone derivatives, cleavage was preceded by an aldol-type dimerisation to a ketol.⁷

We report a mild and efficient alkaline cleavage of methyl- and ethyl aryl ketones.⁸ After treating the ketones with an excess of KOH in DMF at 65-68 °C, TLC revealed that more than 80% of the starting material has been consumed within 3 to 6 hours. To ensure complete conversion, however, heating was continued for a further 12 hours. In all test cases (Tables 1 and 2) the bond between the alkyl chain and the carbonyl group had been cleaved exclusively, irrespective of the substituents on the aromatic ring. The procedure was tested using a variety of ketones containing phenyl, naphthalene, and phenanthrene rings with a variety of substituents, and with either a methyl or an ethyl group as the alkyl side chain. The corresponding aryl carboxylic acids were isolated in reasonable yields. The resultant acids were isolated and characterized in the form of free acids or as benzyl esters.

We consider the above transformation to be a general method, complementary to methods of oxidative cleavage of ketones, namely the haloform reaction and the Baeyer-Villiger oxidation, and an extension of the Haller-Bauer reaction. A detailed investigation with the aim to find the scope and limitations of the method and to elucidate the mechanism of the above transformation is in progress.



Table 1. The cleavage of aryl-methyl ketones

Table 2. The cleavage of aryl-ethyl ketones

Acknowledgment. The work was supported by the Ministry of Science and Technology of the Republic of Slovenia.

References and Notes

- 1. Hamlin, K. E.; Weston, A. W. Org. React. 1957, 9, 1-36.
- 2. Gilday, J. P.; Paquette, L. A. Org. Prep. Proc. Int. 1990, 22, 167-201.
- 3. Impastato, F. J.; Walborsky, H. M. J. Am. Chem. Soc. 1962, 84, 4838-4843.
- 4. Lock, G.; Bock, E. Chem. Ber. 1937, 70, 916-25.
- 5. Lock, G.; Schreckeneder, R. Chem. Ber. 1939, 72, 511-517.
- 6. Treboganov, A. D.; Zinkevich, E. P.; Kraevskij, A. A.; Preobrazhenskij, N. A. Zh. Org. Khim. 1967, 3, 1418-1419.
- 7. Mittra, A.; Bhowmik, D. R.; Venkateswaran, R. V. J. Org. Chem. 1998, 63, 9555-9556.
- 8. General procedure for the cleavage of ketones to acids: Each ketone (0.5 mmol) was added to a suspension of finely ground KOH (150 mg, 2.7 mmol) in DMF (12 mL) and gradually heated to 65-68 °C. A Drierite-filled trap was used to exclude moisture. After 3 hours an additional amount of KOH (130 mg, 2.3 mmol) was added and heating was continued for another 12 hours. The reaction mixture was cooled, diluted with water (50 mL) and extracted with methylene chloride (3× 20 mL) to remove nonpolar material. The water layer was acidified with HCl to pH 1 and extracted with methylene chloride (3× 20 mL) to yield the acid. The product was purified by radial chromatography, sublimation, or recrystallisation.
- 9. Adcock, W.; Wells, P. R. Australian J. Chem. 1965, 18, 1351-1364.
- 10. Kendall, J.; Carpenter, C. D. J. Am. Chem. Soc. 1914, 36, 2498-2517.
- 11. Liebermann, C.; Zsuffa, M. Chem. Ber. 1911, 44, 852 858.
- 12. Okamoto, Y.; Inukai, T.; Brown, H. C. J. Am. Chem. Soc. 1958, 80, 4969-4972.
- 13. Rupe, H.; Steinback, A. Chem. Ber. 1910, 43, 3465-3471.
- 14. Bray, L. G.; Dippy, J. F.; Hughes, S. R. C. J. Chem. Soc. 1957, 265-267.
- 15. Eddy, N. B. J. Pharmacol. 1935, 55, 354-364, Chem. Abstr. 1936, 30, 3524.
- 16. Benzyl 6-methoxy-2-naphthoate: m.p. 102-3 °C; Anal.Calcd for C19H16O3: C, 78.06; H, 5.52; Found: C, 77.98; H, 5.54. ¹H NMR δ: 3.94 (3H,s); 5.41 (2H,s); 7.15 (1H,d, J= 2.6); 7.19 (1H, dd, J= 2.6 and 8.9); 7.39 (3H, m); 7.49 (2H, m); 7.75 (1H, d, J= 8.7); 7.83 (1H, d, J= 8.9); 8.06 (1H,dd, J= 1.8 and 8.7); 8.56 (1H, d, J= 1.8).
- 17. Shono, T.; Ishige, O.; Uyama, H., Kashimura, S. J. Org. Chem. 1986, 51, 546-549.