

598. Reactions of Ortho-esters of Titanium. Part IV.*
Alcoholysis Reactions of Alkyl Orthotitanates.

By I. D. VARMA and R. C. MEHROTRA.

Reactions of $\text{Ti}(\text{OMe})_4$, $\text{Ti}(\text{OEt})_4$, $\text{Ti}(\text{OPr}^i)_4$, and $\text{Ti}(\text{OBu}^t)_4$ with alcohols (MeOH , EtOH , Pr^iOH , and Bu^tOH) have revealed the following order of reactivity in replacement reactions: $\text{OMe} \gg \text{OEt} > \text{OPr}^i > \text{OBu}^t$. It has therefore been possible to synthesise the following pure mixed ortho-titanates: $\text{MeO} \cdot \text{Ti}(\text{OR})_3$, $(\text{MeO})_2\text{Ti}(\text{OR})_2$ ($\text{OR} = \text{OEt}$, OPr^i , OBu^t), and $(\text{MeO})_3\text{Ti} \cdot \text{OR}$ ($\text{OR} = \text{OEt}$, OPr^i). The mixed ethyl methyl orthotitanates crystallise from benzene and distil unchanged under reduced pressure. Isopropyl methyl and methyl t-butyl titanates are liquids and undergo disproportionation, mainly into monomethyl tri-isopropyl or tri-t-butyl and tetramethyl titanates. Fresh light has been thrown on the nature of tetramethyl titanate.

A FEW mixed ortho-esters of titanium¹ and of zirconium² have been reported. Higher ortho-esters of titanium have been prepared by alcohol-interchange reactions^{3,4} which have now been shown to be reversible. Data presented in Tables 1 and 2 exhibit

TABLE 1. Reaction: $\text{Ti}(\text{OR})_4 + \text{MeOH}$ (molar ratio 1 : 3).

Ester	$\text{Ti}(\text{OBu}^t)_4$	$\text{Ti}(\text{OPr}^i)_4$	$\text{Ti}(\text{OEt})_4$
$\text{MeO} : \text{Ti}$ (in product)	2·3	2·1	2·0

TABLE 2. Reaction: $\text{Ti}(\text{OR})_4 + \text{R}'\text{OH}$ (molar ratio 1 : 4).

OR' : Ti (in product)					OR' : Ti (in product)				
Ester	MeOH	EtOH	Pr ⁱ OH	Bu ^t OH	Ester	MeOH	EtOH	Pr ⁱ OH	Bu ^t OH
$\text{Ti}(\text{OMe})_4$	—	1·5	1·5	—	$\text{Ti}(\text{OPr}^i)_4$...	—	—	—	2·3
$\text{Ti}(\text{OEt})_4$	—	—	—	1·6	$\text{Ti}(\text{OBu}^t)_4$...	2·7	2·4	1·8	—

the following comparative reactivity of the alkoxy-groups in alcoholysis reactions: $\text{MeO} \gg \text{EtO} > \text{Pr}^i\text{O} > \text{Bu}^t\text{O}$.

* Part III, *J. prakt. Chem.*, in the press.


¹ Ghosh, Ghosh Mazumdar, Bose, and Sen Gupta, *J. Indian Chem. Soc.*, 1954, **34**, 683; Nesmeyanov and Nogina, *Izvest. Akad. Nauk S.S.S.R., Otdel. khim. Nauk*, 1954, 41 (*Chem. Abs.*, 1955, **49**, 6084).

² Mehrotra, *J. Indian Chem. Soc.*, 1954, **31**, 905.

³ Bradley, Mehrotra, and Wardlaw, *J.*, 1952, 2027, 4204, 5020.

⁴ Bradley, Mehrotra, Swanwick, and Wardlaw, *J.*, 1953, 2025.

The foregoing and also reactions for the molar ratios 1 : 1 and 1 : 2 are represented in Tables 4—8. By carrying out the reactions in benzene the compounds of types (I) and (II) have been isolated (R = Et, Prⁱ, or Bu^t).



The identity of some of these orthotitanates (particularly the ethyl methyl orthotitanates) has been established by distilling them unchanged (Table 3). In other cases the mixed orthotitanates have been found to disproportionate when distilled under reduced pressure. The isopropyl (or t-butyl) methyl titanates, in general, yield mainly a mono-methyl derivative and tetramethyl titanate.

TABLE 3.

Alkyl ortho-titanate	B. p./mm.	M (obs.)	Mol. complexity	Alkyl ortho-titanate	B. p./mm.	M (obs.)	Mol. complexity
Ti(OEt) ₄	121°/0.8	556	2.4	Ti(OEt)(OMe) ₃ ...	142°/0.3	657	3.5
Ti(OEt) ₃ (OMe) ...	143°/1.5	553	2.6	Ti(OMe) ₄	190°/1.5	—	4.0
Ti(OEt) ₂ (OMe) ₂	130°/0.2	604	3.0		(Sublimn. temp.)		(extrap.)

Tetramethyl titanate is unique among the orthotitanates, in being an infusible white solid, insoluble in most organic solvents. The mixed methyl titanates are all soluble in benzene. Of these the ethyl methyl titanates crystallise as white solids from benzene, and their individual identity was further established by the unchanged analyses of the products after repeated crystallisation. Attempts have been made to explain the insoluble nature of tetramethyl titanate by its highly polymerised nature or by strong inductive effects of the methyl group. The molecular weights of ethyl methyl orthotitanates (Table 3) have been measured ebullioscopically by a method already described,³ and appear to indicate that the main controlling factor in the unique properties of tetramethyl titanate is not its highly polymerised character but the inductive effects of methyl group.

EXPERIMENTAL

Apparatus.—Moisture was excluded in all the experiments.⁵

Materials.—Alcohols⁶ and benzene⁵ were purified as described elsewhere. Ti(OEt)₄ and Ti(OPrⁱ)₄ (Peter Spence products) were purified by distillation.⁵ Ti(Obu^t)₄⁷ and Ti(OMe)₄⁸ were prepared and purified by distillation and sublimation respectively.

Analytical Methods.—Titanium was estimated as TiO₂. Ethoxy- and isopropoxy-contents of esters were determined by oxidation with dichromate.⁹

Reaction between Methanol and Tetraethyl Titanate.—Methanol (0.74 g., 23 mmoles) was added with constant stirring to Ti(OEt)₄ (5.15 g., 23 mmoles); heat was evolved and a white precipitate settled. Benzene (15.0 g.) was distilled into the reaction mixture. After refluxing gently at 90° for 3 hr., the clear solution was allowed to crystallise overnight at room temperature. After the supernatant liquor had been decanted, the white crystals were dried at 30°/1 mm. (1.6 g.) (Found: Ti, 22.3%). The supernatant liquor was evaporated to dryness at 30°/1 mm. A white solid (2.9 g.) was obtained (Found: Ti, 22.55%). The two solids were mixed and recrystallised from benzene (Found: Ti, 22.3. Calc. for C₇H₁₈O₄Ti: Ti, 22.4%). The above compound (2.2 g.) distilled at 143°/1.5 mm. and a wax-like solid (1.8 g.) was obtained (Found: Ti, 22.25%). It was recrystallised from benzene (Found: Ti, 22.3%). Other reactions are condensed in Tables 4—9.

⁵ Varma and Mehrotra, *J. prakt. Chem.*, 1959, **8**, 235.

⁶ Vogel, "A Text-book of Practical Organic Chemistry," Longmans, Green and Co., London, 1956, pp. 169, 170.

⁷ Speer, *J. Org. Chem.*, 1949, **14**, 655.

⁸ Bischoff and Adkins, *J. Amer. Chem. Soc.*, 1924, **46**, 256.

⁹ Mehrotra, *J. Indian Chem. Soc.*, 1953, **30**, 585; 1954, **31**, 904; *J. Amer. Chem. Soc.*, 1954, **76**, 2266.

TABLE 4. Reaction of methanol with tetraethyl or diethyl dimethyl titanate.

Reactants (g.)			Product		Action of heat on product			
Ti(OEt) ₄	MeOH	C ₆ H ₆ (g.)	Yield (g.)	Ti (%) *	Distillate		Found *	
					B. p./mm.	g.	Ti (%)	Solid (g.)
5.15	0.74 (1 mol.)	15.0	4.5 ^a	22.3	143°/1.5	1.8 ^b	22.25	2.2
5.65	1.59 (2 mol.)	5.0	5.1 ^a	23.0	130°/0.2	4.0 ^b	23.8	4.4
7.82	3.29 (3 mol.)	8.2	6.9 ^a	23.8	130°/0.2	5.6 ^b	23.9	6.0
Ti(OMe) ₂ (OMe) ₂								
3.61	2.52 (>4 mol.)	6.0	3.1 ^a	25.5	142°/0.3	2.4 ^b	25.55	2.6
6.63	18.7 (†)	—	5.4 ^c	27.6	Subl. 190°/1.5	—	27.6 ^d	—

* Ti (%) calc. for: Ti(OEt)₃(OMe), 22.4; for Ti(OEt)₂(OMe)₂, 23.95; for Ti(OEt)(OMe)₃, 25.75; for Ti(OMe)₄, 27.85.

Key for Tables 4—9.

^a White powder sol. in C₆H₆.

^b Wax-like solid sol. in C₆H₆.

^c White solid insol. in C₆H₆.

^d Found in sublimate.

^e Reaction mixture became clear on refluxing.

^f Colourless liquid miscible with C₆H₆.

^g Thick colourless liquid sol. in C₆H₆.

^h Cooling occurs when reagents are mixed.

ⁱ Heat evolved when reagents are mixed.

† Large excess.

TABLE 5. Reaction of methanol with tetraisopropyl titanate.

Reactants (g.)			Product		Action of heat on product			
Ti(OPr ⁱ) ₄	MeOH	C ₆ H ₆ (g.)	Yield (%)	Ti (%) *	Distillate		Sublimate	
					B. p./mm.	g.	Ti (%) *	Ti (%) *
6.34	0.71 (1 mol.)	—	5.7 ^f	18.7	5.1	101°/1.5	4.3 ^f	17.8
5.5	1.22 (2 mol.)	—	4.6 ^g	20.3	3.8	120°/1.5	2.7 ^f	18.3
6.19	2.10 (3 mol.)	50	4.9 ^g	21.25	4.17	126°/2	2.83 ^f	18.8
3.56	1.87 (>4 mol.)	10	2.8 ^g	24.3	2.2	145°/2.5	1.1 ^f	18.9
4.50	10.30 (†) ^e	—	2.2 ^e	27.55	—	Subl. 190°/1.5	—	—

* Ti (%) calc. for: Ti(OMe)(OPrⁱ)₃, 18.7; for Ti(OMe)₂(OPrⁱ)₂, 21.0; for Ti(OMe)₃(OPrⁱ)₁, 23.95.

TABLE 6. Reaction of methanol with tetra-*t*-butyl or dimethyl di-*t*-butyl titanate.

Reactants (g.)			Product		Action of heat on product			
Ti(OBu ^t) ₄	MeOH	C ₆ H ₆ (g.)	Yield (g.)	Ti (%) *	Distillate		Sublimate	
					B. p./mm.	g.	Ti (%) *	Ti (%) *
6.25	0.59 (1 mol.)	—	5.6 ^f	16.0	4.9	99°/1.2	4.1 ^f	15.0
5.57	1.05 (2 mol.)	20	4.5 ^f	18.5	4.1	84°/0.7	2.95 ^f	14.7
5.88	1.73 (3 mol.)	15	3.9 ^f	19.6	2.7	105°/1.5	1.5 ^f	16.0
1.63	0.70 (4 mol.)	6.0	1.1 ^g	21.0	—	—	—	—
Ti(OMe) ₂ (OBu ^t) ₂								
2.75	15.0 (†) ^e	—	1.9 ^e	28.3	—	Subl. 190°/1.5	—	27.5

* Ti (%) calc. for: Ti(OMe)(OBu^t)₃, 16.1; Ti(OMe)₂(OBu^t)₂, 18.7; Ti(OMe)₃(OBu^t)₁, 22.4; Ti(OMe)₄, 27.85.

TABLE 7. Reaction of *t*-butanol with tetraethyl or tetraisopropyl titanate [Ti(OR)₄].

Reactants (g.)			Product					Action of heat on product				
Ti(OEt) ₄	Bu ^t OH	C ₆ H ₆ (g.)	Yield (g.)	Ti (%)	OR (%)	OR : Ti	Product taken (g.)	Distillate B. p./mm.	g.	Ti (%)	OR (%)	OR : Ti
5.61	1.87 (1 mol.)	14	6.1 ^f	19.3	60.15	3.3	—	97—140°/9—10	3.6	17.3	38.6	2.4
7.90	5.19 (2 mol.)	19	9.0 ^f	18.5	51.9	3.0	5.9	—	—	—	—	—
5.24	6.80 (4 mol.)	19.5	6.4 ^f	17.1	38.2	2.4	—	—	—	—	—	—
Ti(OPr ⁱ) ₄												
7.30	1.98 (1 mol.)	15	7.6 ^f	16.2	63.4	3.1	4.3	103.5°/8.5	4.0 ^f	16.2	62.5	3.1
7.52	3.93 (2 mol.)	11	8.0	15.7	50.2	2.6	—	—	—	—	—	—
4.60	4.80 (4 mol.)	18	5.9	14.9	32.6	1.7	—	—	—	—	—	—

TABLE 8. *Reaction of alcohols (R·OH) with tetra-*t*-butyl titanate.*

Reactants (g.)		Product					Action of heat on product						
Ti(Obut) ⁴	PrOH	C ₆ H ₆ (g.)	Yield (%)	Ti (%)	OR (%)	OR : Ti	Product taken (g.)	Distillate		Found in distillate			
								B. p./mm.	g.	Ti (%)	OR (%)	OR : Ti	
5.50	1.05 (1 mol.)	10 ^h	5.3 ^f	14.8	18.8	1.0	2.0	103°/3.5	1.8 ^f	14.6	16.1	0.9	
5.34	1.87 (2 mol.)	10	5.1	14.9	26.1	1.4	—	—	—	—	—	—	
5.96	4.18 (4 mol.)	19	5.7 ^e	15.2	34.4	1.8	—	—	—	—	—	—	
2.58	7.30 (†)	—	2.2 ^f	16.3	61.0	3.0	—	—	—	—	—	—	
EtOH													
6.42	3.53 (4 mol.)	20	5.2 ^f	17.6	40.2	2.4	—	—	—	—	—	—	
3.97	8.56 (†)	—	2.9	19.9	63.6	3.4	2.1	120°/0.8	1.8 ^f	19.9	65.7	3.5	

TABLE 9. *Reaction of alcohols (R·OH) with tetramethyl titanate.*

Reactants (g.)		Product		Action of heat on product			
Ti(OMe) ₄	EtOH	C ₆ H ₆ (g.)	Yield (g.)	Ti (%)	Product taken (g.)	Distillate B. p./mm.	Ti (%)
1.33	1.43 (4 mol.)	25	1.55 ^a	24.9	—	—	—
1.70	6.70 (†)	—	1.0 ^a	23.85	0.8	130°/0.2	0.6 ^b
PrOH							
1.32	1.92 (4 mol.)	25	1.7 ^e	22.2	—	—	—
1.65	16.5 (†)	—	2.0 ^f	18.2	1.5	61°/0.3	1.1 ^f

Ti (%) calc. for: Ti(OMe)₂(OEt)₂, 23.95; for Ti(OPr)₄, 16.85.

We thank Messrs. Peter Spence Ltd. for a gift of titanium alkoxides. One of us (I. D. V.) is indebted to the Scientific Research Committee, Uttar Pradesh, Allahabad, for the award of a scholarship during the tenure of which this work was carried out.

UNIVERSITIES OF LUCKNOW AND GORAKHPUR.

[Received, January 11th, 1960.]