


This article was downloaded by: [Northeastern University]

On: 20 November 2014, At: 23:44

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

### Analysis of Chemical Neutralization Products of Phosphonothiolates by Gas Chromatography Mass Spectrometry

A. K. Gupta<sup>a</sup>, M. Palit<sup>a</sup>, D. K. Dubey<sup>a</sup> & S. K. Raza<sup>a</sup>

<sup>a</sup> Defence Research & Development Establishment, Gwalior, India

Published online: 27 Oct 2010.

To cite this article: A. K. Gupta, M. Palit, D. K. Dubey & S. K. Raza (2003) Analysis of Chemical Neutralization Products of Phosphonothiolates by Gas Chromatography Mass Spectrometry, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 178:8, 1631-1637, DOI: [10.1080/10426500307845](https://doi.org/10.1080/10426500307845)

To link to this article: <http://dx.doi.org/10.1080/10426500307845>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

## ANALYSIS OF CHEMICAL NEUTRALIZATION PRODUCTS OF PHOSPHONOTHIOLATES BY GAS CHROMATOGRAPHY MASS SPECTROMETRY

A. K. Gupta, M. Palit, D. K. Dubey, and S. K. Raza  
Defence Research & Development Establishment, Gwalior, India

(Received November 27, 2001; accepted February 24, 2002)

*A series of phosphonothiolates, including the highly toxic O-Ethyl-S-(2-diisopropylamino) ethyl methylphosphonothioate (VX), have been subjected to chemical neutralization reaction with metallic sodium. The phosphonothiolates decompose to their respective phosphonic and phosphonothioic acids and this results in the detoxification of VX. GC/MS technique in both EI and CI mode has been applied for reaction monitoring and final identification of the neutralization products formed in this reaction.*

**Keywords:** Chemical ionization; chemical neutralization; disulphide; electron ionization; GC/MS; phosphonic acid; phosphonothiolates; sodium; thiophosphonic acid; VX

Chemical warfare agents either can be destroyed by incineration or neutralized by chemical reaction to produce nonlethal or nontoxic products, which can be disposed of safely, or postprocessed.<sup>1</sup> O-Alkyl methylphosphonothiolates constitute one of the most lethal classes of “nerve agents” represented by O-ethyl S(2-diisopropylamino)ethyl methylphosphonothiolate commonly known as VX (**1**) which is a highly toxic chemical and can stop respiratory and nervous functions and can kill in minutes.<sup>2</sup>

VX can be neutralized effectively by the use of bleach at low pH, where VX readily dissolves in acidic solutions; on the other hand the solubility of VX is reduced significantly at higher pH. Hydrolysis catalyzed

The authors thank Dr. D. K. Jaiswal, Director, DRDE and Dr. M. P. Kaushik, Head PTD Division for providing necessary facilities and encouragement.



Address correspondence to S. K. Raza, Defence Research & Development Establishment, Gwalior-474 002, India. E-mail: skraza@hotmail.com

by base is one of the most widely used neutralization reaction for nerve agents. VX ( $pK_a = 9$ ) dissolves in pure water to form a basic solution and the solubility of VX decreases significantly as the solution becomes more basic, leading to very slow rate of hydrolysis.<sup>3</sup> Another approach for the neutralization of VX is oxidation of the sulfur atom. But on oxidation, nitrogen present in VX is predominantly oxidized to the stable N-oxide, which is still toxic.<sup>4</sup> Thus, the neutralization of VX by oxidation is not the method of choice. Nano-size MgO, CaO, and Al<sub>2</sub>O<sub>3</sub> also have been used for the neutralization of VX.<sup>5-7</sup> These reactions are analogous to their solution behavior,<sup>3</sup> with two exceptions: The first is that the corresponding nontoxic phosphonate products reside as surface bound complexes; the second, and most notable exception, is that toxic EA-2192 which forms under basic hydrolysis<sup>8,9</sup> is not observed in these reactions. VX also has been neutralized and decontaminated by using metal ions supported on zeolites. Silver and sodium exchanged zeolites have been used to enhance the rate of decomposition of VX and related compounds at ambient temperature.<sup>10,11</sup> Recently, desulfurization of alkyl and alkyl-aryl organ sulfur compounds with sodium was reported resulting in the formation of corresponding alkanes and small amounts of alkylmercaptans.<sup>12</sup> Our interest in detoxification of highly-toxic VX and related phosphonothiolates prompted us to apply this method for the chemical neutralization of these compounds. In this study we selected VX (**1**), and some structurally related phosphonothiolates as shown in Figure 1.

## RESULTS AND DISCUSSION

The total chemical neutralization was achieved within 120 min. The chemical neutralization of VX (**1**) yielded nontoxic products namely N,N-diisoprylaminoethyl disulfide (**1a**), methylphosphonic acid (**1b**), O-ethyl methylphosphonic acid (**1c**), methylthiophosphonic acid (**1d**), and O-ethyl methylthiophosphonic acid (**1e**). All the phosphonic acids were identified as their trimethylsilyl derivatives both in EI and CI modes. **1a** was observed only while monitoring the neutralization reaction with GC/MS-EI, but after doing the work-up of the reaction mixture, **1a** could not be detected either in organic or aqueous layer. This may have been due to further desulfurization.

The neutralization of S,S-dialkyl alkylphosphonothiolates (**2-8**) produced their respective disulfides (**2a-8a**), alkylphosphonic acids (**2b-8b**), and alkylthiophosphonic acid (**2c-8c**). The acids were identified on the basis of EI and CI data of their silylated derivatives. In **5** we detected S-butyl isopropylthiophosphonic acid (**5d**) as its trimethylsilyl




	<u>X</u>	<u>R<sub>1</sub></u>	<u>R<sub>2</sub></u>
(2)	O	CH <sub>3</sub>	<i>n</i> -C <sub>4</sub> H <sub>9</sub>
(3)	O	C <sub>2</sub> H <sub>5</sub>	<i>n</i> -C <sub>4</sub> H <sub>9</sub>
(4)	O	<i>n</i> -C <sub>3</sub> H <sub>7</sub>	<i>n</i> -C <sub>4</sub> H <sub>9</sub>
(5)	O	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	<i>n</i> -C <sub>4</sub> H <sub>9</sub>
(6)	O	C <sub>2</sub> H <sub>5</sub>	<i>n</i> -C <sub>3</sub> H <sub>7</sub>
(7)	O	<i>n</i> -C <sub>3</sub> H <sub>7</sub>	<i>n</i> -C <sub>3</sub> H <sub>7</sub>
(8)	O	<i>i</i> -C <sub>3</sub> H <sub>7</sub>	<i>n</i> -C <sub>3</sub> H <sub>7</sub>
(9)	S	CH <sub>3</sub>	<i>n</i> -C <sub>3</sub> H <sub>7</sub>

**FIGURE 1** Structures of phosphonothiolates subjected to chemical neutralization reaction.

derivative which was formed most probably due to incomplete desulphurisation.

The neutralization reaction of S,S-dipropyl methylthiophosphonothiolate (**9**) yields not only its corresponding disulfide (**9a**) but also tri- and tetra-sulfides (**9b** and **9c**) respectively. The other neutralization products obtained from **9** were methylthiophosphonic acid (**9d**) and methyl phosphonic acid (**9e**) identified as their trimethylsilyl derivative by GC/MS.

In all these neutralization reactions, we observed disulfides only while monitoring the reaction products before work-up; however, after giving extended reaction time or after the addition of methanol, these disulfides disappear and only corresponding alkylphosphonic acids and alkylthiophosphonic acids were detected by GC/MS as their



**FIGURE 2** Products obtained on chemical neutralization reaction of **1-9** as confirmed by GC/MS.

**TABLE I** EI and CI MS Data of the Compounds Detected from Chemical Neutralization of **1–9**

Compd. no.	Molecular weight	EI mass spectrum (m/z)	CI MS (M + H) <sup>+</sup>
<b>1a</b>	320	42, 72, 89, 114 (100%), 160, 174, 189, 285	321
<b>1b, 2b, 9d</b>	240	59, 73, 105, 133, 147, 181, 209, 225 (100%), 240 (M <sup>+</sup> )	241
<b>1c</b>	196	61, 75, 121, 137, 153 (100%), 169, 181, 196 (M <sup>+</sup> )	197
<b>1d, 2c, 9e</b>	256	73, 93, 113, 133, 147, 241 (100%), 256 (M <sup>+</sup> )	257
<b>1e</b>	212	56, 73, 121, 137, 153, 169 (100%), 197, 212 (M <sup>+</sup> )	213
<b>2a, 3a, 4a, 5a</b>	178	57, 87, 122, 178 (100%), (M <sup>+</sup> )	179
<b>6a, 7a, 8a, 9a</b>	150	66, 78, 108, 150 (100%) (M <sup>+</sup> )	151
<b>3b, 6b</b>	254	73, 121, 135, 147, 195, 211, 226, 239 (100%), 254 (M <sup>+</sup> )	255
<b>4b, 7b</b>	268	73, 135, 195, 211, 225, 237, 253 (100%), 268 (M <sup>+</sup> )	269
<b>5b, 8b</b>	268	73, 135, 147, 195, 211, 226, 253 (100%), 268 (M <sup>+</sup> )	269
<b>5d</b>	284	73, 121, 133, 147, 209 (100%), 242, 269, 284 (M <sup>+</sup> )	285
<b>9b</b>	182	55, 64, 100, 111, 140, 182 (100%) (M <sup>+</sup> )	183
<b>9c</b>	214	65, 70, 104, 130, 142, 193, 214 (100%) (M <sup>+</sup> )	215

trimethylsilyl derivatives. The disulphides perhaps undergo further desulfurization to give alkenes, which were not detected. The EI and CI mass spectral data of various phosphonothiolates obtained from the neutralization reaction are presented in Table I.

Chemical neutralization with sodium metal may thus be a useful reaction for the complete decontamination of thiophosphonates including the highly toxic compounds such as VX. The mass spectral data in both EI and CI for various degradation products can be used for the detection of the compounds even in the reaction mixture.

## EXPERIMENTAL

### Synthesis of Phosphonothiolates

O-Ethyl-S- [2-(diisopropylamino) ethyl] methylphosphonothiolate (VX) was synthesized inhouse according to literature procedure<sup>13</sup> and the final structure and purity was checked by GC/MS and <sup>1</sup>H-NMR. (*Caution*: VX is synthesized in a special laboratory, built mainly for synthesis of highly-toxic compounds. Full body protection including breathing protection is worn when handling the agent, which includes both synthesis work and the neutralization reaction.)

Dialkyl alkylphosphonothiolates (**2–8**) and dialkyl alkylthiophosphonothiolate (**9**) were synthesized according to the literature

procedure<sup>14–16</sup> and the final structure and purity were checked by GC/MS and <sup>1</sup>H-NMR.

### Desulphurization Reaction

Sodium (0.1 mmol) and phosphonothiolates (0.01 mmol) were taken in a solvent (10 mL) [decane for **1**, **2**, **5** and dodecane for **3**, **4**, **6–9**] in a 50 mL two-necked, round-bottom flask fitted with a condenser. The reaction mixture, monitored by GC/MS in EI mode, was vigorously stirred and refluxed till the substrate was totally consumed. After cooling the reaction mixture to room temperature, unreacted sodium was destroyed with methanol at 0°C. Saturated aqueous NH<sub>4</sub>Cl (50 mL) was added to the mixture.

### Extraction and Derivatization

After the addition of saturated aqueous NH<sub>4</sub>Cl solution, the aqueous and organic layers were separated using separating funnel. The organic layer was analyzed by GC/MS in both EI and CI modes as such while the aqueous layer was extracted with diethyl ether and analyzed after concentration. This layer was later silylated with BSTFA at 80°C for 120 min and subjected to GC/MS analysis.

One portion of the aqueous layer was evaporated to dryness on rotavapour followed by reconstitution with acidic methanol (HCl/methanol –0.5N). It was further evaporated to dryness and silylated for GC/MS analysis. Another portion of the aqueous layer after evaporation to dryness was reconstituted with alkaline methanol (1% triethylamine/methanol). It was further evaporated to dryness and silylated for GC/MS analysis.

### GC/MS Analysis

The GC/MS analyses were performed by Varian 3400 GC coupled to a TSQ 7000 mass spectrometer (Finnigan Mat). GC operating conditions were as follows: injector temperature 250°C, transfer line temperature 280°C, column temperature programming 50°C (2 min)—@10°C/min–280°C (5 min), carrier gas helium at pressure of 10 psi. EI mass spectrometric operating conditions were as follows: ion source pressure  $1.5 \times 10^{-6}$  torr, source temperature 150°C, electron energy 70 eV, and emission current 400  $\mu$ A. CI mass spectrometric operating conditions were as follows: ion source pressure with methane as the reagent gas  $1.5 \times 10^{-3}$  torr, source temperature 150°C, electron energy 100 eV, and emission current 300  $\mu$ A.

## REFERENCES

- [1] Yu-Chu Yang, *Chemistry and Industry*, 334 (1995).
- [2] J. A. F. Compton, in *Military Chemical and Biological Agents, Chemical and Toxicological Properties* (The Telford Press, Caldwell, NJ, 1987).
- [3] Y.-C. Yang, J. A. Baker, and J. R. Ward, *Chem. Rev.*, **92**, 1729 (1992).
- [4] Y.-C. Yang, L. L. Szafraniec, W. T. Beaudry, and D. K. Rohrbaugh, *J. Am. Chem. Soc.*, **112**, 6621 (1990).
- [5] G. W. Wagner, P. W. Bartram, O. Koper, and K. J. Klabunde, *J. Phys. Chem. B*, **103**, 3225 (1999).
- [6] G. W. Wagner, O. Koper, E. Lucas, S. Decker, and K. J. Klabunde, *J. Phys. Chem. B*, **104**, 5118 (2000).
- [7] G. W. Wagner, L. R. Procell, R. J. O'Connor, S. Munavalli, C. L. Carnes, P. N. Kapoor, and K. J. Klabunde, *J. Am. Chem. Soc.*, **123**, 1636 (2001).
- [8] Y.-C. Yang, *Acc. Chem. Rev.*, **32**, 109 (1999).
- [9] Y.-C. Yang, L. L. Szafranine, and W. T. Beaudry, *J. Org. Chem.*, **58**, 6964 (1993).
- [10] A. Bellamy, *J. Chem. Soc., Perkin Trans. 2*, 2325 (1994).
- [11] G. W. Wagner and P. W. Bartram, *Langmuir*, **15**, 8113 (1999).
- [12] Zhengkun Yu and J. G. Verkade, *Tet. Lett.*, **39**, 2671 (1998).
- [13] F. Siegfried, *Lehrbuch der Militarchemie*, Band 1-2, 2 ueberarb. Aufl.; militarverlag der DR: Berlin, 1977.
- [14] E. Schegk, H. Schlor, and G. Schrader, *DAS 1*, **058**, 992 (1957).
- [15] G. Schrader, *DAS 1*, **032**, 247 (1954).
- [16] K. N. Anisimov, N. E. Kolobova, and A. N. Nesmejanov, *Izv. Akad-Nauk SSSR, Ser. Khim.*, 669 (1955).