Science Foundation (Grant CHE 82-17950). The NMR facilities used in this research were supported in part through the University of Chicago Cancer Center Grant No. NIH-CA-14599.

Registry No. 1, 113507-51-0; 2, 89317-73-7; 3, 113507-52-1; 4, 113507-53-2; 5, 113507-54-3; 6, 107987-07-5; 7, 62442-83-5; 8, 75511-41-0; 9, 113507-55-4; 10, 84782-16-1; 11, 597-04-6; 12, 4749-12-6; 13,

55424-74-3; 20, 112818-18-5; 21, 32811-25-9; 22, 35293-39-1; 23, 90646-75-6; 24, 113507-56-5; coenzyme B₁₂, 13870-90-1.

Supplementary Material Available: Table VIII containing representative material balances for the reaction of various bromides with $(n-Bu)_3$ SnH (1 page). Ordering information is given on any current masthead page.

Thermal Decomposition and Cobalt–Carbon Bond Dissociation Energies of Organocobalamins: Neopentyl-, (Cyclopentylmethyl)-, (Cyclohexylmethyl)-, (Tetrahydrofurfuryl)- and ((Tetrahydro-2*H*-pyryl)methyl)cobalamin

Sook-Hui Kim, Hui Lan Chen, Natalie Feilchenfeld, and Jack Halpern*

Contribution from the Department of Chemistry, The University of Chicago, Chicago, Illinois 60637. Received September 18, 1987

Abstract: The title compounds were prepared and characterized and their thermal decomposition reactions were studied in aqueous solutions of varying pH and containing varying concentrations of cob(II)alamin (B12,) and of bis(dimethylglyoximato)cobalt(II), $[Co(DH)_2]$. Homolytic cobalt-carbon bond dissociation pathways were identified through competitive trapping of the resulting free radicals according to the following scheme: $R-B_{12} \rightleftharpoons R^* + B_{12}$; $R^* + Co(DH)_2 \rightarrow R-Co(DH)_2$. Determination of the kinetics of these reactions yielded (after correction for the base on-base off equilibria) the following values of the activation parameters: R = neopentyl, $\Delta H^* = 26.7 \pm 1.2$ kcal/mol, $\Delta S^* = 15 \pm 5$ cal/(mol K); R = cyclopentylmethyl, $\Delta H^* = 26.8 \pm 2 \text{ kcal/mol}; \Delta S^* = 3 \pm 6 \text{ cal/(mol K)}$. From these measurements the Co-C bond dissociation energies of the base-on forms of neopentyl-B₁₂ and cyclopentylmethyl-B₁₂ were deduced to be ca. 24 kcal/mol. In ethylene glycol ΔH^* for homolytic dissociation of the Co-C bond of neopentyl-B₁₂ was found to be about 4 kcal/mol higher than in aqueous solution.

It is now widely accepted that coenzyme B_{12} dependent rearrangements are initiated by enzyme-induced homolytic dissociation of the coenzyme cobalt-carbon bond to generate a 5'-deoxyadenosyl radical.1-5

Accordingly, a knowledge of the Co-C bond dissociation energy of this and related organocobalt compounds, and of the factors that influence such bond dissociation energies and that may contribute to enzyme-induced bond weakening and dissociation, is important to an understanding of the coenzyme's role. In this context we have previously determined the Co-C bond dissociation energies of a variety of organocobalt complexes (coenzyme B₁₂ models!) in which the nature of the equatorial ligand (porphyrin, Schiff base, etc.), the cobalt-bonded alkyl group, and the transaxial ligand (amine, phosphine, etc.) were varied.⁶⁻⁹ Such prior studies have included determination of the Co-C bond dissociation energy of coenzyme B_{12} itself ((5'-deoxyadenosyl)cobalamin), for which values ranging from 26 to 30 kcal/mol have been reported.^{10,11} Attention also is directed to several other earlier studies on the thermal decomposition of cobalamins.¹²

In this paper we report studies on the thermal decomposition of several other cobalamins $(R-B_{12})$ and, in some cases, determination of their Co-C bond dissociation energies. A particular objective was to attempt to identify distinctive features that might be associated with the 5'-deoxyadenosyl moiety as compared with other organic substituents. These studies also bear on the validity of one of the procedures that has previously been employed to determine the Co-C bond dissociation energy of coenzyme B₁₂.¹⁰

Results and Discussion

General. The general approach used in these studies parallels that employed in several earlier investigations.^{7,8,10,11,12a,13} Co-R bond dissociation energies were deduced from measurements of the kinetics of the homolytic bond dissociation reactions in the presence of a radical trap (in the present study bis(dimethyl-

0002-7863/88/1510-3120\$01.50/0 © 1988 American Chemical Society

For leading references, see: (a) Dolphin, D., Ed. B₁₂; Wiley: New York, 1982. (b) Vitamin B₁₂, Proc. Eur. Symp., 3rd, 1979.
 (2) Halpern, J. Science (Washington, D.C.) 1985, 227, 869 and references

cited therein.

⁽³⁾ Abeles, R. H. Ref 1b, p 373.
(4) Babior, B. M. Acc. Chem. Res. 1975, 8, 376.

⁽⁷⁾ Boolof, B. M. ACC. Chem. Res. 19(5, 8, 3/6).
(5) Golding, B. T. Ref 1a, Vol. 1, p 543 and references cited therein.
(6) (a) Ng, F. T. T.; Rempel, G. L.; Halpern, J. J. Am. Chem. Soc. 1979, 101, 7124.
(b) Ng, F. T. T.; Rempel, G. L.; Halpern, J. Ibid. 1982, 104, 621.
(c) Ng, F. T. T.; Rempel, G. L.; Halpern, J. Inorg. Chim. Acta. 1983, 77, L165.

⁽⁷⁾ Tsou, T. T.; Loots, M.; Halpern, J. J. Am. Chem. Soc. 1982, 104, 623 (8) (a) Geno, M. K.; Halpern, J. J. Am. Chem. Soc. 1987, 109, 1238. (b)
 Geno, M. K.; Halpern, J. J. Chem. Soc., Chem. Commun. 1987, 1052.
 (9) Halpern, J. Bull. Chem. Soc. Jpn. 1988, 61, 13.

⁽¹⁰⁾ Halpern, J.; Kim, S. H.; Leung, T. W. J. Am. Chem. Soc. 1984, 106, 8317.

^{(11) (}a) Finke, R. G.; Hay, B. P. Inorg. Chem. 1984, 23, 3041. (b) Hay,
B. P.; Finke, R. G. J. Am. Chem. Soc. 1986, 108, 4820.
(12) (a) Blau, R. J.; Espenson, J. H. J. Am. Chem. Soc. 1985, 107, 3530.

 ⁽b) Schrauzer, G. N.; Grate, J. H. *Ibid.* 1981, 103, 541. (c) Grate, J. H.;
 Schrauzer, G. N. *Ibid.* 1979, 101, 4601.
 (13) Finke, R. G.; Smith, B. L.; Mayer, B. J.; Molinero, A. A. *Inorg.*

Chem. 1983, 22, 3677.

Table I. Base On-Base Off Equilibria of Organocobalamins (R-B₁₂)

R	pK₂ (25 °C)	temp range of measts, °C	$\Delta H'$ (pH), kcal/mol	$\Delta S'$ (pH), cal/(mol K)
(CH ₃) ₃ CCH ₂	5.07 ± 0.03	10.2-30.6	5.9 ± 0.3 (6.9)	20 ± 1 (6.9)
CH2	3.99 ± 0.05	26.8-54.6	$7.2 \pm 0.3 (7.6)$ $3.9 \pm 0.2 (5.0)$	$20 \pm 1 (7.6)$ $12 \pm 1 (5.0)$
CH2	4.08 ± 0.02			
_0_Сн ⁵	3.44 ± 0.01			

glyoximato)cobalt(II), abbreviated Co(DH)₂, which forms significantly stronger cobalt-carbon bonds than cobalamin)^{8a} according to the scheme of eq 1-3.

$$\mathbf{R} \cdot \mathbf{B}_{12} \xleftarrow{k_1}{k_{-1}} \mathbf{R}^* + \operatorname{cob}(II) \operatorname{alamin}(\mathbf{B}_{12_r})$$
(1)

$$\mathbf{R}^{\bullet} + \operatorname{Co}(\mathbf{DH})_2 \xrightarrow{\kappa_2} \mathbf{R} \cdot \operatorname{Co}(\mathbf{DH})_2$$
(2)

overall:
$$\overline{\text{R-B}_{12} + \text{Co}(\text{DH})_2} \rightarrow \text{R-Co}(\text{DH})_2 + B_{12_r}$$
 (3)

Under pseudo-first-order conditions $(Co(DH)_2 \text{ and } B_{12}, \text{ con-}$ stant) this scheme yields

$$-\frac{1}{[\mathbf{R}-\mathbf{B}_{12}]}\frac{\mathbf{d}[\mathbf{R}-\mathbf{B}_{12}]}{\mathbf{d}t} = k_{\text{obsd}} = \frac{k_1k_2[\operatorname{Co}(\mathbf{DH})_2]}{k_{-1}[\mathbf{B}_{12}] + k_2[\operatorname{Co}(\mathbf{DH})_2]} \quad (4)$$

which rearranges to

$$\frac{1}{k_{\text{obsd}}} = \frac{1}{k_1} + \frac{k_{-1}[\mathbf{B}_{12_r}]}{k_1 k_2 [\text{Co}(\mathbf{DH})_2]}$$
(5)

The Co-R bond dissociation energy $(D_{\text{Co-R}})$ may be derived from the kinetic measurements through the relation

$$D_{\rm Co-R} = \Delta H_1^* - \Delta H_{-1}^* \tag{6}$$

In cases where information is available it has been concluded that the recombination reaction (k_{-1}) is diffusion controlled.¹⁴ Thus, ΔH_{-1}^{\dagger} corresponds to the activation enthalpy for a diffusion-controlled reaction, which, in turn, can be approximated by ΔH^* for viscous flow.¹⁵ For water, this ranges from about 3 kcal/mol at 20 °C to 2 kcal/mol at 100 °C.16

Correction for Base On-Base Off Equilibria. Cobalamins are known to undergo reversible dissociation of the pendant axial 5,6-dimethylbenzimidazole and thus to exist in solution as an equilibrium mixture of the "base-on" [RCo-N], "base-off" [R-Co N], and "protonated-base-off" [RCo NH⁺] forms. The concentrations of these species are related through the equations

$$[R-B_{12}] = [RCo-N] + [RCo N] + [RCo NH^+]$$
(7)

$$K_{\text{on-off}} = [RCo N] / [RCo-N]$$
(8)

$$K_{a} = ([\dot{RCo} \dot{N}] + [\dot{RCo} \dot{N}])[H^{+}]/[\dot{RCo} \dot{N}H^{+}]$$
 (9)

$$K'_{\text{on-off}} = ([R'_{\text{Oo}} N] + [R'_{\text{Oo}} NH^+])/[R'_{\text{Oo}} N]$$
(10)

Values of pK_a , K_{on-off} , and K'_{on-off} , determined as described in the Experimental Section, are listed in Table I. The pK_a of free 5,6-dimethylbenzimidazole (and, hence, presumably of [RCo N]) has been estimated to be ca. 5.5 (25 °C, I = 0.1 M).¹⁷ Thus,



Figure 1. Plots of $[k_{obsd}]^{-1}$ vs $[B_{12,r}]/[Co(DH)_2]$ for the reaction of $(CH_3)_3CCH_2$ - B_{12} with $[Co(DH)_2]$ at 40.5 °C, pH 6.8. (O) 1.2×10^{-3} M $B_{12,r}$; (\Box) 2.6 × 10⁻³ M $Co(DH)_2$; (Δ) 1.5 × 10⁻³ M $Co(DH)_2$; (\diamond) limiting rate with excess $Co(DH)_2 (k_{obsd} \sim k_1)$.

the degree of protonation of [RCo N] is small at pH \gtrsim 7 and, under these conditions, $K_{\text{on-off}} \sim K'_{\text{on-off}}$. In view of the much higher thermal stability of the base-off

and protonated-base-off forms,¹⁸ their contributions to the Co-C bond dissociation reaction (eq 1) may be neglected and the measured values of k_1 must be corrected to reflect only the contributions of the base-on form; i.e.

$$(k_1)_{\text{corr}} = k_1 [\text{R-B}_{12}] / [\text{RCo-N}] = k_1 (1 + K'_{\text{on-off}})$$
 (11)

Dissociation of Co(DH)_2. Because $Co(DH)_2$ undergoes OH--induced irreversible decomposition at the higher temperatures required for the study of some of these reactions, 19 it was necessary, in some cases, to lower the pH of the reaction solutions to the range 4-5. Under these conditions $Co(DH)_2$ undergoes reversible dissociation, according to eq 12 and 13, so that the solution contains a mixture of $Co(DH)_2$ and $Co^{2+.20}$

$$\operatorname{Co}(\mathrm{DH})_2 + 2\mathrm{H}^+ \rightleftharpoons \mathrm{Co}^{2+} + 2\mathrm{DH}_2 \tag{12}$$

$$\frac{[\text{Co}^{2+}][\text{DH}_2]^2}{[\text{Co}(\text{DH})_2][\text{H}^+]^2} = K_{\text{DH}_2}$$
(13)

This equilibrium could be monitored spectrophotometrically by measuring the intensity of the absorbance of $Co(DH)_2$ at 462 nm ($\epsilon_{max} = 1.9 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$). From such measurements, encompassing the initial pH range 4.3–5.0 and initial concentration ranges 1.0×10^{-3} to 6.3×10^{-3} M Co(DH)₂ and 0 to 8.4×10^{-3} M excess DH₂, K_{DH_2} was determined to be $(3.0 \pm 0.7) \times 10^5$ at 25 °C in 0.3 M acetate buffer. This value was insensitive to temperature variations over the range 30-100 °C.

Neopentylcobalamin. (a) In Aqueous Solution at pH 6.8. The reaction of neopentyl- B_{12} with $Co(DH)_2$ in aqueous solution at pH 6.8 (where dissociation of $Co(DH)_2$ according to eq 12 is negligible) was found to exhibit the stoichiometry of eq 14.

neopentyl- B_{12} + Co(DH)₂ \rightarrow neopentyl-Co(DH)₂ + B_{12r} (14)

This is confirmed by the following observations: (i) The UVvisible spectrum of a solution containing initially 1.4×10^{-4} M each of neopentyl-B₁₂ and Co(DH)₂ (0.1 M phosphate buffer, pH 6.8), after reacting at 80 °C for 5 h, matched that of a synthetic solution containing 1.4×10^{-4} M each of neopentyl-Co(DH)₂ and \mathbf{B}_{12r} (ii) In several experiments it was found by ¹H NMR that,

⁽¹⁴⁾ Halpern, J. Acc. Chem. Res. 1982, 15, 238 and references cited therein (15) Schuh, H. H.; Fischer, H. Helv. Chim. Acta 1978, 61, 2130 and

references cited therein. (16) Deduced from data in: Handbook of Chemistry and Physics, 60th

Ed.; CRC: Boca Raton, FL, 1979.

⁽¹⁷⁾ Brown, K. L.; Hakimi, J. M.; Nuss, D. M.; Montejano, Y. D.; Ja-

 ⁽¹⁾ Bioman, R. E., Hannin, 1984, 23, 1463.
 (18) (a) Chemaly, S. M.; Pratt, J. M. J. Chem. Soc., Dalton Trans. 1980, 2274. (b) Schrauzer, G. N.; Grate, J. H. J. Am. Chem. Soc. 1981, 103, 541.

⁽¹⁹⁾ Simandi, L. I.; Nemeth, S.; Bodo-Zahonyi, E. Inorg. Chim. Acta
1980, 45, L143 and references cited therein.
(20) (a) Adin, A.; Espenson, J. H. Inorg. Chem. 1972, 11, 686. (b) Gjerde,
H. B.; Espenson, J. H. Organometallics 1982, 1, 435.

Table II. Stoichiometries of the Reactions of Neopentylcobalamin with "Co(DH)2"

temp, time,		time.	$10^3 \times \text{init concn, M}$			$10^3 \times \text{final}$			
pН	°Ċ	h	(CH ₃) ₃ CCH ₂ -B ₁₂	"Co(DH)2"	B _{12a}	(CH ₃) ₃ CCH ₂ -B ₁₂	(CH ₃) ₃ CCH ₂ -Co(DH) ₂	$\sum \mathbf{B}_{12}$	$\sum (CH_3)_3 CCH_2$
6.8	45	1.5	1.1	7.8	1.584	0	1.29 ^b	1.58	1.29
6.8	80	5	3.5	5.0	3.6	0	3.6	3.6	3.6
6.8	45	1.5	1.1	7.8	1.584	0	1.29	1.58	1.29
4.3	60	29	1.0	1.5	0.9	0.1	0.89	1.0	0.99
4.3	60	9	1.0	5.0	1.03	0	1.06	1.03	1.06
4.3	70	18	1.0	1.6	0.91	0.1	0.86	1.01	0.96

 a 0.44 × 10⁻³ M B_{12a} present initially; Δ [B_{12a}] = 1.14 M. b 0.30 × 10⁻³ M (CH₃)₃CCH₂-B₁₂ present initially; Δ [(CH₃)₃CCH₂-B₁₂] = 0.99 M.

following reaction of neopentyl- B_{12} with $Co(DH)_2$ (and after passing O_2 through the solution to oxidize the paramagnetic B_{12_r} product to B_{12_a}), the concentrations of neopentyl- $Co(DH)_2$) and B_{12_a} correspond to the concentration of reacted neopentyl- B_{12} (Table II).

The kinetics of reaction 14 were examined at 40.5 °C. The results of experiments encompassing the initial concentration ranges, 3.0×10^{-4} to 1.2×10^{-3} M B₁₂, and 2.0×10^{-3} to 6.8×10^{-3} M Co(DH)₂, conformed to eq 5 as depicted in Figure 1. These measurements yield $k_1 = (1.65 \pm 0.15) \times 10^{-3} \text{ s}^{-1}$ and $k_{-1}/k_2 = 3.5 \pm 0.3$.

 k_1 was determined at other temperatures (25-45 °C) from measurements under limiting conditions (large excess Co(DH)₂, no added B₁₂,) such that $k_2[Co(DH)_2] \gg k_{-1}[B_{12}]$; hence $k_{obsd} \simeq k_1$. The results, listed in Table II, yield $\Delta H_1^* = 23.1 \pm 0.7$ kcal/mol and $\Delta S_1^* = 2 \pm 2$ cal/(mol K). These values are in good agreement with those reported in an earlier study using O₂ as radical scavenger, $\Delta H_1^* = 23.4 \pm 0.2$ kcal/mol, $\Delta S_1^* = 1.6 \pm 0.1$ cal/(mol K).^{12b} After correction of the rate constants in Table II according to eq 11, $(\Delta H_1^*)_{corr} = 26.7 \pm 1.2$ kcal/mol and $(\Delta S_1^*)_{corr} = 15 \pm 5$ cal/(mol K). Subtracting $\Delta H_{-1}^* \sim 3$ kcal/mol leads to a value of 24 ± 2 kcal/mol for the neopentyl-B₁₂ bond dissociation energy, $D_{neopentyl-B_{12}}$. (b) In Aqueous Solution at pH 4.3. To establish whether Co-

(b) In Aqueous Solution at pH 4.3. To establish whether Co- $(DH)_2$ also was effective as a radical trap at lower pHs where it was partially or largely dissociated into Co²⁺ and DH₂ (eq 12),^{10,11,20} the reaction between neopentyl-B₁₂ and Co(DH)₂ also was examined at pH 4.3.

The data in Table II demonstrate that, under these conditions, the stoichiometry of eq 14 still is obeyed, transfer of the neopentyl radical from B_{12} to $Co(DH)_2$ being essentially quantitative. The kinetics of the reaction at 35.2 °C were examined with initial solution compositions ranging from 3.0×10^{-3} to 6.9×10^{-3} M "Co(DH)₂" and 3.1×10^{-4} to 1.2×10^{-3} M B_{12r} . Under these conditions, the reaction was found to obey the rate law of eq 15,

$$k_{\text{obsd}} = \frac{k_1' k_2' [\text{``Co(DH)}_2"]}{k_{-1}' [\text{B}_{12_r}] + k_2' [\text{``Co(DH)}_2"]}$$
(15)

which is essentially the same as eq 4 but with ["Co(DH)₂"] = $[Co(DH)_2] + [Co^{2+}]$. These measurements yield the values $k_1' = (1.7 \pm 0.3) \times 10^{-4} \text{ s}^{-1}$ and $k_{-1}'/k_2' = 13 \pm 3$ (compared with $k_{-1}/k_2 = 3.5$ at 40.5 °C). Thus, we estimate the efficiency of "Co(DH)₂" as a radical trap at pH 4.3 to be about one-quarter that at pH 6.8 where dissociation of Co(DH)₂ is negligible.

Two interpretations may be accorded to the trapping of the neopentyl radicals at pH 4.3:

(i) Trapping is effected wholly by undissociated $Co(DH)_2$ which is present under these conditions, in accord with eq 2.

(ii) Trapping also is effected by the Co^{2+} ion to yield a transient neopentyl- Co^{2+} adduct which reacts rapidly with DH₂ to form the observed neopentyl- $Co(DH)_2$ product (eq 16 and 17). (Because of the reversibility of reaction 16, Co^{2+} is not expected to be an effective radical trap in the absence of DH₂.)

$$\mathbf{R}^{\bullet} + \mathbf{Co}^{2+} \stackrel{k_2}{\longleftrightarrow} [\mathbf{R} - \mathbf{Co}^{2+}]$$
(16)

$$[\mathbf{R}-\mathbf{Co}^{2+}] + 2\mathbf{DH}_2 \xrightarrow{\text{tast}} \mathbf{R}-\mathbf{Co}(\mathbf{DH})_2 + 2\mathbf{H}^+ \qquad (17)$$

Because more than 90% of the $Co(DH)_2$ is dissociated into Co^{2+} and DH_2 at pH 4.3, it seems unlikely that the relatively efficient radical trapping under these conditions is due entirely to undissociated $Co(DH)_2$. Accordingly, we favor the interpretation of eq 16 and 17, which invokes radical trapping also by Co^{2+} . Indeed, Kochi²¹ has reported evidence for such trapping of alkyl radicals by Co^{2+} and formation of transient $[R-Co^{2+}]$ adducts. In any event, it is clear that " $Co(DH)_2$ " does function as an efficient radical trap at pH 4.3 and can be used to determine the kinetics of alkyl-B₁₂ bond dissociation under these conditions. This point is noteworthy, since the appropriateness of such a procedure has been questioned.^{11b}

(c) In Ethylene Glycol Solution. Because of discrepancies in earlier measurements on the thermal decomposition of (5'-deoxyadenosyl)cobalamin in aqueous and in ethylene glycol solutions,^{10,11} it was of interest to extend these studies on neopentyl-B₁₂, where both the stoichiometry and mechanisms are unequivocal, to ethylene glycol.

Decomposition of neopentyl- B_{12} in ethylene glycol over the temperature range 20-50 °C, under limiting conditions (initially ca. 2×10^{-4} neopentyl-B₁₂, 5×10^{-3} M Co(DH)₂), exhibited excellent pseudo-first-order kinetics and yielded the values of k_1 listed in Table III and the activation parameters $\Delta H_1^* = 26.8$ ± 0.3 kcal/mol and $\Delta S_1^* = 13 \pm 1$ cal/(mol K) (vs 23.1 ± 0.7 kcal/mol and 2 ± 2 cal/(mol K) in aqueous solution at pH 6.8). Preliminary measurements yield estimates of $10 \pm 1 \text{ kcal/mol}$ and 34 ± 5 cal/(mol K) for $\Delta H'$ and $\Delta S'$, respectively, for neopentyl- B_{12} in ethylene glycol. Using these values to correct for the base on-base off equilibrium yields $(\Delta H_1^{\dagger})_{corr} = 33 \pm 2$ kcal/mol and $(\Delta S_1^*)_{corr} = 35 \pm 10 \text{ cal/(mol K)}$ (vs 26.7 ± 1.2 kcal/mol and 15 ± 5 cal/(mol K) in aqueous solution at pH 6.8). The differences, $\Delta \Delta H_1^* = 3.7 \pm 1$ kcal/mol and $(\Delta \Delta H_1^*)_{corr} =$ 6 ± 3 kcal/mol, parallel those found earlier for the homolytic Co-C bond dissociation reactions of several other organocobalt compounds (benzyl-Co(DH)₂ and benzyl-Co(porphyrin) in toluene and ethylene glycol $(\Delta\Delta H^* = 4-6 \text{ kcal/mol})$,^{8a} presumably reflecting the higher viscosity of ethylene glycol compared with that of water and toluene. Accordingly, a difference of several kcal/mol between ΔH^* for the homolytic dissociation of the Co-C bond in 5'-deoxyadenosyl- B_{12} in ethylene glycol and in water is not unexpected.10,11

(Cyclopentylmethyl)cobalamin (c- $C_5H_9CH_2-B_{12}$). (a) In the Absence of Co(DH)₂. Since higher temperatures (60–80 °C) were required for the decomposition of c- $C_5H_9CH_2-B_{12}$ to proceed at conveniently measurable rates, this reaction was studied at pH 5.0. Under these conditions the extent of protonation of c- $C_5H_9CH_2-B_{12}$ is less than 10%. At the same time "Co(DH)₂" was found to be stable.

At 74 °C, c-C₅H₉CH₂-B₁₂ decomposed in the absence of "Co(DH)₂", with a half-life of ca. 50 min, to form B₁₂, in quantitative yield (spectrophotometric).²² The kinetic behavior, described below, and the pattern of organic products are accommodated by Scheme I.

In the absence of "Co(DH)₂", the principal organic product, after thermolysis of $c-C_5H_9CH_2-B_{12}$ for 6 h at 75 °C (identified by ¹H NMR following extraction with C₆D₆), was found to be

⁽²¹⁾ Kochi, J. K.; Rust, F. F. J. Am. Chem. Soc. 1961, 83, 2017.

⁽²²⁾ B_{12_a} decomposes spontaneously in acidic solutions at high temperatures to form $B_{12_r}^{11b}$ However, the rate is much lower than the time scale of this reaction: $t_{1/2} \sim 360$ min at 100 °C, pH 4.3. Hence it would appear that B_{12_r} is the primary product of the reaction.

Table III.	Kinetic Data	for Homoly	ic $R-B_{12}$	Bond	Dissociation ⁴
------------	--------------	------------	---------------	------	---------------------------

			uncorrected			corrected ^b		
R	pН	temp, °C	$\frac{10^4k_1}{s^{-1}}$	ΔH_1^* , kcal/mol	$\frac{\Delta S_1^*}{\operatorname{cal}/(\operatorname{mol} K)}$	$\frac{10^4 k_1}{s^{-1}}$	ΔH_1^* , kcal/mol	ΔS_1^* , cal/(mol K)
(CH ₁) ₁ CCH ₂	6.8	25.4	2.12			4.70		
(CH ₁) ₁ CCH ₂	6.8	30.0	3.69			8.97		
(CH ₁),CCH ₂	6.8	30.7	4.26			10.5		
(CH ₃) ₃ CCH ₃	6.8	35.5	7.45			20.3		
(CH ₃),CCH ₂	6.8	35.2	8.46			23.0		
(CH,),CCH,	6.8	40.0	14.7			44.0		
(CH ₃),CCH ₃	6.8	40.6	16.5			49.4		
(CH ₃),CCH ₃	6.8	45.4	24.8			81.8		
(CH ₃),CCH ₃	6.8	45.6	24.9			82.1		
(CH ₃),CCH ₃	6.8			23.1 ± 0.7	2 ± 2		26.7 ± 1.2	15 ± 5
(CH ₃),CCH ₃	с	20.4	0.40			0.92		
(CH ₃),CCH ₃	с	20.8	0.42			0.90		
(CH ₃),CCH ₃	с	30.0	1.65			5.43		
(CH),CCH	с	30.8	1.80			5.44		
(CH ₃),CCH ₂	с	40.0	6.82			29.1		
(CH ₁),CCH ₂	с	40.6	7.71			33.6		
(CH ₃),CCH ₃	с	50.0	27.3			172		
(CH ₃),CCH ₃	с			$26.8 \pm 0.3^{\circ}$	$13 \pm 1^{\circ}$		33 ± 2^{c}	35 ± 10°
c-C.HCH	5.0	58	0.26			0.47		
c-C,Ho-CH	5.0	63	0.45			0.86		
c-C,H,-CH	5.0	68	0.98			1.95		
c-CiHi-CH	5.0	73	1.30			2.70		
c-C,H,-CH,	5.0	78	2.49			5.35		
c-C ₅ H ₂ -CH ₂	5.0			25.2 ± 1.7	-3 ± 5		26.8 ± 2	3 ± 6

^a In aqueous solution unless designated otherwise. ^bCorrected for base on-base off equilibrium; data for base-on form. ^cIn ethylene glycol.

Scheme I Possible Pathways for the Decomposition of $c-C_5H_9CH_2-B_{12}$ in Acetate Buffer



1-methylcyclopentane (together with unidentified products associated with resonances at 1.71, 1.40, 1.02, and 1.01 ppm). The absence of methylcyclopentane under these conditions is not unexpected. Since the formation of the cyclopentylmethyl radical is reversible, and of methylenecyclopentane irreversible, most of the product would be expected to be derived from the latter. Failure to observe methylenecyclopentane also is expected. In a control experiment methylenecyclopentane $(5 \times 10^{-3} \text{ M})$ in 0.1 M pH 5 acetate buffer was heated at 75 °C for 1 h and found to be unstable. In addition to unreacted methylenecyclopentane the ¹H NMR spectrum of the C_6D_6 extracts exhibited resonances at 2.51, 2.33, and 2.02 ppm attributable to methylcyclopentene and resonances at 1.71, 1.40, 1.03, and 1.02 ppm, corresponding to those of the unidentified products of the thermal decomposition of $c-C_5H_9CH_2-B_{12}$. Thus, it would appear that the latter arise from further reactions of methylenecyclopentane formed according



Figure 2. Dependence of k_{obsd} on $[B_{12_r}]^{-1}$ for the decomposition of c-C₅H₂-B₁₂ (pH 5.0).

Fable IV. Kinetic Data fo	r (Cyclopentylmet)	hyl)cobalamin (pH 5.0)
----------------------------------	--------------------	------------------------

temp, °C	$10^4 k_0, \mathrm{s}^{-1}$	$10^4 (k_{obsd})_{lim}, s^{-1}$	$10^4 k_1$, s ⁻¹
58	0.20	0.46	0.26
63	0.37	0.82	0.45
68	0.63	1.61	0.98
73	1.15	2.46	1.30
78	1.92	4.41	2.49

to Scheme I. Such possible products include the hydroxide or acetate of the 1-methylcyclopentyl carbonium ion or of the addition product of the latter to methylenecyclopentane.

According to Scheme I it should be possible to suppress the homolytic dissociation pathway by addition of B_{12} , and, in this way, to determine, in the limit, the rate constant of the olefin elimination pathway, k_0 . The plots of k_{obsd} vs $[B_{12}]^{-1}$ in Figure 2 are consistent with this and yield the values of k_0 listed in Table IV and the activation parameters $\Delta H_0^* = 25.4 \pm 0.3$ and $\Delta S_0^* = -4 \pm 2$ cal/(mol K).

(b) In the Presence of Co(DH)₂. Scheme I anticipates that, in the presence of "Co(DH)₂", the cyclopentylmethyl radical produced in the homolytic dissociation step should be trapped to form $c-C_5H_9CH_2-Co(DH)_2$. To test this, $c-C_5H_9CH_2-B_{12}$ (1.0



[B₁₂]/[Co(DH)₂]

Figure 3. Plots of $(k_{obsd} - k_0)^{-1}$ vs $[B_{12,}]/[\text{``Co(DH)}_2)$ "] for the decomposition of $c\text{-}C_5H_9CH_2\text{-}B_{12}$ at 73.0 °C, pH 5.0. (O) 4.0 × 10⁻³ M "Co(DH)₂"; (\Box) 2.0 × 10⁻³ M "Co(DH)₂"; (Δ) 1.0 × 10⁻³ M B₁₂; (\diamond) limiting value with excess "Co(DH)₂" ($k_{obsd} - k_0 \approx k_1$).

× 10⁻³ M in D₂O containing 0.1 M pD 5.0 CD₃COO⁻ buffer) was heated at 75 °C for 20 min in the presence of a large excess (5 × 10⁻³ M) of "Co(DH)₂" followed by passing O₂ through for 30 min to oxidize B₁₂ and unreacted "Co(DH)₂". The ¹H NMR spectrum of the resulting solution contained signals corresponding to a quantitative yield (1.0 × 10⁻³ M) of B₁₂ and 4.7 × 10⁻⁴ M c-C₃H₉CH₂-Co(DH)₂. Thus it is concluded that 47% of the c-C₃H₉CH₂-B₁₂ decomposes by the homolytic pathway under these conditions.

The kinetic scheme for the reaction of $c-C_5H_9CH_2-B_{12}$ in the presence of "Co(DH)₂" and B₁₂, corresponds to

$$CH_2 - B_{12}$$

$$CH_2 - B_{12}$$

$$CH_2 - B_{12}$$

$$CH_2 - CH_2 - C$$

The resulting rate law

$$-\frac{1}{[c-C_{5}H_{9}CH_{2}-B_{12}]}\frac{d[c-C_{5}H_{9}CH_{2}-B_{12}]}{dt} = k_{obsd} = k_{0} + \frac{k_{1}k_{2}'["Co(DH)_{2}"]}{k_{0} + \frac{k_{1}k_{2}'["Co(DH)_{2}"]}{k_{-1}[B_{12}] + k_{2}'["Co(DH)_{2}"]}$$
(19)

rearranges to

$$(k_{\rm obsd} - k_0)^{-1} = \frac{1}{k_1} + \frac{k_{-1}[\mathbf{B}_{12,r}]}{k_1 k_2' [\text{``Co(DH}_{2,r}]]}$$
(20)

The rates of the reaction of c-C₅H₉CH₂-B₁₂ were monitored spectrophotometrically at 73 °C and pH 5.0 in a series of experiments encompassing the initial concentration ranges 1.0×10^{-4} to 2.0×10^{-4} M B₁₂, and 6.5×10^{-4} to 4.0×10^{-3} M "Co-(DH)₂". The kinetics conformed to pseudo-first-order behavior, yielding values of k_{obsd} . Combining these with the values of k_0 in Table III leads to the linear plot of $(k_{obsd} - k_0)^{-1}$ vs [B₁₂,]/["Co(DH)₂"] depicted in Figure 3 in accord with eq 20. These measurements yield $k_1 = (1.30 \pm 0.11) \times 10^{-4}$ s⁻¹ and $k_{-1}/k_2' = 3.8 \pm 0.3$.

In the limit when $k_2'[\text{``Co(DH)}_2"] \gg k_{-1}[B_{12,}]$, eq 19 reduces to $k_{obsd} = k_0 + k_1$. Values of k_{obsd} determined under these limiting conditions at temperatures ranging from 58 to 78 °C are listed in Table IV together with the values of k_0 determined earlier and the resulting values of k_1 obtained by difference. These yield ΔH_1^* = 25.2 ± 1.7 kcal/mol and $\Delta S_1^* = -3 \pm 5$ cal/(mol K). Correcting for the base on-base off equilibrium yields (ΔH_1^*)_{corr} = 26.8 ± 2 kcal/mol and (ΔS_1^*)_{corr} = 3 ± 6 cal/(mol K). Thus,



Figure 4. IUPAC numbering scheme for carbon atoms of alkylcobalamins (X = alkyl group).

we deduce the Co-C bond dissociation energy to be 24 ± 2 kcal/mol.

A knowledge of the values of k_0 and of k_1 and k_{-1}/k_2' permits prediction of the product distribution. Under the limiting conditions just described, the yield of c-C₅H₉CH₂-Co(DH)₂ should correspond to k_1/k_{obsd} , i.e., $1.30 \times 10^{-4} \text{ s}^{-1}/2.46 \times 10^{-4} \text{ s}^{-1} = 0.53$ at 73 °C. This is close to the yield of 47% at 75 °C reported above.

It is striking that the activation parameters for k_0 and k_1 are virtually identical. Thus, $\Delta H_0^* = 25.3 \text{ kcal/mol}$, $\Delta H_1^* = 25.2 \text{ kcal/mol}$; $\Delta S_0^* = -4 \text{ cal/(mol K)}$, $\Delta S_1^* = -3 \text{ cal/(mol K)}$. This suggests that the two processes share a common rate-determining step, namely homolytic Co-C bond dissociation to yield a geminate $\overline{c-C_5H_9CH_2^*,B_{12}}$, radical pair. The two products, i.e., radical and olefin, could then arise from competing geminate pair separation and cage H-atom transfer, i.e., $\overline{c-C_5H_9CH_2^*,B_{12}} \rightarrow \overline{c}$. $\overline{C_5H_8=}CH_2 + H-B_{12}$; $H-B_{12} \rightarrow B_{12}$, $\pm 1/_2H_2$. We have previously proposed a similar mechanism for the olefin-forming reaction PhCH(CH₃)-Co(DH)₂ \rightarrow Co(DH)₂ + PhCH==CH₂ + $1/_2H_2$.^{6b} Other olefin-elimination reactions have been similarly interpreted.²³

(Cyclohexylmethyl)cobalamin (c- $C_6H_{11}CH_2-B_{12}$). The behavior of c-C₆H₁₁CH₂-B₁₂ closely paralleled that of c-C₅H₉CH₂-B₁₂. The two compounds exhibited similar pK_a 's (4.06 and 3.99, respectively). A pH 6.8 solution of $c-C_6H_{11}CH_2-B_{12}$ was stable at room temperature for at least 24 h, and higher temperatures were required to observe decomposition. At pH 5.0 (in the absence of "Co(DH)₂"), c-C₆H₁₁CH₂-B₁₂ decomposed at 75 °C with a half-life of ca. 70 min to form B_{12_r} in quantitative yield (spectrophotometric). Following decomposition at 75 °C for 6 h the ¹H NMR spectrum of a C_6D_6 extract contained signals due to methylenecyclohexane and 1-methylcyclohexene, in the ratio 5:7, together with minor unidentified resonances at 1.68, 1.415, 1.04, and 1.02 ppm. The last three signals also could be observed in the spectrum of a sample of authentic methylenecyclohexane decomposed under the same conditions. No methylcyclohexane could be detected.

As in the case of $c-C_5H_9CH_2-B_{12}$, the decomposition of $c-C_6H_{11}CH_2-B_{12}$ (in the absence of "Co(DH)₂") was inhibited by added B_{12} ; thus, at 75 °C, pH 5, $k_{obsd} = 1.39 \times 10^{-4}$, 1.29×10^{-4} , and $1.14 \times 10^{-4} s^{-1}$ at 5.0×10^{-4} , 1.0×10^{-3} , and 2.0×10^{-3} M

⁽²³⁾ Baldwin, D. A.; Betterton, E. A.; Chemaly, S. M.; Pratt, J. M. J. Chem. Soc., Dalton Trans. 1985, 1613.

Table V. ¹H NMR Spectra of Alkylcobalamins^a

		neopentyl	cyclopentylmethyl			cyclohexyl- methyl	tetrahydro- furfuryl	(tetrahydro-2 <i>H</i> - pyryl)methyl
assigt [*]	ethyl ^c	pD 4.3 ^d	pD 6.8*	pD 5.0 ^d	pD 4.3 ^d	pD 5.0 ^d	pD 6.8*	(pD 6.8)
B(2)H	6.991	9.190	7.110	7.428	8.236	7.744	6.960, 6.984	6.983, 6.965
B(4)H	6.245	7.413	6.376	6.537	6.976	6.696	6.260, 6.360	6.235, 6.222
B(7)H	7.166	7.510	7.162	7.215	7.354	7.264	7.176, 7.164	7.164, 7.157
B(10)H ₃	2.212		2.220	2.191	2.200/2.180		2.216	2.219
$B(11)H_{3}$	2.212	2.194/2.160	2.220	2.191		2.188/2.180	2.216	2.219
R(1)H	6.245	6.500	6.250	6.283	6.390	6.330	6.260	6.265
$Pr(3)H_3$	1.209	1.172	1.220	1.205	1.185	1.196	1.218	1.224
C(10)H	6.041	6.978	6.128	6.240	6.552	6.380	5.988, 6.048	6.035, 5.995
C(20)H ₁	0.520	0.614	0.438	0.483	0.604	0.528	0.504, 0.418	0.508, 0.454
C(25)H ₁	1.386	(1.624) ^g	1.400	(1.428)8	(1.624) ^g	(1.456)8	(1.420)8	$(1.467)^{g}$
C(35)H ₃	2.495	2.384	2.504	2.483	2.440	2.440	2.500	2.507
C(36)H ₃	1.778	1.838	1.820	1.815	1.820	1.820	1.786, 1.858	1.833, 1.783
C(46)H ₃	1.043	1.210	1.244	1.226	1.180	1.234	1.178, 1.280	1.289, 1.213
C(47)H ₃	1.441	1.418	1.344	1.356	1.384	1.366	(1.350)8	$(1.367, 1.357)^{g}$
C(53)H ₃	2.476	2.384	2.480	2.463	2.430	2.450	2.500	2.493
C(54)H ₁	1.386	(1.392)8	1.400	(1.397)8	(1.392)8	(1.400)8	(1.396)8	(1.399, 1.393)*
$\mathbf{X}(1)^{h}$	0.846/0.566	i/0.920				1		
$\mathbf{X}(2)^h$	-0.594		0.360	-0.430	-0.540	-1.007	1.070, 0.991	
$X(3)^h$		0.086	(0.380, ^j 0.560 ^j	(0.376, ^j 0.536 ^j	(0.312, 0.412)	1	0.980, 0.560, 0.380	0.9960, 0.820
$\mathbf{X}(4)^h$			0.908, ^j 0.991 ^j	0.896, 0.946	0.804, 0.816	1		0.705
$\mathbf{X}(5)^h$			1.070	1.056, ^j 1.137 ^j	0.980, 1.090	1.642/1.558	3.58, 3.40	
$\mathbf{X}(6)^h$			-	•		1		3.69, 3.46, 3.34, 2.93
$\mathbf{X}(7)^h$						1		

^a Chemical shifts (δ) downfield from (trimethylsilyl)propionic acid (TSP). ^b Based on numbering scheme in Figure 4 (cf. ref 27 and 28). ^c From ref 27. ^d0.1 M acetate buffer in D₂O at 36 ^oC. ^c0.1 M phosphate buffer in D₂O at 36 ^oC. ^e0.1 M phosphate buffer in D₂O at 33 ^oC. ^eTentative assignments. ^kAlkyl group carbon atoms numbered in sequence starting from Co-bonded carbon atom. ⁱNot observed due to overlap with other resonances. ^jThese single-proton resonances belonging to the cyclopentylmethyl group could not be assigned. ^kOverlap with adjacent large peak. ^jSignals for these 10 protons appear between 0.9 and 0.3 ppm.

 $B_{12,r}$, respectively. Extrapolation to $[B_{12_r}]^{-1} = 0$ yielded $k_0 = 1.1 \times 10^{-4} \text{ s}^{-1}$, close to the value $1.15 \times 10^{-4} \text{ s}^{-1}$ at 73 °C previously determined for c-C₅H₉CH₂-B₁₂. Because of the similarity of behavior to the latter, the study of c-C₆H₁₁CH₂-B₁₂ was not pursued further.

(Tetrahydrofurfuryl)cobalamin (THF- B_{12}). The study of this compound was of interest because it incorporates an analogue of one of the structural elements (the deoxyribose moiety) of (5'deoxyadenosyl)cobalamin. THF- B_{12} was only moderately stable in aqueous solution at pH 6.8, decomposition being detectable after 10 h at 15 °C under an inert atmosphere. Decomposition was accelerated by acid, so that the half-life for decomposition at 25 °C was ca. 60 min at pH 5.0 and less than 5 min at pH 2.2. The product of decomposition was aquocob(III)alamin (B_{12}). Thus, it appears that decomposition is induced by protonation of the oxygen atoms in accord with eq 21

$$\begin{array}{c} \overset{O}{\longrightarrow} CH_2 \xrightarrow{H^+} B_{12} \xrightarrow{H^-} CH_2 \xrightarrow{H^+} B_{12} \xrightarrow{H^-} CH_2 \xrightarrow{H^+} B_{12} \xrightarrow{H^-} CH_2 \xrightarrow{H^+} B_{12a} \xrightarrow{H^+} CH_2 \xrightarrow{H^+} CH_2 \xrightarrow{H^+} B_{12a} \xrightarrow{H^+} CH_2 \xrightarrow{H$$

Because of this instability it proved impossible to study the homolytic Co–C bond dissociation reaction or to measure the pK_a of THF-B₁₂. It is remarkable that the stability of the structurally related (5'-deoxyadenosyl)cobalamin is so much greater than that of THF-B₁₂ notwithstanding probably more severe steric crowding.

((Tetrahydro-2H-pyryl)methyl)cobalamin (THP-B₁₂). The behavior of this compound paralleled that of THF-B₁₂, although it exhibited somewhat greater stability. In aqueous solution at pH 6.8 it was stable for ca. 2 days at 25 °C. The half-life for its decomposition (to B₁₂) at 36 °C was 36 h at pH 4.3 and 12.5 h at pH 2.5. Decomposition was sufficiently slow to permit measurement of the pK_a (3.44 ± 0.01) but not to study the homolytic Co-C bond dissociation.

Experimental Section

General. All manipulations were performed in the dark under N_2 with rigorous exclusion of air.

was measured relative to a standardized Orion Ag/AgCl electrode using a Radiometer PHM 64 pH meter.

Starting Materials. Vitamin B_{12} (cyanocobalamin), dimethylglyoxime, cyclopentylmethanol, tetrahydrofurfuryl alcohol, methylcyclopentane, methylenecyclopentane, and methylcyclohexane were used without purification. Neopentyl bromide, cyclohexylmethyl iodide, and 2-(bromomethyl)tetrahydro-2*H*-pyran were purified by vacuum distillation. Cyclopentylmethyl iodide and tetrahydrofurfuryl iodide were prepared by the procedure of Rydon.²⁴

Cobalamins. Cob(II)alamin (B_{12_1}) ,²⁵ hydroxocob(III)alamin (B_{12_a}) ,²⁶ and neopentylcobalamin^{12b} were prepared by literature procedures.

(Cyclopentylmethyl)cobalamin. B_{12_4} (0.25 g, 0.186 mmol) was dissolved in 10 mL of N₂-saturated 5% NH₄Br methanol solution. Zn dust (1 g) was added to the cherry red solution and the color changed immediately to dark green, indicating the formation of B_{12_4} . Cyclopentylmethyl iodide (0.03 mL, 0.285 mmol) was injected into the B_{12_5} solution, resulting in orange (cyclopentylmethyl)cobalamin within a few minutes. After the Zn dust was removed by centrifugation, the orange supernatant was mixed with 100 mL of 1 M aqueous HCl solution in a separatory funnel. The organocobalamin was extracted into 15 mL of a 1:1 chloroform-phenol mixture and precipitated by addition of 50 mL of ether. The orange precipitate was collected by centrifugation and washed repeatedly with ether until the phenol was completely removed. The recovered solid was dried in vacuo overnight. Yield, 0.18 g (0.127 mmol, 68% based on B_{12_6}). UV-visible, λ_{max} (10⁻³ ϵ_{max}): pH 7.6, 522.5 (8.58), 450.0 (3.95); pH 5.0, 518.0 (7.65), 449.0 (4.115), 380.0 (7.50).

(Cyclohexylmethyl)cobalamin. B_{12_a} (0.1 g, 0.074 mmol) was dissolved in 5 mL of N₂-saturated 5% NH₄Br methanol solution. Zn dust (0.4 g) was added to the cherry red solution, resulting in an immediate color change to greenish black, indicating formation of B_{12_4} . Addition of 0.03 mL of cyclohexylmethyl bromide resulted in a color change to red-orange within a few minutes. After centrifugation of the orange suspension, the supernatant was mixed with 50 mL of 1 M aqueous HCl. The organocobalamin was extracted into 10 mL of a 1:1 mixture of chloroform and phenol and precipitated by addition of 100 mL of ether. The orange solid was recovered by centrifugation and washed with ether until phenol was completely removed. The product was dried in vacuo overnight. Yield, 0.06 g (0.042 mmol, 57% based on B_{12_8}). UV-visible, λ_{max} (10⁻³ ϵ_{max}): pH 6.8, 518 (6.70), 440 (4.14); pH 5.0, 518 (6.50), 442 (4.25), 380 (6.99).

(Tetrahydrofurfuryl) cobalamin. B_{12_s} (0.25 g, 0.19 mmol) was dissolved in 10 mL of N₂-saturated methanol. Addition of 0.05 g of NaBH₄ (0.13 mmol) to the cherry red solution resulted in an immediate color change to dark green, indicating the formation of B_{12_s} . Tetrahydrofurfuryl iodide (0.03 mL, 0.29 mmol) was injected into the brown solution, resulting in a color change to pink. After 1 h of stirring to complete the reaction, 80 mL of water was added, and the organocobalamin was

Instrumentation. ¹H NMR spectra were recorded in D_2O or C_6D_6 solution using The University of Chicago 500-MHz spectrometer. Chemical shifts are referenced with respect to internal sodium 3-(trimethylsilyl)propionate- d_4 (TSP- d_4) or TMS, respectively. The solvent peak was suppressed by irradiating the water signal. UV-visible spectra were recorded on a Perkin-Elmer Lambda 7 or Lambda 5 spectrophotometer, equipped with a thermostated cell connected to a circulating constant-temperature bath (±0.2 °C). The pH of the buffer solutions

⁽²⁴⁾ Rydon, N. H. Org. Synth. 1971, 51, 44.

⁽²⁵⁾ Blaser, H. U.; Halpern, J. J. Am. Chem. Soc. 1980, 102, 1684.

⁽²⁶⁾ Bernhaur, K.; Wagner, O. Biochem. Z. 1963, 357, 366.

extracted into 15 mL of 1:1 phenol-chloroform. Ether (150 mL) was added to precipitate the organocobalamin. The pink solid was recovered by centrifugation, washed repeatedly with ether, and dried under vacuum. Yield, 0.15 g (0.11 mmol, 56% based on B_{12_4}). UV-visible, λ_{max} (10⁻³ ϵ_{max}): pH 6.8, 520 (8.41), 368 (10.80), 355 (13.67).

((Tetrahydro-2H-pyryl)methyl)cobalamin. B₁₂ (0.25 g, 0.19 mmol) was dissolved in 10 mL of methanol. The resulting cherry red solution was saturated with N₂, and 0.05 g of NaBH₄ (0.13 mmol) was added to produce a dark green solution of B_{12} . Addition of 0.03 mL of 2-(bro-momethyl)tetrahydro-2*H*-pyran resulted in an immediate color change to pinkish red. After 1 h of stirring under N_2 to complete the reaction, the solution was mixed with 100 mL of water, and the organocobalamin was extracted into 15 mL of 1:1 phenol-chloroform. Ether (150 mL) was added to the phenol-chloroform layer to precipitate the organocobalamin. The pink-red solid was recovered by centrifugation and washed with ether. The product was dried in vacuo overnight. UV-visible, λ_{max} $(10^{-3}\epsilon_{max})$: pH 6.8, 521 (6.76), 372 (9.02), 340 (11.72).

The ¹H NMR spectra of the cobalamins are detailed in Table V. Chemical shift assignments are based on comparisons with the published spectra of ethylcobalamin²⁷ and (5'-deoxyadenosyl)cobalamin²⁸ and on the pH dependencies of the resonances.

Organobis(dimethylglyoximato)cobalt(III) Complexes ([R-Co(DH)2]). Dimethyl sulfide adducts of neopentyl-Co(DH)₂ and c-C₅H₉CH₂-Co-(DH)₂ were prepared by literature procedures.²⁹ The dimethyl sulfide ligands were removed by dissolving the adducts in N2-saturated water and evaporating to dryness under vacuum.

Solvents and Buffers. Buffer solutions (0.1 M) for UV-visible spectral studies were prepared by using CH₃COOH-CH₃COONa for pH 3-5 and K₂HPO₄-KH₂PO₄ for pH 6-8 in deionized distilled water that was previously boiled and stored under N₂. Buffer solutions (0.1 M) for ¹H NMR spectral studies were prepared with CD₃COOD-CD₃COONa for pD 3-5 and K₂HPO₄-KH₂PO₄ for pD 6-8 in 99.8% D₂O. Ethylene glycol was purified by a literature procedure.³⁰

Identification and Quantification of Thermolysis Products by ¹H NMR. Samples for ¹H NMR analysis were prepared in a flask equipped with a high-vacuum valve and a side arm. After the desired quantity of the solid compound was weighed and dissolved in N2-saturated D2O buffer solution, the valve was closed and the flask placed in a thermostated oil bath (±0.5 °C). The reaction mixture was cooled and sampled periodically under a N_2 atmosphere. Subsequently, the sample was oxidized by air, and a known concentration of internal reference $(TSP-d_4)$ was

(27) Hensens, O. D.; Hill, H. A. O.; McClelland, C. E.; Williams, R. J.

P. Ref 1a, Vol. I, p 463. (28) Summers, M. F.; Marzilli, L. G.; Bax, A. J. Am. Chem. Soc. 1986, 108. 4285.

(29) Schrauzer, G. N. Inorg. Synth. 1968, 11, 61.
(30) Perrin, D. D.; Arnarego, W. L. F.; Perrin, D. R. In Purification of Laboratory Chemicals; Pergamon: Oxford, 1966; p 163.

added. The thermolysis products were identified by comparison with the ¹H NMR spectra of authentic compounds and quantified by integration and comparison with the internal reference.

Kinetics Measurements. Samples for kinetic studies were prepared in a 1-cm Pyrex glass cell equipped with a high-vacuum valve and a side arm. The desired quantity of the solid compound was dissolved in the N₂-saturated aqueous buffer solution or ethylene glycol in the cell. The cell was closed and placed in the thermostated cell compartment of the UV-visible spectrophotometer or (with concordant results) in a thermostated oil bath. In the former case, the disappearance of the alkylcobalamins and appearance of B_{12} , were continuously monitored by measuring the absorbance at 650 nm. (No decomposition was observed when organocobalamin solutions were irradiated at this wavelength for 24 h at 25 °C.) In the latter case, the cell was removed from the oil bath periodically, cooled immediately, and placed in the thermostated cell compartment (25 °C), and the absorbance at 650 nm was recorded. These operations were performed in the dark.

The kinetic measurements were performed under pseudo-first-order conditions and the reactions followed for at least 3 half-lives. First-order rate constants were computed by linear least-squares regression.

Base On-Base Off Equilibrium Constants. Alkylcobalamin solutions $(1 \times 10^{-4} \text{ to } 2 \times 10^{-4} \text{ M})$ were prepared as for the kinetic measurements. Spectra were recorded over the pH range 2-7 and pK_a values, defined by eq 9, were determined by fitting spectral titration curves at ca. 522nm. Values of $\Delta H'$ and $\Delta S'$ listed in Table I were determined by linear least-squares regression fitting of the temperature dependence of the absorbance (A_{obsd}) at 522.5 nm (520 nm in the case of ethylene glycol) to the equation

$$\ln \left[(A_{\alpha} - A_{\text{obsd}}) / (A_{\text{obsd}} - A_0) \right] = -\Delta H' / RT + \Delta S' / R \quad (22)$$

The absorbance of the base-off form (A_0) was assumed to be that of

the limiting protonated form, i.e., [Co NH⁺], at low pH. The absorbance of the base-on form (A_{α}) was determined from the fitting procedure. The measured absorbances were corrected for thermal expansion of the solutions. Values of $(k_1)_{corr}$ were computed according to eq 11.

Acknowledgment. Support of this research through a grant from the National Institutes of Health (Grant DK 13339) is gratefully acknowledged. The NMR facilities were supported in part through The University of Chicago Cancer Center Grant No. N1H-CA-14599.

Registry No. B_{12_4} , 13422-51-0; Co(DH)₂, 36451-49-7; neopentyl-cobalamin, 71721-47-6; (cyclopentylmethyl)cobalamin, 113779-31-0; (cyclohexylmethyl)cobalamin, 113779-32-1; (tetrahydrofurfuryl)cobalamin, 75642-83-0; ((tetrahydro-2*H*-pyranyl)methyl)cobalamin, 113792-90-8; cyclopentylmethyl iodide, 27935-87-1; cyclohexylmethyl bromide, 2550-36-9; tetrahydrofurfuryl iodide, 5831-70-9; 2-(bromomethyl)tetrahydro-2H-pyran, 34723-82-5.

Reduction of Hydrogen Peroxide by the Ferrous Iron Chelate of Diethylenetriamine-N,N,N',N'',N''-pentaacetate

Siham Rahhal and Helen W. Richter*

Contribution from the Department of Chemistry, The University of Akron, Akron, Ohio 44325. Received April 27, 1987

Abstract: Ferrous iron complexed with diethylenetriamine-N,N',N'', pentaacetate (DTPA) reacts with hydrogen peroxide in neutral aqueous solution at room temperature to yield an oxidizing species that is not scavenged by tert-butyl alcohol, under conditions where >90% of hydroxyl radical would be scavenged. Further, the oxidizing species reacts with $Fe^{II}DTPA^{3-}$ and H_2O_2 at virtually the same rate, in contrast to the known behavior of the hydroxyl radical. These observations, the effects of methanol on the reaction, and the kinetic and stoichiometric results of the experiments clearly establish that the oxidizing species produced is not the hydroxyl radical, but an iron-oxo species such as the ferryl ion. The oxidizing species is formed in an apparent first-order reaction, when either hydrogen peroxide or Fe^{II}DTPA³⁻ is in kinetic excess. The bimolecular reaction rate constant is (1.37 \pm 0.07) \times 10³ M^{-1} s⁻¹.

The reduction of hydrogen peroxide by iron chelates is a well-known reaction.¹ In acid solution, it appears that the reduction produces the hydroxyl radical (HO[•]) (eq 1). This is the

 $Fe(II) + H_2O_2 = Fe(III) + HO^- + HO^-$ (1)

key reaction in the production of hydroxylated products by Fenton's reagent in strongly acid solutions. The rate in acid solution has been measured.²⁻⁸ Acid solution values have been

⁽¹⁾ Walling, C. Acc. Chem. Res. 1975, 8, 125-131.