

Note

THERMAL STUDIES ON COMPLEXES OF ZINC(II) AND CADMIUM(II) WITH 5,6-BENZOQUINOLINE-*N*-OXIDE

A.K. SRIVASTAVA, ANUPAMA SRIVASTAVA and P.C. JAIN

Meerut College, Meerut-250 001 (India)

(Received 8 January 1988)

In recent years, a number of complexes of 5,6-benzoquinoline [1–5] and aromatic amine-*N*-oxides [6–9] have been isolated and characterised. However, the thermal investigations of these complexes have been carried out only to a limited extent. In this paper we report the thermal decomposition characteristics, i.e. cleavage of metal–ligand bond and accompanying enthalpy change, as determined by thermogravimetry (TG), differential thermal gravimetry (DTG) and differential thermal analysis (DTA), of zinc(II) and cadmium(II) complexes of 5,6-benzoquinoline-*N*-oxide and their intermediates.

EXPERIMENTAL

5,6-Benzoquinoline was obtained from Merck and its oxide was prepared by the method described for quinoline-*N*-oxide [10]. The metal salts and other chemicals used were of BDH AnalaR grade.

The metal complexes were prepared by the following general method: an ethanolic solution of the corresponding metal salt (0.01 mole in 40 ml) was refluxed with a solution of the ligand (0.04 mole in 30 ml ethanol) on a waterbath for about 0.5 h. From the refluxate the complexes were separated either on standing or on concentrating and cooling to room temperature. The product was collected, washed with cold ethanol and finally with diethylether and dried in vacuo over P_4O_{10} .

The elemental analyses and physical measurements were made as reported earlier [11]. The differential thermal analysis and thermogravimetry units employed were similar to those described earlier [12]. All experiments were carried out in static air. Heating rates employed were $10^{\circ}\text{C min}^{-1}$ for DTA and $5^{\circ}\text{C min}^{-1}$ for TGA.

RESULTS AND DISCUSSION

The composition together with the analytical and partial IR data of Zn(II) and Cd(II) complexes of 5,6-benzoquinoline-*N*-oxide (BenzqnO) are

TABLE 1

Analytical, partial IR and thermal decomposition data of 5,6-benzoquinoline-*N*-oxide complexes of Zn(II) and Cd(II)


Compound	Metal (%)		N (%)		Anion (%)		$\nu(\text{NO})$	$\delta(\text{NO})$	Decomp. temp. (°C)		Decomp. product	Wt. loss (%)	
	Found	calcd.	Found	calcd.	Found	calcd.			Initial	Final		Found	Calcd.
BenzqnO	—	—	—	—	—	—	1240sh 1235s	840s	—	—	—	—	—
ZnCl ₂ ·4BenzqnO	7.31 7.09	6.23 6.11	7.51 7.75	—	1215m	—	835m	450m	210 310 530	290 390 630	ZnCl ₂ ·2BenzqnO ZnCl ₂ ZnO	44.12 86.95 92.39	42.57 85.15 91.15
ZnBr ₂ ·4BenzqnO	6.57 6.46	5.70 5.57	15.61 15.92	—	1220s	—	830m	445m	205 305 540	285 390 610	ZnBr ₂ ·2BenzqnO ZnBr ₂ ZnO	40.16 79.32 93.32	38.80 77.61 91.57
ZnI ₂ ·4BenzqnO	6.16 5.91 6.92 6.76	5.29 5.09 8.92 8.74	22.76 23.11 11.73 12.07	—	1225s	—	830m	440m	—	—	—	—	—
Zn(NCS) ₂ ·4BenzqnO	—	—	—	—	1225m	—	835m	435m	200 320 545	295 390 620	Zn(NCS) ₂ ·2BenzqnO Zn(NCS) ₂ ZnO	42.16 83.02 93.16	40.58 81.16 91.57
Zn(CH ₃ COO) ₂ ·4BenzqnO	6.91 6.74	5.96 5.81	— —	—	1225s	—	830m	440m	190 330 520	280 395 615	Zn(CH ₃ COO) ₂ ·2BenzqnO Zn(CH ₃ COO) ₂ ZnO	42.16 82.36 93.85	40.49 81.16 91.58
CdCl ₂ ·4BenzqnO	11.82 11.63	5.95 5.81	7.19 7.37	—	1220s	—	835s	445s	215 305 515	295 395 620	CdCl ₂ ·2BenzqnO CdCl ₂ CdO	42.10 83.16 88.92	40.49 80.99 86.70
CdBr ₂ ·4BenzqnO	10.80 10.64	5.51 5.32	14.82 15.20	—	1225m	—	830m	440m	210 360 510	300 405 625	CdBr ₂ ·2BenzqnO CdBr ₂ CdO	39.32 76.20 89.16	37.07 74.14 87.83
CdI ₂ ·4BenzqnO	9.91 9.77	5.02 4.88	21.56 22.16	—	1230m	—	835m	445m	—	—	—	—	—
Cd(NCS) ₂ ·4BenzqnO	11.29 11.11	8.56 8.33	11.32 11.50	—	1235m	—	840m	450m	200 320 520	290 390 620	Cd(NCS) ₂ ·2BenzqnO Cd(NCS) ₂ CdO	40.16 79.32 88.90	38.69 77.38 87.30

reported in Table 1. The IR assignments are largely based on the assignments in benzoquinoline and quinoline-*N*-oxide [13]. The decrease in the frequency of the $\nu(\text{NO})$ vibrations from 1235 cm^{-1} in the ligand to $1220 \pm 5\text{ cm}^{-1}$ in the complexes is attributed to a change in the nature of the nitrogen–oxygen bond as a result of metal–oxygen coordination [6–9]. The $\delta(\text{NO})$ bending vibration, observed at 840 cm^{-1} in BenzqnO, also shows a slight shift supporting the metal–oxygen coordination [6–9]. The bands observed in the $450\text{--}430\text{ cm}^{-1}$ region have been tentatively assigned to metal–oxygen stretching vibration. In the thiocyanate complexes, the $\nu(\text{C--S})$ vibrations lie in the $860\text{--}820\text{ cm}^{-1}$ region and $\nu(\text{C--N})$ in the $2080\text{--}2060\text{ cm}^{-1}$ region, strongly suggesting N-bonded NCS [14].

Thermal studies

The TG and DTG curves do not show the presence of water. TG analysis of the complexes shows that all the Zn(II) and Cd(II) complexes are stable up to about 200°C , beyond which decomposition starts and continues up to about 300°C . Analysis of the thermograms shows that two moles of BenzqnO are lost at this stage. It is therefore assumed that two molecules of the ligand are loosely bonded to the metal in the complexes. On further heating in the temperature range $305\text{--}405^\circ\text{C}$ the complexes $\text{MX}_2 \cdot 2\text{BenzqnO}$ ($\text{M} = \text{Zn, Cd}$) lose all the remaining ligand molecules. Finally, metal oxide is obtained as residual mass at about 600°C . In all cases, the second decomposition reaction is indicated by the endo peak and the final decomposition to metal oxide by the exo peak in the DTG curve.

Thus, the analysis of the thermograms indicates that the decomposition process is completed in the following three steps



($\text{M} = \text{Zn, Cd}$; $\text{X} = \text{Cl, Br, CH}_3\text{COO, NCS}$).

If the minimum TG decomposition temperature is taken as a rough criterion of thermal stability, then the order for the complexes reported herein is: $\text{Cl} > \text{Br} > \text{NCS} > \text{CH}_3\text{COO}$.

REFERENCES

- 1 S.S.A. Zaidi and K.S. Siddiqi, *Anorg. Allg. Chem.*, 379 (1970) 329.
- 2 S.S.A. Zaidi and K.S. Siddiqi, *J. Inorg. Nucl. Chem.*, 35 (1973) 655.
- 3 S.S.A. Zaidi, K.S. Siddiqi and N.S. Neelam, *Indian J. Chem.*, 14A (1976) 209.
- 4 S.S.A. Zaidi, T.A. Khan and N.S. Neelam, *Indian J. Chem.*, 19A (1980) 169.
- 5 R.K. Agarwal and S.K. Gupta, *Thermochim. Acta*, in press.
- 6 R.G. Garvey, J.A. Nelson and R.O. Ragsdale, *Coord. Chem. Rev.*, 3 (1968) 375.
- 7 M. Orchin and P.J. Schmidt, *Coord. Chem. Rev.*, 3 (1968) 345.
- 8 N.M. Karayannis, *Coord. Chem. Rev.*, 11 (1973) 93.

- 9 N.M. Karayannis, A.N. Speca, D.E. Chesani and L.L. Pytleneski, *Coord. Chem. Rev.*, 20 (1976) 37.
- 10 E. Ochiai, *J. Org. Chem.*, 18 (1953) 548.
- 11 A.K. Srivastava, R.K. Agarwal, Veena Kapoor and P.C. Jain, *J. Indian Chem. Soc.*, 60 (1983) 496.
- 12 R.K. Agarwal, A.K. Srivastava and T.N. Srivastava, *Transition Met. Chem.*, 5 (1980) 95.
- 13 V. Krishnan and C.C. Patel, *Can. J. Chem.*, 44 (1966) 972.
- 14 J.L. Burmeister, *Coord. Chem. Rev.*, 1 (1966) 205; 3 (1968) 225.