

Synthesis of Ketones from Aliphatic Nitriles and Phenylmagnesium Bromide

BY CHARLES R. HAUSER, WILBERT J. HUMPHLETT AND MARTIN J. WEISS

Shriner and Turner¹ have shown that, in the synthesis of acetophenone from acetonitrile and phenylmagnesium bromide, the best result is obtained with a 300% excess of the Grignard reagent. Accordingly these workers employed this large excess of the reagent in corresponding reactions with higher homologs of acetonitrile. Apparently on this basis the statement has been made in texts² that, with the higher aliphatic nitriles, as well as acetonitrile, the large excess of the Grignard reagent is necessary for satisfactory yields of ketones. However, Bary³ had reported previously that a 90% yield of butyrophenone is obtained from butyronitrile with a 50% excess of phenylmagnesium bromide.

In the present investigation, high yields of ketones have been obtained from propionitrile and higher homologs through *n*-capronitrile with only a 10% excess of phenylmagnesium bromide. These results, and also the corresponding cases reported by Shriner and Turner,¹ are given in Table I. With the three higher nitriles, our results appear even slightly better than those from the large excess of the Grignard reagent.

It should be pointed out that, with ethylmagnesium bromide and propionitrile, we have obtained only a 23% yield of diethyl ketone (b. p. 101–105°) using a 10% excess of the reagent. Earlier workers⁴ have similarly reported only fair yields of ketones with this aliphatic Grignard reagent and propionitrile or butyronitrile.

TABLE I

YIELDS OF KETONES FROM ALIPHATIC NITRILES AND PHENYLMAGNESIUM BROMIDE

Nitrile	Ketones, 10% excess Grignard				Ketones, 300% excess Grig. ^a			
	B. p. °C.	Mm.	Yield, %	<i>n</i> _D ²⁰	B. p. °C.	Mm.	Yield, %	<i>n</i> _D ²⁰
Aceto	201–205		33	1.5344 ^a	202–205		70	
Propio	105–106	17	83	1.5270 ^b	115–120	21	91	
<i>n</i> -Butyro	121–123	20	82	1.5203 ^c	125–130	21	77	
<i>n</i> -Valero	139–141	24	83	1.5146 ^d	135–140	25	79	
<i>n</i> -Capro	137–138	13	89	1.5116	145–150	19	83	

^a "International Critical Tables," Vol. VII (compiled by C. J. West, National Research Council, McGraw-Hill Book Co., New York, N. Y., 1930, p. 43) gives 1.53427 at 19.1°. ^b Wallach, *Ann.*, **332**, 317 (1904). ^c "Handbook of Chemistry and Physics" (Chemical Rubber Publishing Co., 28th edition, 1944, p. 680) gives 1.52016 at 18.25°. ^d Layraud [*Bull. soc. chim.*, (3) **35**, 223 (1906)] gives 1.5152 at 19°. ^e Ref. 1.

Procedure.—In a 1-liter three-necked round-bottomed flask equipped through ground-glass joints with a mercury-sealed stirrer, dropping funnel, and a reflux condenser

(having a drying tube) was placed 300 ml. of an ether solution containing 0.275 mole of phenylmagnesium bromide. To the stirred refluxing solution was added, during fifteen to twenty minutes, 0.250 mole of the nitrile in 125 ml. of dry ether. The stirring and refluxing was continued for one to six hours⁵; during the first hour a precipitate formed. The cooled mixture was decomposed with ice and acid and the ether removed on the steam-bath under an ether still. After heating for an hour longer to ensure hydrolysis of the ketimine, the ketone was extracted with four 150-ml. portions of ether. The solvent was distilled from the dried combined ether solutions and the residue distilled through a 15 cm. Vigreux column.

(5) The yields given in Table I were obtained on six-hour runs but an equally good yield (85%) has been obtained in a one-hour run with butyronitrile.

DEPARTMENT OF CHEMISTRY
DUKE UNIVERSITY
DURHAM, N. C.

RECEIVED AUGUST 18, 1947

Some Azeotropes of Alkylacetylenes and Ethyl Alcohol¹

BY G. F. HENNION AND J. J. GROEBNER

Azeotropic distillation is a well-known means for the separation and purification of various hydrocarbons and data are available for many hydrocarbon-carrier systems. Particular attention has been given to the azeotropes of paraffins, naphthenes, olefins, diolefins and aromatics. Since there is no information on constant boiling mixtures of higher acetylenes, we have investigated four acetylene-ethyl alcohol binaries. The data are given in Table I. When the boiling points of the azeotropes are plotted against their compositions a smooth curve is obtained, similar to, but appreciably above, the one for paraffins.²

The lowering of the hydrocarbon boiling point achieved by azeotropic distillation is greatest for paraffins and least for aromatics; the other hydrocarbons are intermediate and usually in the order cited above.³ When the data for the acetylenes are compared with those of the other hydrocarbons it appears that the acetylenes fall between the aromatics and the others. Thus the

TABLE I

BINARY AZEOTROPES OF ALKYLACETYLENES AND ETHYL ALCOHOL

Acetylene ^a	Properties of azeotropes				
	B. P., °C.	<i>n</i> _D ²⁰	<i>d</i> ₄ ²⁰	Mole %	Wt. %
<i>n</i> -Butyl	62.8	1.3880	0.7264	65.0	76.8
Diethyl	67.5	1.3922	.7399	51.7	65.6
Isoamyl	71.0	1.3857	.7490	42.0	60.2
<i>n</i> -Amyl	74.2	1.3818	.7586	28.5	45.4

^a The boiling points of the pure hydrocarbons were: *n*-butylacetylene, 70.2°; diethyl-, 80.5°; isoamyl-, 90.8°; *n*-amyl-, 99.5°. Other physical properties are cited by Hennion and Banigan, *THIS JOURNAL*, **68**, 1381 (1946).

(1) Paper LI on the chemistry of substituted acetylenes; previous paper, *THIS JOURNAL*, **68**, 1381 (1946).

(2) Mair, Glasgow and Rossini, *J. Research Natl. Bur. Standards*, **27**, 47 (1941).

(3) Rossini, Mair and Glasgow, *Oil Gas J.*, **188**, Nov. 14 (1940).

(1) Shriner and Turner, *THIS JOURNAL*, **52**, 1267 (1930).

(2) Hickinbottom, "Reactions of Organic Compounds," Longmans, Green and Co., New York, N. Y., 1936, p. 258; Fuson and Snyder, "Organic Chemistry," John Wiley and Sons, Inc., New York, N. Y., 1942, p. 270.

(3) Bary, *Bull. soc. chim. Belg.*, **31**, 397 (1922).

(4) See ref. 3 and Baerts, *ibid.*, **31**, 184 (1922).