

**140. The Donor-Acceptor-Acceptor Purine Analog:
Transformation of 5-Aza-7-deaza-1*H*-isoguanine (= 4-Aminoimidazo-
[1,2-*a*]-1,3,5-triazin-2(1*H*)-one) to 2'-Deoxy-5-aza-7-deaza-isoguanosine
Using Purine Nucleoside Phosphorylase**

by Johannes J. Voegel, Michael M. Altorfer, and Steven A. Benner*

Institut für Organische Chemie, ETH Zürich, Universitätstrasse 16, CH-8092 Zürich

(20.V.93)

A new synthesis is reported for 4-aminoimidazo[1,2-*a*]-1,3,5-triazin-2(1*H*)-one (= 5-aza-7-deaza-isoguanosine; **8**), a purine analog that, when incorporated into an oligonucleotide chain, presents a H-bond donor-acceptor-acceptor pattern to a complementary pyrimidine analog. A protected ribose derivative was coupled to **8** to yield 4-amino-8-(β -D-ribofuranosyl)imidazo[1,2-*a*]-1,3,5-triazin-2(8*H*)-one (= 5-aza-7-deaza-isoguanosine; **11**) after deprotection. Alternatively, direct synthesis of both the ribo derivative **11** and the corresponding deoxyribo derivative **17** as the β -D-anomers was achieved using the enzyme purine nucleoside phosphorylase in a one-pot reaction. This adapts a known synthetic approach to yield a new strategy for obtaining diastereoisomerically pure deoxyribonucleoside analogs on 1-gram scales.

Introduction. – In the preparation of ribonucleosides and their analogs, anchimeric assistance from substituents at the 2'-position of the D-ribose ring readily yields products having a β -D-configuration at the anomeric center [2–5]. In deoxyribonucleosides, similar assistance is not available, and reactions that couple a heterocycle and a 2'-deoxy-D-ribose derivative generally yield mixtures of α - and β -D-isomers [6–13]. This is the key problem in the synthesis of many deoxyribosides, and much work has been devoted to solving it.

Enzyme-catalyzed coupling of deoxyribose derivatives and heterocycles offers one possible strategy for obtaining anomerically pure nucleoside analogs. *E.g.*, both *N*-deoxyribosyltransferase and purine nucleoside phosphorylase (PNPase) transfer heterocycles on and off deoxy-D-ribose derivatives, with the β -D-anomer being the exclusive product [14] [15]. The use of enzymes as synthetic tools often encounters other difficulties, however, including inconvenient reactants and products, narrow substrate specificity, and idiosyncrasies and unreliability of reaction procedures.

Hennen and *Wong* [1] recently reported a particularly elegant procedure for addressing the first difficulty in ribose transferase reactions. In their procedure, 7-methylguanosine (**12**) serves as the glycosyl donor (see below, *Scheme 3*). Ribose 1-phosphate is formed *in situ* with the concomitant precipitation of essentially insoluble 7-methylguanine (**14**) as by-product. The ribose 1-phosphate can then accept a wide range of heterocycles (see below, *Scheme 3*).

The narrow substrate specificity of many glycotransferases remains problematic. For example, *N*-deoxyribosyltransferase does not accept substrates that differ only slightly in structure from the natural substrates (*e.g.* 7-deazapurine) [16]. Thus, this enzyme cannot

serve as a general tool for preparing nucleoside analogs. This problem is alleviated using PNPase, which can accept a wider range of heterocycles [17–21]. Further, 2'-deoxyribonucleosides are substrates for PNPase [17] [20] [22] [23]. However, the *Hennen-Wong* procedure with 2'-deoxy-7-methylguanosine (**18**) as glycosyl donor has, to our knowledge, never been explored with deoxyribose derivatives and PNPase.


As part of work in this laboratory aimed at expanding the genetic alphabet [24], we had need for both 9-(ribosyl) [25] and 9-(2'-deoxyribosyl)-5-aza-7-deaza-9*H*-isoguanine (**11** and **17**, resp., see below *Scheme 3*). The heterocycle borne by these nucleoside analogs presents a H-bond donor-donor-acceptor pattern to a pyrimidine analog on a complementary strand of DNA.

Rosemeyer and *Seela* prepared a small amount of a different (but related) β -D-nucleoside, 2'-deoxy-5-aza-7-deazaguanosine, by incubating 2'-deoxy- α -D-ribofuranose 1-phosphate, the heterocycle, and PNPase [11]. Therefore, we thought it should be relatively straightforward to use the procedure of *Hennen* and *Wong* to prepare 5-aza-7-deazaisoguanosine (**11**) and 2'-deoxy-5-aza-7-deazaisoguanosine (**17**). Unfortunately, our work encountered the third problem commonly observed in enzyme-assisted synthetic reactions: idiosyncrasies. When 7-methylguanosine (**12**) [26] was incubated with PNPase under the conditions of *Hennen* and *Wong* (0.25M phosphate buffer, pH 7.8 room temperature) [1] in the presence of 5-aza-7-deaza-1*H*-isoguanine, essentially none of the desired product was obtained.

Notably, the enzyme used by *Hennen* and *Wong* was obtained from *Toyoba Chemical Co.*, ours was from *Sigma*. While we do not know whether this difference fully accounts for the failure of the reaction with 5-aza-7-deaza-1*H*-isoguanine and for the extremely low conversion rates with adenine in our hands using *Hennen-Wong* conditions (see below), the *Sigma* enzyme is more accessible to many workers. Therefore, an optimal set of reaction conditions was developed using the *Sigma* enzyme with adenine as a substrate. Then, these conditions were applied to the preparation of 5-aza-7-deazaisoguanosine (**11**).


Results and Discussion. – For comparison purposes, 5-aza-7-deazaisoguanosine (**11**) was first prepared by a classical approach starting from aminoimidazolium sulfate **1**. The free base **2** of **1** was obtained by neutralization of an aqueous solution of **1** with Na_2CO_3 , followed by extraction with EtOH or MeCN [27]. In analogy to syntheses performed with 2-aminothiazole [28] [29], the condensed triazine ring system **5** was synthesized by reaction of 2-amino-1*H*-imidazole (**2**) with (phenylthio)carbonyl isothiocyanate (**4**), which was prepared *in situ* from *S*-phenyl chlorothioformate (**3**; *Scheme 1*). The polarity of the

Scheme 1



product made it difficult to isolate. Therefore, crude **5** was treated directly with benzyl bromide, yielding dibenzylated product **6**, which was easily separated from other alkylated products by chromatography. The benzyl-sulfide group was then replaced with ammonia in MeOH, yielding the benzylated nucleoside base **7**. Conditions could not be found for removal of the benzyl group of **7** by catalytic hydrogenation using Pd catalysis [30–33]. But reaction of **7** with Na in liquid ammonia [34] yielded the desired 5-aza-7-deaza-1*H*-isoguanine (**8**; Scheme 2).

Scheme 2




The glycosylation of 5-aza-7-deaza-1*H*-isoguanine (**8**) failed, remarkably, under the conditions of the silyl modification of the *Hilbert-Johnson* reaction using trimethylsilyl triflate as catalyst developed by *Vorbrüggen* and coworkers [3–5]. Therefore, the classical method (silylated base, SnCl₄/dichloroethane) [35], already applied successfully to the synthesis of 5-aza-7-deazaguanosine by *Kim et al.* [36], was used. This procedure applied to **8** and **9** led to the protected ribonucleoside **10** in a yield of 60%. Cleavage of the benzoyl protecting groups with NH₃ in MeOH yielded the target 5-aza-7-deazaisoguanosine (**11**) as a crystalline white solid (Scheme 2). The analytical data agreed with those reported by *Prisbe et al.* [25], and this material served as a standard in the enzymatic work.

A systematic study was then made to find optimal conditions for enzymatic coupling of adenine to ribose, the latter being generated as the 1-phosphate derivative from 7-methylguanosine (**12**) according to *Hennen* and *Wong* [1] (Scheme 3). The high concentrations of phosphate (250 mM) recommended by *Hennen* and *Wong* reduced the yield of the expected adenosine (**13**) by ca. 80% with the *Sigma* enzyme employed by us. Of three buffer concentrations tested (250, 100, and 50 mM), best results were achieved with 50 mM. The yield was insensitive to the pH between 7.0 and 7.8. The enzyme tolerated temperatures of 60°, and the yield of product **13** was highest at this temperature (the by-product 7-methylguanine **14** precipitated).

The optimal reaction conditions (phosphate buffer 0.05M, pH 7.0, 60°) were then used to prepare 5-aza-7-deazaisoguanosine (**11**) from **12** and 5-aza-7-deaza-1*H*-isoguanine (**8**) according to Scheme 1. HPLC analysis showed more than 50% product formation after

Scheme 3



10 days (Fig. 1), establishing that **11** can be prepared enzymatically. As the chemical synthesis appears preferable for the preparation of large amounts of **11** the workup was not optimized. The structure of the product was proven by comparison with material synthesized *via* the nonenzymatic route. A comparison between adenine and **8** showed that **8** reacted to **11** *ca.* 5 times slower than adenine to **13**.

This procedure was then directed towards the synthesis of the corresponding deoxyribonucleoside. The 2'-deoxy-7-methylguanosinium iodide (**15**) was prepared by alkylation of 2'-deoxyguanosine [26] and used as the glycosyl donor (Scheme 3). Reaction conditions were optimized using **15** and adenine for the formation of 2'-deoxyadenosine




Fig. 1. Enzymatic formation of 5-aza-7-deazaisoguanosine (**11**) by Route A (see Scheme 3). Reaction conditions: 5-aza-7-deaza-1H-isoguanine (**8**), 8 μ mol; 7-methylguanosine (**12**), 32 μ mol; PNPase, 4 u; phosphate conc., 0.05M; volume, 1.66 ml; pH 7.0; temp. 60°. Yields by HPLC.




Fig. 2. Enzymatic formation of 2'-deoxyadenosine (**16**) by Route B (see Scheme 3). Reaction conditions: adenine, 8 μ mol; 2'-deoxy-7-methylguanosinium iodide (**15**), 32 μ mol; PNPase, 9 u; phosphate conc., 0.05M; volume, 1.66 ml; pH 7.4; time 16 h. Yields by HPLC.

(**16**) (see Fig. 2). The optimal temperature was found to be 45°; at higher temperature, product formation decreased rapidly. Moreover, rather high enzyme concentrations (23 units/8 μ mol of heterocycle) were necessary to achieve high conversion rates (70 % of **16**) in a reasonable time (16 h).

The optimal reaction conditions (45°, 23 units PNPase, 50 mM phosphate, 32 μ mol of **15**, 8 μ mol free base, 1.66 ml) were then used for the synthesis of 2'-deoxy-5-aza-7-deazaisoguanosine (**17**) from **15** and **8** (Scheme 3). A comparison between **8** and adenine showed that 2'-deoxy-5-aza-7-deazaisoguanosine (**17**) was formed *ca.* three times slower than 2'-deoxyadenosine (**16**). With 2'-deoxy-7-methylguanosinium iodide (**15**), yields of **17** (based on heterocycle) were never higher than 20%, even with very high enzyme concentrations. Variation of pH and temperature did not increase the yields. Interestingly, however, the glycosyl donor influenced greatly the yield of the reaction. Yields after 24 h using **15** were *ca.* 50% below those obtained under analogous conditions with the zwitterionic 2'-deoxy-7-methylguanosine (**18**; Scheme 3), prepared from **15** by neutralization with MeNH₂/H₂O [26]. Moreover, with iodide salt **15**, the pH of the reaction mixture changed upon cleavage of 7-methylguanine, making high concentrations of this glycosyl donor incompatible with low concentrations of buffer.

By varying the concentration of 2'-deoxy-7-methylguanosine (**18**) and PNPase, conditions useful for the synthesis of 2'-deoxy-5-aza-7-deazaisoguanosine (**17**) could be found. Doubling the glycosyl-donor concentration led to an increase in the yield of **17** of 40%. Further increase in the concentration of **18** increased yields only slightly. The amount of **17** isolated was a linear function of the amount of enzyme added, and high enzyme concentrations were used for an efficient synthesis of **17**. After equilibrium was attained, the enzyme was recovered by ultrafiltration and recycled. Typically, *ca.* 5% of enzyme activity was lost during each recycling step. The starting concentration of PNPase was chosen so that at least five cycles could be run. By this technique, 1120 units of PNPase and 780 μ mol of 5-aza-7-deaza-1*H*-isoguanine (**8**) yielded 54% of isolated deoxyribonucleoside **17**.

In the deoxy series, a NOE from H–C(1') to H–C(4') established the β -D-configuration of the principal product (a NOE from H–C(1') to H–C(3') expected in the α -D-isomer was missing) [37]. The position of glycosylation (N(8)) of the imidazotriazine base was assigned by comparison of the ¹H- and ¹³C-NMR data of **17** with the corresponding ribonucleoside **11** and the N(1)-linked ribonucleoside, both known by independent chemical synthesis [25] [38].

This work showed PNPase to be a servicable tool for preparing β -D-configured purine nucleoside analogs on a 1-gram scale. Further, by using deoxymethylguanosine **18** as a glycosyl donor, a glycosyl exchange reaction proved to be satisfactory, avoiding the need for a second enzyme to generate the intermediate ribose 1-phosphate.

The authors are indebted to Brigitte Brandenburg, Marco Sperl, and PD Dr. Bernhard Jaun for NMR measurements, to Rolf Häfliger, Oswald Greter, and Dr. Walter Amrein for obtaining MS data, to Michael Schneider and Dieter Manser for performing elemental analysis, to Hans-Ulrich Hediger for IR spectroscopy, to Dr. Engelbert Zass for CAS on-line searches, and to Anne Preisig for administrative assistance. J. J. V. thanks Dr. Rosenkranz for a postgraduate scholarship. Financial support from the Swiss National Science Foundation and Sandoz AG is gratefully acknowledged.

Experimental Part

General. *Fluka* solvents (*puriss. p.a.*, *puriss.*, or *purum*) were used for all syntheses. MeCN was distilled over CaH₂ and DMF dried over molecular sieves (4 Å, 2–3 mm). The 7-methylguanosine (**12**) was synthesized according to [26]. The following chemicals were purchased from *Fluka*: adenine (*purum*), 1-*O*-acetyl-2',3',5'-tri-*O*-benzoyl-β-D-ribofuranose (*purum*), 2'-deoxyguanosine monohydrate (> 98%), benzyl bromide (*purum*), 1,1,1,3,3,3-hexamethyldisilazane (HMDS; *purum*, freshly distilled before use), MeNH₂ (*purum*), Na₂CO₃, and Et₃N. From *Aldrich* were purchased: 2-aminoimidazolesulfate and SnCl₄. KSCN was obtained from *Mallinckrodt Chemical Works*, (phenylthio)carbonyl chloride (= *S*-phenyl chlorothioformate) from *Alfa Products*, and KH₂PO₄ from *Merck*. From *Sigma* were purchased: guanosine (*Sigma Grade*) and bacterial purine nucleoside phosphorylase (= PNPase; lyophilized powder (60% enzyme), 14 u/mg; 1 unit is the amount of enzyme that phosphorylates 1.0 μmol of inosine to hypoxanthine and ribose 1-phosphate at 25° and pH 7.4). PNPase was recycled *via* ultrafiltration through a *Centricon-10* filtration apparatus purchased from *Amicon*, *Diaflo* ultrafiltration membranes of the type *PM 10* were used. Column chromatography: reversed-phase silica gel (*C18*, 40 μm) from *J. T. Baker*. Flash chromatography (FC): standard flash silica gel *60*. Ion-exchange chromatography: *Dowex 1 × 8 resin*, 200–400 mesh, OH form (*Bio-Rad*) 1.5 × 16 cm column; 10–15 ml fractions; elution with linear gradient 0–1.0M (Et₃NH)HCO₃, total volume 400 ml; detection by UV at 254 nm, single fractions by HPLC or HPTLC. HPLC: all enzymatic reactions with PNPase were followed by measuring the UV absorbance at 254 and 277 nm in the HPLC; the integration of the UV absorbance at 254 nm formed the basis for calculating the yields of reactions that were not worked up; for deoxyadenosine, and anal. *C8* reversed-phase column (*Brownlee Labs Aquapore RP-300*, 22 cm × 4.6 mm, particle size 7 μm) was used, flow rate 1 ml/min, elution buffer 0.1M (Et₃NH)OAc, pH 7.5, without gradient, for all other reactions, a semi-prep. *C18* reversed-phase column (*Supelco LC-18-DB*, 25 cm × 10 mm, particle size 5 μm) was used, flow rate 2 ml/min, elution buffer 0.1M (Et₃NH)Ac, pH 7.5, with gradient *A* (0–4% MeCN in 5 min, then increase to 8% MeCN in another 25 min) or gradient *B* (0–4% MeCN in 30 min). UV spectra: λ_{max}(ε) in nm. IR Spectra: in cm⁻¹. NMR Spectra: at 93.94 kg (400 MHz for ¹H, 100 MHz for ¹³C) or 70.46 kg (300 MHz for ¹H, 75 MHz for ¹³C), δ in ppm rel. to Me₄Si (= 0.00 ppm) as internal standard; OH and NH assignments were confirmed by D₂O exchange. MS: in *m/z* (rel. intensity in %).

Nonenzymatic Syntheses. – 2-Amino-1H-imidazole (**2**). Following literature procedures [27], **1** (3.30 g, 25 mmol) was dissolved in H₂O (15 ml) together with Na₂CO₃ (2.65 g, 25 mmol). The H₂O was removed by evaporation and the residue extracted with EtOH (6 × 20 ml). The solvent was evaporated and the remaining brown oil dried overnight under high vacuum: **2** (1.65 g, 79%), which was used in the next reaction.

A similar reaction on larger scale of **1** (13.77 g, 104 mmol) gave **2** (6.51 g, 75%).

3,4-Dihydro-4-thioxoimidazo[1,2-*a*]-1,3,5-triazin-2(1H)-one (**5**). KSCN (8.00 g, 75 mmol) was suspended in abs. MeCN (70 ml). *S*-Phenyl chlorothioformate (**3**; 9.40 ml, 75 mmol) was added dropwise at r.t. The product **4**, formed *in situ* after 5 min stirring, was immediately used in the subsequent cyclization. A larger-scale reaction was also carried out with a larger amount of **3** (21 ml, 168 mmol) and KSCN (18.65 g, 192 mmol).

To the suspension of **4** in MeCN was added Na₂CO₃ (2.65 g, 25 mmol) and then dropwise a soln. of **2** (1.65 g, 20 mmol) in MeCN (15 ml). The mixture was stirred for 1 h. The yellow precipitate was filtered off, washed with MeCN and Et₂O, and dried under high vacuum: crude **5** (10.03 g). Yellow brown solid, which was used directly for the next reaction.

A larger-scale reaction with more **2** (6.50 g, 78 mmol) gave **5** (22.05 g).

1-Benzyl-4-(benzylthio)imidazo[1,2-*a*]-1,3,5-triazin-2(1H)-one (**6**). In a soln. of **5** (10.03 g) in DMF (70 ml), Na₂CO₃ (2.65 g, 25 mmol) was suspended and benzyl bromide (5.95 ml, 50 mmol) injected. The soln. was stirred 1.5 h at r.t., then diluted with CH₂Cl₂, and extracted with sat. aq. NH₄Cl soln., H₂O, and sat. aq. NaCl soln. The org. phase was dried (MgSO₄) and evaporated. Chromatography (silica gel (300 g), Et₂O/hexane 8:2) yielded **6** (1.06 g, 15% over 2 steps). Light yellow solid, which was used directly for the next reaction. Ring-benzylated products, not useful for subsequent synthesis, were the primary side products. An anal. sample of **6** was obtained *via* recrystallization from Et₂O/hexane 3:1. M.p. 134–135°. UV (EtOH): 234 (11 300), 260 (10 700), 300 (2600). IR (CHCl₃): 3060, 3025, 1680, 1610, 1525, 1490, 1450, 1370, 1350, 1300, 1170, 1040, 960. ¹H-NMR (CDCl₃): 4.59 (*s*, CH₂S); 5.34 (*s*, CH₂N); 7.04 (*d*, *J* = 1.9, H-C(6)); 7.07 (*d*, *J* = 1.9, H-C(7)); 7.24–7.43 (*m*, 4 H_{ar}, 4 H_m); 7.57–7.61 (*m*, 2 H_p). ¹³C-NMR (CDCl₃): 35.74 (*t*, CH₂S); 46.94 (*t*, CH₂N); 109.35 (*d*, C(6)); 128.10 (*d*, arom. CH); 128.29 (*d*, arom. CH); 128.53 (*d*, arom. CH); 128.97 (*d*, arom. CH); 129.34 (*d*, arom. CH); 129.42 (*d*, arom. CH); 131.07 (*d*, C(7)); 134.50 (*s*, C_{ipso}); 135.60 (*s*, C_{ipso}); 144.47 (*s*, C(8a)); 151.01 (*s*, C(2)); 160.40 (*s*, C(4)). MS: 349 (6, [M+]⁺), 348 (24, M⁺), 257 (8), 200 (12), 91 (100). Anal. calc. for C₁₉H₁₆N₄OS (348.43): C 65.50, H 4.63, N 16.08; found: C 65.26, H 4.70, N 16.07.

A scaled-up reaction with **5** (22.0 g) gave **6** (3.32 g, 12% over 2 steps).

4-Amino-1-benzylimidazof[1,2-a]-1,3,5-triazin-2(1H)-one (7). To **6** (835 mg, 2.4 mmol) in a dry flask, MeOH sat. with NH_3 (40 ml) was injected at low temp. The flask was sealed and the mixture stirred for 1 h at r.t. The NH_3 was evaporated and the solvent removed under vacuum to yield **7** (0.594 g). Slightly impure white solid, which was used without further purification for the next reaction. An anal. sample of **7** was obtained *via* recrystallization from MeCN. M.p. 238° (dec.). UV (EtOH): 232 (9100), 263 (3700). IR (KBr): 3340, 3150, 3125, 3025, 2740, 1680, 1635, 1540, 1490, 1370, 1230, 1080, 1050. $^1\text{H-NMR}$ ((D_6) DMSO): 5.13 (s, CH_2); 7.04 (d, $J = 1.8$, H-C(6)); 7.21–7.37 (m, 5 arom. H); 7.63 (d, $J = 1.8$, H-C(7)); 8.19–8.49 (br. s, NH_2). $^{13}\text{C-NMR}$ ((D_6) DMSO): 44.34 (t, CH_2); 108.53 (d, C(6)); 126.03 (d, C_m); 126.58 (d, C_o); 127.10 (d, C_p); 128.18 (d, C(7)); 136.08 (s, C_{ipso}); 144.25 (s, C(8a)); 148.52 (s, C(2)); 152.07 (s, C(4)). MS: 242 (s, $[M + 1]^+$), 241 (31, M^+), 199 (16), 91 (100). Anal. calc. for $\text{C}_{12}\text{H}_{11}\text{N}_5\text{O}$ (241.25): C 59.74, H 4.60, N 29.03; found: C 59.72, H 4.85, N 29.14.

An analogous reaction with more **6** (3.30 g, 9.5 mmol) gave **7** (2.48 g).

4-Aminoimidazof[1,2-a]-1,3,5-triazin-2(1H)-one (= 5-Aza-7-deaza-1H-isoguanine; 8). To a suspension of **7** (372 mg, 1.5 mmol) in liquid NH_3 (30 ml), Na was added in small pieces until the blue color persisted. Excess Na was destroyed with NH_4Cl and the NH_3 evaporated. The residual solid was dissolved in H_2O (7 ml) and the pH of the soln. adjusted to 5 with 2M HCl. The soln. was cooled and the resulting precipitate isolated by filtration: **8** (0.203 g, 85% over 2 steps). White solid. An anal. sample was obtained by recrystallization from MeOH/ H_2O 8:1. M.p. > 300°. UV (EtOH): 239 (6500), 255 (2500). IR (KBr): 3250, 3130, 2760, 1710, 1660, 1570, 1470, 1300, 1205, 1080. $^1\text{H-NMR}$ ((D_6) DMSO): 6.97 (d, $J = 1.8$, H-C(6)); 7.55 (d, $J = 1.8$, H-C(7)); 8.08 (br. s, NH_2). $^{13}\text{C-NMR}$ ((D_6) DMSO): 108.06 (d, C(6)); 128.88 (d, C(7)); 144.83 (s, C(8a)); 150.53 (s, C(2)); 154.24 (s, C(4)). FAB-MS (thioglycerine): 152 ($[M + 1]^+$). Anal. calc. for $\text{C}_5\text{H}_5\text{N}_5\text{O} \cdot \frac{1}{2} \text{H}_2\text{O}$ (160.14): C 37.49, H 3.78, N 43.74; found: C 37.58, H 3.54, N 43.60.

An analogous reaction of **7** (1.30 g, 5.4 mmol) gave **8** (668 mg, 87% over 2 steps).

4-Amino-8-(2',3',5'-tri-O-benzoyl- β -D-ribofuranosyl)imidazof[1,2-a]-1,3,5-triazin-2(8H)-one (10). Dry **8** (23.7 mg, 0.157 mmol) was suspended in freshly distilled HMDS (0.5 ml). A small amount of NH_4Cl was added and the mixture refluxed for 24 h. The excess HMDS was removed by distillation under high vacuum. The resulting white solid was dried and then dissolved in dichloroethane (0.4 ml). A soln. of **9** (79.2 mg, 0.157 mmol) in dichloroethane (0.4 ml) was added, followed by SnCl_4 (25 μl). The mixture was stirred for 4 h at r.t., then diluted with CH_2Cl_2 and extracted with sat. aq. NaHCO_3 and NaCl solns. The org. phase was dried (Na_2SO_4) and the solvent evaporated. The isolated oil was purified by chromatography (small silica-gel column, $\text{CH}_2\text{Cl}_2/\text{EtOH}$ 9:1): **10** (56 mg, 60%) as light brown solid. UV (EtOH): 218 (42400), 230 (46400). IR (KBr): 3420, 3130, 3060, 1730, 1690, 1600, 1540, 1450, 1315, 1270, 1180, 1090, 1025. $^1\text{H-NMR}$ ((D_6) DMSO): 4.64–4.83 (m, H-C(4'), 2 H-C(5')); 6.03 (dd, $J = 5.5, 5.9$, H-C(3')); 6.19 (dd, $J = 5.1, 5.9$, H-C(2')); 6.30 (d, $J = 5.1$, H-C(1')); 7.42–7.70, 7.88–8.04 (2m, 15 arom. H, H-C(6), H-C(7)); 7.79 (br. s, NH_2). $^{13}\text{C-NMR}$ ((D_6) DMSO): 63.62 (t, C(5')); 70.56 (d, C(3')); 72.55 (d, C(2')); 78.99 (d, C(4')); 85.39 (d, C(1')); 107.42 (d, C(6)); 115.73 (d, C(7)); 128.19 (s, C_{ipso}); 128.45 (s, C_{ipso}); 128.68 (d, C_o , C_m); 129.13 (s, C_{ipso}); 129.26 (d, C_o , C_m); 133.44 (d, C_p); 133.79 (d, C_p); 133.92 (d, C_p); 149.36 (s, C(8a)); 150.49 (s, C(2)); 161.53 (s, C(4)); 164.43 (s, Bz); 164.51 (s, Bz); 165.41 (s, Bz). FAB-MS (3-nitrobenzyl alcohol): 618 ($[M + \text{Na}]^+$), 596 ($[M + 1]^+$), 445 (ribose part), 152 (free base).

An analogous reaction with **8** (50 mg, 0.33 mmol) gave **10** (82 mg, 44%) as light brown foam.

4-Amino-8-(β -D-ribofuranosyl)imidazof[1,2-a]-1,3,5-triazin-2(8H)-one (= 5-Aza-7-deazaisoguanosine; 11). Compound **10** (51 mg) was suspended in EtOH saturated with NH_3 (0.5 ml). The flask was sealed and the mixture stirred overnight at r.t. After evaporation of excess of NH_3 , the soln. was cooled to 0° and the flocculant solid recovered by filtration and dried under high vacuum: **11** (17 mg, 70%). White solid. M.p. 240° (dec.). UV (EtOH): 245 (12200). IR (KBr): 3430, 3320, 3280, 3200, 3080, 2850, 1675, 1635, 1600, 1560, 1450, 1360, 1255, 1130, 1080, 1040. $^1\text{H-NMR}$ ((D_6) DMSO): 3.51–3.64 (m, 2 H-C(5')); 3.89–3.91 (m, H-C(4')); 4.05–4.06 (m, H-C(3')); 4.28–4.31 (m, H-C(2')); 5.18 (br. s, 2 OH); 5.48 (br. s, OH); 5.69 (d, $J = 5.6$, H-C(1')); 7.49 (d, $J = 2.5$, H-C(6)); 7.53 (d, $J = 2.5$, H-C(7)); 7.82 (br. s, NH_2). $^1\text{H-NMR}$ ((D_6) DMSO, D_2O exchange): 3.53 (dd, $J = 3.7, 12.0$, H_a -C(5')); 3.62 (dd, $J = 3.5, 12.0$, H_b -C(5')); 3.91 (ddd, $J = 3.5, 3.5, 3.7$, H-C(4')); 4.06 (dd, $J = 3.5, 5.0$, H-C(3')); 4.29 (dd, $J = 5.0, 5.7$, H-C(2')); 5.71 (d, $J = 5.7$, H-C(1')); 7.49 (d, $J = 2.9$, H-C(6)); 7.54 (d, $J = 2.9$, H-C(7)). $^1\text{H-NMR}$ (D_2O): 3.77 (dd, $J = 4.0, 12.7$, H_a -C(5')); 3.87 (dd, $J = 3.0, 12.7$, H_b -C(5')); 4.19 (ddd, $J = 3.0, 4.0, 4.3$, H-C(4')); 4.33 (dd, $J = 4.3, 5.3$, H-C(3')); 4.56 (dd, $J = 5.3, 5.4$, H-C(2')); 5.92 (d, $J = 5.4$, H-C(1')); 7.43 (d, $J = 3.0$, H-C(6)); 7.47 (d, $J = 3.0$, H-C(7)). $^{13}\text{C-NMR}$ ((D_6) DMSO): 61.16 (t, C(5')); 70.10 (d, C(2')); 73.18 (d, C(3')); 85.28 (d, C(4')); 86.85 (d, C(1')); 106.70 (d, C(6)); 115.74 (d, C(7)); 149.36 (s, C(8a)); 150.34 (s, C(2)); 161.74 (s, C(4)). FAB-MS (glycerine): 284 ($[M + 1]^+$), 152 (free base).

An analogous reaction with **10** (300 mg, 0.5 mmol) gave **11** (98 mg, 70%) as a white solid. The anal. data agreed with the data reported in [25].

Enzymatic Synthesis. – **Adenosine (13).** To a soln. of adenine (1.1 mg, 8 μ mol) and 7-methylguanosine (**12**; 11.1 mg, 32 μ mol) in aq. phosphate buffer (1.50 ml; 0.05M KH_2PO_4 , pH 7.0) was added PNPase (4 units, 0.16 ml of a freshly prepared soln. containing 25 μ /ml). Immediately a white precipitate formed. The mixture was allowed to stand at 60° in the water bath with gentle shaking. HPLC monitoring (gradient *A*): adenine (t_R 22 min), **14** (t_R 23 min), **13** (t_R 29 min). The mixture was not worked up. HPLC analysis after 22 h gave a yield of 70% for **13**.

4-Amino-8-(β -D-ribofuranosyl)imidazo[1,2-a]-1,3,5-triazin-2(8H)-one (= 5-Aza-7-deazaisoguanosine; **11**). As described for **13**, with **8** (50.0 mg, 0.3 mmol), **12** (416.0 mg, 1.3 mmol), phosphate buffer (50 ml; 0.05M KH_2PO_4 , pH 7.0), and PNPase (9.0 mg, 126 units). HPLC monitoring (gradient *B*): **8** (t_R 16 min), **11** (t_R 17 min), **14** (t_R 26 min). After 230 h, the mixture was worked up. The precipitate was recovered by centrifugation and the supernatant soln. evaporated until it turned slightly cloudy. The soln. was then loaded on an ion-exchange column (for exper. details, see *General*): **11** (0.35–0.45M $(\text{Et}_3\text{NH})\text{HCO}_3$) followed by **8** (0.5–0.6M $(\text{Et}_3\text{NH})\text{HCO}_3$). The ribonucleoside fractions were lyophilized. The white lyophilizate was dissolved in H_2O and **11** precipitated with MeCN. Drying under high vacuum gave **11** (19 mg, 22%) as white powder. Anal. data: in agreement with those given above.

2'-Deoxy-7-methylguanosinium Iodide (15). The synthetic procedures [26] were modified to achieve reproducible results in our hands. Deoxyguanosine hydrate (20.0 g, 70.1 mmol) was dissolved in DMSO (150 ml) at r.t. MeI (20 ml, 315 mmol) was added at r.t. and the mixture stirred at r.t. for 1 h. The mixture turned orange. Excess MeI was removed by purging with N_2 . The soln. was diluted with CHCl_3 (1200 ml) at r.t. and then cooled to 0° for 3 h. The resulting precipitate was isolated by filtration and washed with cold EtOH (60 ml) and cold Et_2O (80 ml). Drying gave **15** (22.0 g, 77%) as a white powder.

2'-Deoxy-7-methylguanosine (18). Following a slightly modified procedure of [26], **15** (22.0 g, 54 mmol) was suspended in EtOH (1100 ml) at 0° and treated with 40% $\text{MeNH}_2/\text{H}_2\text{O}$ (220 ml). The suspension was stirred at 0° for 1 h. The light white precipitate was isolated by centrifugation and washed with EtOH (350 ml) and Et_2O (400 ml). Drying under high vacuum gave **18** (10.9 g, 66%) as white powder.

2'-Deoxyadenosine (16). To a soln. of adenine (1.1 mg, 8 μ mol) and **15** (13 mg, 32 μ mol) in aq. phosphate buffer (1.50 ml; 0.05M KH_2PO_4 , pH 7.4) was added PNPase (23 units, 0.16 ml of a freshly prepared soln. containing 145 μ /ml). A white precipitate was immediately formed, clouding the initially clear soln. The mixture was allowed to stand at 45° in the water bath under gentle shaking by HPLC monitoring: **14** (t_R 8 min), adenine (t_R 11 min), **16** (t_R 20 min). The mixture was not worked up. HPLC analysis after 16 h gave a yield of 70% for **16**.

4-Amino-8-(β -D-2'-deoxyribofuranosyl)imidazo[1,2-a]-1,3,5-triazin-2(8H)-one (= 2'-Deoxy-5-aza-7-deaza-isoguanosine; **17**). The reaction was performed in 5 cycles. After reaching the reaction equilibrium, the enzyme was recovered and used for the next cycle. Each cycle was performed as follows: To a soln. of **8** (25 mg, 156 μ mol) and **18** (380 mg, 1.25 mmol) in aq. phosphate buffer (32 ml; 0.05M in KH_2PO_4 , pH 7.4) was added PNPase (80 mg, 1120 μ). Immediately a white precipitate formed making the soln. cloudy. The mixture was allowed to stand at 45° in the water bath with gentle shaking. HPLC monitoring (gradient *B*): **8** (t_R 16 min), **17** (t_R 19 min), **14** (t_R 26 min). After reaching equilibrium (first cycle 24 h, later up to 36 h), the mixture was worked up. The precipitate was recovered by centrifugation and the enzyme recycled (for exper. details, see *General*) from the supernatant soln. After 5 cycles, the supernatant solns. were pooled and lyophilized. The lyophilizate was dissolved in little H_2O and loaded on an ion-exchange column (for exper. details, see *General*): **17** (0.25–0.4M $(\text{Et}_3\text{NH})\text{HCO}_3$) followed by **8** (0.5–0.6M $(\text{Et}_3\text{NH})\text{HCO}_3$). The deoxyribonucleoside fractions were lyophilized. The lyophilizate was desalted by reversed-phase chromatography (30 g of silica gel *RP-C18*, 3 ml fractions, H_2O (70 ml), then $\text{H}_2\text{O}/\text{MeOH}$ 4:1 (10 ml)). Again the deoxyribonucleoside fractions were lyophilized. Drying under high vacuum gave **17** (112 mg, 54%) as white powder. An anal. sample was obtained by recrystallization from EtOH. M.p. 210° (dec.). UV (EtOH): 246 (12900). IR (KBr): 3380, 3330, 3140, 3100, 3010, 2940, 2880, 1675, 1635, 1610, 1530, 1460, 1380, 1280, 1245, 1120, 1070. $^1\text{H-NMR}$ (D_2O): 2.46 (ddd, $J = 4.0, 6.6, 14.0$, $\text{H}_a\text{-C}(2'')$); 2.58 (ddd, $J = 6.6, 6.6, 14.0$, $\text{H}_b\text{-C}(2'')$); 3.71 (dd, $J = 4.8, 12.6$, $\text{H}_a\text{-C}(5'')$); 3.79 (dd, $J = 3.5, 12.6$, $\text{H}_b\text{-C}(5'')$); 4.04–4.07 (m, $\text{H-C}(4'')$); 4.52–4.55 (m, $\text{H-C}(3'')$); 6.28 (dd, $J = 6.6, 6.6$, $\text{H-C}(1'')$); 7.39 (d, $J = 3.0$, $\text{H-C}(6)$); 7.42 (d, $J = 3.0$, $\text{H-C}(7)$). $^{13}\text{C-NMR}$ (D_2O): 41.08 (t, $\text{C}(2'')$); 64.10 (t, $\text{C}(5'')$); 73.49 (d, $\text{C}(3'')$); 86.46 (d, $\text{C}(4'')$); 89.72 (d, $\text{C}(1'')$); 110.45 (d, $\text{C}(6)$); 119.23 (d, $\text{C}(7)$); 152.93 (s, $\text{C}(2)$); 154.08 (s, $\text{C}(8a)$); 168.60 (s, $\text{C}(4)$). FAB-MS (3-nitrobenzyl alcohol: 268 ($[M + 1]^+$), 152 (free base). Anal. calc. for $\text{C}_{10}\text{H}_{13}\text{N}_5\text{O}_4 \cdot \text{H}_2\text{O}$ (285.26): C 42.10, H 5.30, N 24.55; found: C 42.18, H 5.16, N 24.26.

A scaled-up run was performed in 8 cycles using **8** (800 mg, 4.96 mmol) overall and yielded **17** (830 mg, 59%) as a white solid, contaminated with Et_3NH^+ salts. After resolution of the mixture with ion-exchange chromatography, the fractions containing **17** were lyophilized. The product was then recovered by precipitation with EtOH. This simplified procedure avoided the desalting by reversed-phase chromatography.

REFERENCES

- [1] W. J. Hennen, C-H. Wong, *J. Org. Chem.* **1989**, *54*, 4692.
- [2] E. Lukevics, A. Zablocka, 'Nucleoside Synthesis. Organosilicon Methods 1991', Ellis Horwood Limited, London, 1991.
- [3] H. Vorbrüggen, K. Krolikiewicz, B. Bennua, *Chem. Ber.* **1981**, *114*, 1234.
- [4] H. Vorbrüggen, G. Höfle, *Chem. Ber.* **1981**, *114*, 1256.
- [5] H. Vorbrüggen, B. Bennua, *Chem. Ber.* **1981**, *114*, 1279.
- [6] C. Hildebrand, G. E. Wright, *J. Org. Chem.* **1992**, *57*, 1808.
- [7] H. Kawakami, T. Ebata, K. Koseki, H. Matsushita, Y. Naoi, K. Itoh, N. Mizutani, *Heterocycles* **1990**, *31*, 569.
- [8] J. N. Freskos, *Nucleos. Nucleot.* **1989**, *8*, 549.
- [9] P. Garner, S. Ramakanth, *J. Org. Chem.* **1988**, *53*, 1294.
- [10] F. Seela, H. Rosemeyer, W. Bourgeois, *Nucleic Acids Res. Symp. Series* **1987**, *18*, 49.
- [11] H. Rosemeyer, F. Seela, *J. Org. Chem.* **1987**, *52*, 5136.
- [12] A. J. Hubbard, A. S. Jones, R. T. Walker, *Nucleic Acids Res.* **1984**, *12*, 6827.
- [13] Z. Kazimierczuk, H. B. Cottam, G. R. Revankar, R. K. Robins, *J. Am. Chem. Soc.* **1991**, *106*, 6379.
- [14] D. G. Drueckhammer, W. J. Hennen, R. L. Pederson, C. F. Barbas, C. M. Gautheron, T. Krach, C-H. Wong, *Synthesis* **1991**, 499.
- [15] D. W. Hutchinson, *TIBTECH* **1990**, *8*, 348.
- [16] J. Holguin, R. Cardinaud, C. A. Salemink, *Eur. J. Biochem.* **1975**, *54*, 515.
- [17] A. Bzowska, E. Kulikowska, D. Shugar, *Z. Naturforsch., C* **1990**, *45*, 59.
- [18] H. Shirae, K. Yokozeki, K. Kubota, *Agric. Biol. Chem.* **1988**, *52*, 295.
- [19] T. A. Krenitsky, J. L. Rideout, E. Y. Chao, G. W. Koszalka, F. Gurney, R. C. Crouch, N. K. Cohn, G. Wolberg, R. Vinegar, *J. Med. Chem.* **1986**, *29*, 138.
- [20] T. A. Krenitsky, G. W. Koszalka, J. V. Tuttle, *Biochemistry* **1981**, *20*, 3615.
- [21] J. Dosekocil, A. Holy, *Collect. Czech. Chem. Commun.* **1977**, *42*, 370.
- [22] H. Shirae, K. Yokozeki, *Agric. Biol. Chem.* **1991**, *55*, 1849.
- [23] T. Utagawa, H. Morisawa, S. Yamanaka, A. Yamazaki, F. Yoshinaga, Y. Hirose, *Agric. Biol. Chem.* **1985**, *49*, 3239.
- [24] J. A. Piccirilli, T. Krauch, S. E. Moroney, S. A. Benner, *Nature (London)* **1990**, *343*, 33.
- [25] E. J. Prisbe, J. P. H. Verheyden, J. G. Moffatt, *J. Org. Chem.* **1978**, *43*, 4784.
- [26] J. W. Jones, R. K. Robins, *J. Am. Chem. Soc.* **1963**, *85*, 193.
- [27] R. G. Fargher, F. L. Pyman, *J. Chem. Soc., Perkin Trans.* **1919**, *115*, 217.
- [28] R. Esmail, F. Kurzer, *Synthesis* **1975**, 301.
- [29] M. Nagano, J. Tobitsuka, T. Matsui, K. Oyamada, *Chem. Pharm. Bull.* **1972**, *20*, 2618.
- [30] S. Ram, R. E. Eherenkauf, *Synthesis* **1988**, 91.
- [31] M. K. Anwer, F. Spatola, *Synthesis* **1980**, 929.
- [32] B. El Amin, G. M. Aranthanamaiah, G. P. Royer, G. E. Mens, *J. Org. Chem.* **1979**, *44*, 3442.
- [33] A. M. Felix, E. P. Heimer, Th. J. Lambros, Ch. Tzougraki, J. Meienhofer, *J. Org. Chem.* **1978**, *43*, 4194.
- [34] V. du Vigneaud, O. K. Behrens, *J. Biol. Chem.* **1937**, *117*, 27.
- [35] K. A. Watanabe, D. H. Hollenberg, J. J. Fox, *J. Carbohydr. Nucleos. Nucleot.* **1974**, *1*, 1.
- [36] S-H. Kim, D. G. Bartholomew, L. B. Allen, R. K. Robins, G. R. Revankar, P. Dea, *J. Med. Chem.* **1978**, *21*, 883.
- [37] H. Rosemeyer, G. Toth, F. Seela, *Nucleos. Nucleot.* **1989**, *8*, 587.
- [38] E. J. Prisbe, J. P. H. Verheyden, J. G. Moffatt, *J. Org. Chem.* **1978**, *43*, 4774.