



Cyclopropanation of alkenes with $\text{CH}_2\text{I}_2/\text{Et}_3\text{Al}$ by the phase-vanishing method based on fluororous phase screen

Hiroshi Matsubara*, Masaaki Tsukida, Shinji Yasuda, Ilhyong Ryu*

Department of Chemistry, Graduate School of Science, Osaka Prefecture University, Sakai, Osaka 599-8531, Japan

ARTICLE INFO

Article history:

Received 24 March 2008

Received in revised form 5 June 2008

Accepted 5 June 2008

Available online 14 June 2008

This paper is dedicated to Professor Dennis P. Curran for his great contribution to fluororous chemistry.

Keywords:

Fluorous solvent

Phase-vanishing

Simmons–Smith cyclopropanation

Diiodomethane

Triethylaluminum

ABSTRACT


Phase-vanishing (PV) method using perfluorohexanes as a screen phase was applied to cyclopropanation reactions with $\text{CH}_2\text{I}_2/\text{Et}_2\text{Zn}$ and $\text{CH}_2\text{I}_2/\text{Et}_3\text{Al}$. When Et_3Al was used as a carbenoid generator, the reaction proceeded smoothly and desired cyclopropane derivatives were obtained in high yield. The PV cyclopropanation took 2 or 3 days to complete, however, reduction of reaction time by a factor of 2–3 was also achieved by vigorous stirring after the bottom CH_2I_2 layer disappeared.

© 2008 Elsevier B.V. All rights reserved.

1. Introduction

Fluorous phase chemistry has opened up a new fresh ground in the way of organic synthesis and separation process [1]. Perfluorinated compounds are generally immiscible with most organic solvents and are denser than typical organic molecules. By utilizing the unique properties of fluororous solvents inherent to perfluoroalkyl moieties, we recently reported a synthetically convenient triphasic system [2]: a phase-vanishing (PV) method, in which fluororous solvents, such as FC-72 (perfluorohexanes) and Garden HT-135 (polyperfluoroalkyl ethers), act as a screen phase to connect two separated layers. The PV method is demonstrated to be useful in a variety of situations including: for the bromination of alkenes [3] or at benzylic positions [4] with Br_2 , demethylation of methyl ethers [3] with BBr_3 , bromination of alcohols [5] with SOBr_2 or PBr_3 , and Friedel–Crafts acylation of aromatic compounds [6] with SnCl_4 under mild conditions. The PV method has other variations for example using: chlorine gas [7], lighter reagents [5] than fluororous solvents, and solid reagents [8]. We also introduced a “quadraphasic PV” method [9], in which an aqueous phase was added to the original triphasic PV method to remove acidic by-products.


Diiodomethane (CH_2I_2), a denser liquid ($d = 3.33$) than FC-72 (perfluorohexanes: $d = 1.67$), has been used for the Simmons–Smith cyclopropanation reactions as the methylene source [10–12]. The Furukawa variant [11] uses the combination of diethylzinc and CH_2I_2 to give cyclopropane derivatives usually in improved yields as compared with the original Simmons–Smith conditions ($\text{Zn–Cu}/\text{CH}_2\text{I}_2$). The Maruoka/Yamamoto variant [12] uses trialkylaluminum to generate the carbenoid from CH_2I_2 to afford cyclopropanation products in good yield. In both variants, cooling of the reaction mixture by carefully controlled slow addition of CH_2I_2 is required to avoid rapid heat evolution. Herein we wish to



Scheme 1. A concept of phase-vanishing cyclopropanation using diiodomethane.

* Corresponding author. Tel.: +81 72 254 9695; fax: + 81 72 254 9695.
E-mail address: ryu@c.s.osakafu-u.ac.jp (I. Ryu).

Table 1
Control experiments for phase-vanishing cyclopropanation of 1-dodecene



Entry	Conditions	Procedure ^a	Isolated yield of 1 (%)
1	Et ₃ Al (2 mmol), CH ₂ I ₂ (1.3 mmol)	A	78
2	Et ₃ Al (2 mmol), CH ₂ I ₂ (2.0 mmol)	A	98
3	Et ₃ Al (2 mmol), CH ₂ I ₂ (2.0 mmol)	B	98
4	Et ₂ Zn (2 mmol), CH ₂ I ₂ (1.3 mmol)	A	20
5	Et ₂ Zn (2 mmol), CH ₂ I ₂ (2.0 mmol)	A	31
6 ^b	Et ₂ Zn (2 mmol), CH ₂ I ₂ (2.0 mmol)	A	39

^a Procedure A: gentle stirring, 36 h; Procedure B: vigorous stirring, 4 h.

^b Under dry air atmosphere.

report that a phase-vanishing method can be applied to cyclopropanation reaction of olefins using CH₂I₂/Et₃Al or CH₂I₂/Et₂Zn, which can be conducted at ambient temperature without the need for slow addition of reagents (Scheme 1).

2. Results and discussion





We examined the phase-vanishing cyclopropanation of 1-dodecene as a model; the results of this study are summarized in Table 1. Diiodomethane ($d = 3.33$) was overlaid by FC-72 ($d = 1.67$), which in turn was overlaid by a hexane solution of 1-dodecene and triethylaluminum. The bottom layer was gently stirred, taking care

not to mix the three layers. The diiodomethane layer disappeared after 8 h, giving two layers, however the reaction was not complete at this stage. The reaction took almost 2 days to finish (Procedure A). The organic phase was collected and after column chromatography on silica gel decylcyclopropane was obtained in 78% yield (entry 1, Table 1). In a similar procedure, the use of 2 equivalents of diiodomethane increased the yield to 98% (entry 2, Table 1). In contrast, diethylzinc was not observed to be as effective as triethylaluminum in the PV procedure. With this reagent, while extending reaction time or increasing the molar ratio of diiodomethane did not increase the product yield (entries 4 and 5, Table 1), exposure of the reaction mixture to oxygen slightly

Table 2
Phase-vanishing cyclopropanation of alkenes by Et₃Al/CH₂I₂^a

Entry	Substrate	Procedure	Product	Isolated yield (%) ^b
1		A		98
2		B		98
3 ^c		A		98 <i>cis/trans</i> = 2.14/1 ^d
4 ^{c,e}		A		98 <i>cis/trans</i> = 2.12/1 ^d
5		C		93
6		C		97
7		C		99

Table 2 (Continued)

Entry	Substrate	Procedure	Product	Isolated yield (%) ^b
8		C	 6	97
9		C	 7	92
10		D		92
11		C	 8	92

^a General conditions: substrate (1 mmol), CH₂I₂ (2 mmol), hexane solution of Et₂Zn or Et₃Al (1 M, 2 mL), FC-72 (1.5 mL) with aluminum foil protection from light, kept at room temperature with gentle stirring. Procedure A: 36 h; Procedure B: 4 h with vigorous stirring; Procedure C: 58 h; Procedure D: 16 h with vigorous stirring.

^b Isolated yield by silica gel chromatography.


^c *Cis/trans* = 2.43/1.

^d Determined by ¹H-NMR.

^e Galden HT-135 (1.5 mL) was used as a fluorous phase.

improved the yield (entry 6, Table 1). In order to reduce the reaction time for the reaction of Et₃Al/CH₂I₂, after the bottom layer had disappeared, the test tube was stirred vigorously for a further 4 h (Procedure B). This modification also gave a good result (entry 3, Table 1).

Encouraged by the excellent adaptability of the Maruoka–Yamamoto cyclopropanation reaction to the PV method, we embarked on experiments with several other alkenes. In each case 2 equiv. of CH₂I₂ and Et₃Al was used (Table 2). In all cases examined, the cyclopropanation reaction took place to give high yields of the desired cyclopropanes. Styrene derivatives required longer reaction times to complete than aliphatic alkenes (Procedure C: with extended reaction time (58 h) after the bottom phase disappeared) (entries 5–9, 11, Table 2). Coupled with the use of vigorous stirring after the bottom layer disappeared, the total reaction time was able to be shortened by a factor of 2–3 (entries 2 and 10, Table 2). Galden[®] HT-135 is a polyether-type perfluorinated solvent, commercially available as a heat transfer liquid (Solvay Solexis Inc.), bp 135 °C, density = 1.72 g/mL at 25 °C, average molecular weight = 610. This inexpensive polyperfluoro ether solvent was shown to function equally as well with FC-72 (entry 4).



3. Conclusion

Through the use of fluorous media, such as FC-72 (perfluoro-hexanes) and Garden HT-135, as a phase screen, we have demonstrated that cyclopropanation of alkenes can be easily carried out conveniently without the necessity for slow addition of diiodomethane and equipment for temperature control, which are necessary for the conventional glass-flask reaction. Whereas the Furukawa reagent (Et₂Zn/CH₂I₂) did not function satisfactorily for the present test tube-based triphasic system, the Maruoka–Yamamoto reagent (Et₃Al/CH₂I₂) gave the desired cyclopropanation products in high yields. This carbenoid reagent coupled with

the vigorous stirring of the resulting two layers after the bottom layer disappeared significantly reduced the reaction time.

4. Experimental

FC-72 and Galden[®] HT-135 were purchased from Sumitomo 3 M Ltd., and Solvay Solexis Inc., respectively, and dried over molecular sieves 4 Å before use. Reagents and solvents were used as received. Hexane solution of diethylzinc (1 M) and triethylaluminum (1 M) were purchased from Kanto Chemical Co., Inc., and used as received. Products were purified by column chromatography on silica gel (Kanto Chemical Co., Inc., Silica Gel 60N, 70–230 mesh). ¹H NMR spectra were recorded with a JEOL JMN-500 (500 MHz) or a JEOL JMN-400 (400 MHz) spectrometer while ¹³C NMR spectra were recorded with a JEOL JMN-500 (125 MHz) or a JEOL JMN-400 (100 MHz) spectrometer. Infrared spectra were obtained on a JASCO FT/IR-4100 spectrometer. Conventional and high-resolution mass spectra were recorded with a Shimadzu GCMS-QP 5050A instrument and a JEOL MS-700 spectrometer, respectively.

4.1. General procedure for cyclopropanation of olefins by the phase-vanishing method (Table 1, entry 2) (Procedure A)

FC-72 (1.5 mL) was placed in a pyrex test tube (13 mm φ × 105 mm) to which diiodomethane (2.0 mmol, 536 mg) was added slowly using a glass pipette under argon atmosphere. 1-Dodecene (1.0 mmol, 168 mg) was then added slowly, forming three layers. A hexane solution (2 mL) of triethylaluminum (1 M) was added to the top layer, and the test tube was covered with aluminum foil in order to shield the reaction from light and kept it at room temperature. The diiodomethane layer was gently stirred using a magnetic stirrer, taking care not to mix the three layers. The diiodomethane layer disappeared after 8 h, the hexane layer was taken up with a pipette after 44 h. Additional hexane (3 mL × 4) was placed on the residual FC-72 layer, then decanted off. The combined organic layer was washed with aqueous 0.5 M HCl (20 mL) and saturated brine (20 mL), dried over Na₂SO₄, and concentrated. Purification by short column chromatography on silica gel with hexane gave decylcyclopropane (**1**, 178 mg, 98%) as a colorless oil; IR (neat): ν 3000, 2955, 2925, 2850, 1465, 1375, 1040, 1015, 820, 720 cm⁻¹; ¹H NMR (500 MHz, CDCl₃): δ-0.12 (2H, m, CH₂ of cyclopropane), 0.38 (2H, m, CH₂ of cyclopropane), 0.88 (3H, t, J = 6.9 Hz, CH₃), 1.17 (2H, q, J = 7.4, CH₂-c-C₃H₆), 1.26–1.42 (16H,

m, CH₂); ¹³C NMR (125 MHz, CDCl₃): δ 4.35, 10.92, 29.39, 29.54, 59.59, 29.66, 29.69, 29.72, 29.75, 31.96, 33.85, 34.81; EI MS 70 eV, *m/z* (relative intensity): 182 [M]⁺ (10), 168 [M-CH₂]⁺ (13), 154 [M-C₂H₄]⁺ (14), 140 [M-C₃H₆]⁺ (8), 125 (19), 111 (45), 97 (93), 83 (100), 69 (88), 55 (66); HRMS (EI): *m/z* = [M]⁺ calcd for C₁₃H₂₆: 182.2035; found: 182.2036.

All other products **2**[13], **3**[14], **4**[15], **5**[14], **6**[14], **7**[14] and **8**[16] are known compounds and were identified by spectral data with comparison of literature data.

Acknowledgements

We thank JSPS for financial support of this work. We also thank Ms. Sara H. Kyne for useful suggestions on the manuscript.

References

- [1] J.A. Gladysz, D.P. Curran, I.T. Horváth (Eds.), For A General Review of Fluorous Chemistry: Handbook of Fluorous Chemistry, Wiley-VCH, Weinheim, 2004.
- [2] (a) Z. Luo, S.M. Swaleh, F. Theil, D.P. Curran, *Org. Lett.* 4 (2002) 2585–2587; (b) H. Nakamura, B. Linclau, D.P. Curran, *J. Am. Chem. Soc.* 123 (2001) 10119–10120.
- [3] I. Ryu, H. Matsubara, S. Yasuda, H. Nakamura, D.P. Curran, *J. Am. Chem. Soc.* 124 (2002) 12946–12947.
- [4] A. Podgoršek, S. Stavber, M. Zupan, J. Iskra, *Eur. J. Org. Chem.* (2006) 483–488.
- [5] H. Nakamura, T. Usui, H. Kuroda, I. Ryu, H. Matsubara, S. Yasuda, D.P. Curran, *Org. Lett.* 5 (2003) 1167–1169.
- [6] H. Matsubara, S. Yasuda, I. Ryu, *Synlett* (2003) 247–249.
- [7] J. Iskra, S. Stavber, M. Zupan, *Chem. Commun.* (2003) 2496–2497.
- [8] (a) N.K. Jana, J.G. Verkade, *Org. Lett.* 5 (2003) 3787–3790; (b) D.P. Curran, S. Werner, *Org. Lett.* 6 (2004) 1021–1024.
- [9] Md.T. Rahman, N. Kamata, H. Matsubara, I. Ryu, *Synlett* (2005) 2664–2666.
- [10] For leading reviews of the Simmons-Smith reactions.
 - (a) H.E. Simmons, T.L. Cairns, S.A. Vladuchick, C.M. Hoiness, *Org. React.* 20 (1973) 1–131;
 - (b) A.B. Charette, A. Beauchemin, *Org. React.* 58 (2001) 1–415.
- [11] (a) J. Furukawa, N. Kawabata, J. Nishimura, *Tetrahedron Lett.* (1966) 3353–3354; (b) J. Furukawa, N. Kawabata, *J. Nishimura, Tetrahedron* 24 (1968) 53–58.
- [12] K. Maruoka, Y. Fukutani, H. Yamamoto, *J. Org. Chem.* 50 (1985) 4412–4414.
- [13] E.J. O'Connor, S. Brandt, P. Helquist, *J. Am. Chem. Soc.* 109 (1987) 3739–3747.
- [14] J.C. Lorenz, J. Long, Z. Yang, S. Xue, Y. Xie, Y. Shi, *J. Org. Chem.* 69 (2004) 327–334.
- [15] J.R.A. Dulayymi, M.S. Baird, I.G. Bolesov, A.V. Nizovtsev, V.V. Tverezovsky, *J. Chem. Soc. Perkin Trans. 2* (2000) 1603–1617.
- [16] M.-C. Lacasse, C. Poulard, A.B. Charette, *J. Am. Chem. Soc.* 127 (2005) 12440–12441.