Tetrahedron Letters 50 (2009) 1450-1452

Contents lists available at ScienceDirect

Tetrahedron Letters

journal homepage: www.elsevier.com/locate/tetlet

A simple, novel and convenient method for the synthesis of 1-aminophosphinic acids: synthesis of a novel C₂-symmetric phosphinic acid pseudodipeptide

Babak Kaboudin*, Fariba Saadati

Department of Chemistry, Institute for Advanced Studies in Basic Sciences (IASBS), Gava Zang, Zanjan 45195-1159, Iran

ARTICLE INFO	ABSTRACT
Article history: Received 2 November 2008 Revised 23 December 2008 Accepted 13 January 2009 Available online 19 January 2009	A simple, efficient and novel method has been developed for the synthesis of 1-aminophosphinic acids from simple starting materials. Treatment of aromatic aldehydes with ammonia and hypophosphorus
	acid gives novel C ₂ -symmetric 1-aminoarylmethylphosphinic acids. The synthesis of novel C ₂ -symmetric phosphinic acid pseudodipeptides is also discussed.
Available office 13 January 2009	© 2009 Elsevier Ltd. All rights reserved.

Phosphinic acids are of growing importance in understanding and modulating biological processes.¹ In recent years, the synthesis of α -substituted phosphoryl derivatives (phosphonic and phosphinic acids) has attracted significant attention, due to their biological activities with broad application as enzyme inhibitors, antimetabolites and antibiotics.² Among α -functionalized phosphinic acids, α -aminoalkylphosphinic derivatives have potential biological activities, such as anti-bacterial,² herbicidal³ and fungicidal.⁴ 1-Aminoalkylphosphinic acids, the phosphinic acid analogues of 1amino carboxylic acids, are an important class of compounds that exhibit a variety of interesting and useful properties. In addition, the structure of the phosphinic functional group mimics the transition state of peptide hydrolysis, and the symmetric nature of the phosphinic acid derivatives is expected to be of benefit in their binding to the homodimer of HIV-protease having C2-axis symmetry.

A phosphinic moiety (-PO₂-CH₂-) is considered a mimic of the tetrahedral transition state of peptide bond hydrolysis and due to this, phosphinate pseudopeptides have attracted considerable interest providing a wide range of potent inactivators of proteolytic enzymes, particularly metalloproteases.⁵ In contrast to the widely studied 1-aminoalkylphosphonic acid derivatives,⁶⁻⁹ relatively few papers have reported the chemistry of 1-aminoalkylphosphinic acids, although there is evidence that α -aminophosphinic acids are pharmaceutically active. Synthetic routes to 1-aminoalkylphosphinic acids involve Mannich-type reaction of amines with aldehydes in the presence of anhydrous hypophosphorus acid and prolonged heating of anhydrous hypophosphorus acid with a Schiff's base.¹⁰ The addition of hypophosphorus acid to structurally diverse imines appears to be a general method for the preparation of N-substituted 1-aminoalkanephosphinic acids.¹¹ However, both of these methods have drawbacks, such as harsh reaction conditions, anaerobic and anhydrous conditions, long reaction times and also side reactions. On the other hand, the key step in the one-pot synthesis of 1-aminoalkylphosphinic acids is nucleophilic addition of an amine to a carbonyl compound followed by addition of hyphophosphorus acid to the resulting imine. Therefore, the formation of 1-hydroxyalkylphosphinic acids frequently accompanies the formation of 1-aminoalkylphosphinic acids.¹² Hydrophosphinvlation of *N*,*N*'-arylidene bisamides has been reported as a new method for the synthesis of 1-aminophosphinic acids.¹³ However, this method also has problems, and includes difficulties in the preparation of starting materials and requires harsh reaction conditions for hydrolysis of the amide intermediate to the amine product.

Diimines serve as good precursors for the synthesis of numerous organic compounds, especially heterocyclic compounds.14,15 These, easily accessible precursors can be produced by the reaction of aromatic aldehydes with ammonia solution. Recently, we reported the reaction of diimines with diethyl phosphite for the prepartion of bis(1-diethoxyphosphorylalkyl)amines.¹⁶ As part of our efforts to introduce novel methods for the synthesis of organophosphorus compounds,¹⁷ herein we report a new method for the synthesis of α -aminophosphinic acids and novel C_2 -symmetric phosphinic acid pseudodipeptides. We have found that reaction of aromatic aldehvdes with ammonia solution followed by reaction with hypophosphorus acid gives 1-aminophosphinic acids in good yields (Scheme 1).

Thus, the reaction of benzaldehyde, chosen as a model compound, with ammonia followed by treatment with hypophosphorus acid was studied under various reaction conditions. When the reaction was conducted under reflux for 10 h using 5 equiv of



Scheme 1.







^{*} Corresponding author. Tel.: +98 241 4153220; fax: +98 241 4214949. E-mail address: kaboudin@iasbs.ac.ir (B. Kaboudin).

^{0040-4039/\$ -} see front matter © 2009 Elsevier Ltd. All rights reserved. doi:10.1016/j.tetlet.2009.01.055



Scheme 2.

H₃PO₂ (anhydrous), the ³¹P NMR spectrum of the product exhibited two major peaks at δ 17.85 and 17.58 due to the diastereoisomers of **2a**. The ¹H NMR spectrum exhibited two doublets at δ 4.16 and 3.79 indicative of HC–P coupling ($J_{\rm HP}$ = 13.5 Hz). Two doublets at δ 6.88 and 6.83 were indicative of H–P coupling (J_{HP} = 552 Hz). Due to the presence of two stereogenic carbons bonded to a phosphorus atom, and due to the prototopic transfer of the acidic proton between the phosphoryl (P=O) and acidic (P-OH) sites, these compounds exist as two diastereomeric forms: one meso-compound $(R^{\uparrow}, S^{\uparrow}-2\mathbf{a})$ and one racemic pair $(S^{\uparrow}, S^{\uparrow})$ or $R^{\uparrow}, R^{\uparrow}-2\mathbf{a}$, as shown in Scheme 2. When the mixture of diastereoisomers 2a was washed with a solvent mixture of ethanol/water (9:1) and dried in air at room temperature, a single diastereoisomer was obtained. As we were unable to obtain an X-ray crystal of the two diastereoisomers of 2a, their unambiguous stereochemical assignments have not been determined. It should be noted that when the reaction was carried out with 50% hypophosphorus acid, 1-hydroxyphenyl methylphosphinic acid was obtained as the major product.

This process was applied successfully to other aromatic aldehydes as summarized in Table 1. Substituted benzaldehydes reacted with ammonia followed by reaction with hypophosphorus acid to afford the desired products **2b–f** in moderate to good yields. 2-Naphthalene carbaldehyde also reacted with hypophosphorus acid in the presence of ammonia to give compound **2g** in 40% yield. Reaction of aliphatic aldehyde **1h** with hypophosphorus acid in the presence of ammonia gave an unidentified mixture of products.

 Table 1

 Synthesis of 1-aminophosphinic acids 2 from aldehydes

Entry	Ar	Reaction time (h)	Yield ^a (%)
a	C ₆ H ₅ -	10	48
b	p-ClC ₆ H ₄ -	11	71
с	p-FC ₆ H ₄ -	12	61
d	p-MeOC ₆ H ₄ -	12	45
e	p-MeC ₆ H ₄ -	11	64
f	o-FC ₆ H ₄ -	2	42
g	2-Naphthyl	8	40
h	n-C ₆ H ₁₃ -	-	b

^a Isolated yields of mixtures of two diastereoisomers.

^b Unknown products obtained.



1a

Scheme 3.



Scheme 4.

These results prompted us to extend this process to phosphorus acid for the synthesis of 1-aminophosphonic acids (Scheme 3). Initially, we carried out the reaction of benzaldehyde **1a** with ammonia in the presence of phosphorus acid under reflux conditions. Unfortunately, the reaction failed, and no product was detected after 24 h at reflux.

Phosphorus–carbon bond formation via Michael addition is a useful method for the synthesis of phosphinic acid dipeptides with application in organic synthesis and bioorganic chemistry. A novel C_2 -symmetric phosphinic acid dipeptide was synthesized by Michael addition of **2a** to methyl acrylate. 1-Aminophosphinic acid **2a** (single diastereoisomer) was activated by silylation with HMDS¹⁸ then reacted with methyl acrylate to afford adduct **3a** in 85% yield (Scheme 4).

In summary, the simple work-up, mild reaction conditions, good yields and clean reactions with no tar formation make this method an attractive and useful contribution to present methodologies.¹⁹ A novel C_2 -symmetric phosphinic acid dipeptide was also synthesized by Michael addition of **2a** to methyl acrylate.²⁰ Further investigations on the stereochemistry of the products and their complexation ability are now in progress.

Acknowledgement

The Institute for Advanced Studies in Basic Sciences (IASBS) is thanked for supporting this work.

References and notes

- (a) Engel, R. Chem. Rev. 1977, 77, 349–367; (b) Moonen, K.; Laureyn, I.; Stevens, C. V. Chem. Rev. 2004, 104, 6177–6215.
- Collinsova, M.; Jiracek, J. Curr. Med. Chem. 2000, 7, 629–647.
 Kafarski, P.; Lejczak, B.; Tyka, R.; Koba, L.; Pliszczak, E.; Wieczorek, P. J. Plant
- Growth Regul. 1995, 14, 199–203. 4 Ishipuri Y. Yamada Y. Kato T. Sasaki M. Mukai K. Fur. Pat. Appl. FP.82-
- Ishiguri, Y.; Yamada, Y.; Kato, T.; Sasaki, M.; Mukai, K. Eur. Pat. Appl., EP 82-301905, 1982; Chem. Abstr. 1983, 98, 102686.
- (a) Latajka, R.; Krężel Mucha, A.; Jewgiński, M.; Kafarski, P. J. Mol. Struct. 2008, 877, 64–71; (b) Yamagishi, T.; Ichikawa, H.; Haruki, T.; Yokomatsu, T. Org. Lett. 2008, 10, 4347–4350. and references cited therein; (c) Liu, X.; Hu, X. E.; Tian, X.; Mazur, A.; Ebetino, F. H. J. Organomet. Chem. 2002, 646, 212–222.
- (a) Gancarz, R.; Chakraborty, S. Synthesis 1977, 625; (b) Giannousis, P. P.; Bartlett, P. A. J. Med. Chem. 1987, 30, 1603–1609; (c) Maier, L.; Lea, P. J. Phosphorus, Sulfur, Silicon 1983, 17, 1–19; (d) Hilderbrand, R. L. In The Role of Phosphonates in Living Systems; CRC Press: Boca Raton, 1982. F1; (e) Kafarski, P.; Lejczak, B. Phosphorus, Sulfur, Silicon 1991, 63, 193–215; (f) Kukhar, V. P.; Hudson, H. R. In Aminophosphonic and Aminophosphinic Acids; John Wiley & Sons, 2000; (g) Hanessian, S.; Bennani, Y. L. Synthesis 1995, 1272.
- (a) Redmore, D.. In *Topics in Phosphorus Chemistry*; Griffith, E. J., Grayson, M., Eds.; Wiley: New York, 1976; Vol. 8, (b) Atherton, F. R.; Hassall, C. H.; Lambert, R. W. J. Med. Chem. **1986**, 29, 29–40; (c) Allen, M. C.; Fuhrer, W.; Tuck, B.; Wade, R.; Wood, J. M. J. Med. Chem. **1989**, 32, 1652–1661; (d) Hassall, C. H.. In Antibiotics; Hahn, F. E., Ed.; Springer: Berlin, 1983; Vol. VI, pp 1–11; (e) Gancarz, R.; Wieczorek, J. S. Synthesis **1978**, 625; (f) Seyferth, D.; Marmor, R. S.; Hilbert, P. J. Org. Chem. **1971**, 36, 1379–1386; (g) Worms, K. H.; Schmidt-Dunker, M.. In Organic Phosphorus Compounds; Kosolapoff, G. M., Marier, L., Eds.; John Wiley & Sons: New York, 1976; Vol. 7., p 1 (h) Kaboudin, B. Phosphorus, Sulfur, Silicon **2002**, 177, 1749–1751; (i) Bhagat, S.; Chakraborti, A. K. J. Org. Chem. **2007**, 72, 1263–1270.
- (a) Barycki, J.; Mastalerz, P.; Soroka, M. Tetrahedron Lett. **1970**, 36, 3147–3150;
 (b) Kaboudin, B.; Sorbiun, M. Tetrahedron Lett. **2007**, 48, 9015–9017.
- (a) Genet, J. P.; Uziel, J.; Port, M.; Touzin, A. M.; Roland, S.; Thorimbert, S.; Tanier, S. *Tetrahedron Lett.* **1992**, *33*, 77–80; (b) Qian, C.; Huang, T. J. Org. Chem. **1998**, *63*, 4125–4128; (c) Ranu, B. C.; Hajira, A.; Jana, U. Org. Lett. **1999**, *1*, 1141– 1143; (d) Manabe, K.; Kobayashi, S. Chem. Commun. **2000**, *669–670*; (e) Chandrasekhar, S.; Prakash, S. J.; Jagadeshwar, V.; Narsihmulu, C. Tetrahedron Lett. **2001**, *42*, 5561–5563.

- (a) Schmidt, H. Chem. Ber. **1948**, *81*, 477–483; (b) Linfield, W. M.; Jungermann, E.; Guttmann, A. T. J. Org. Chem. **1961**, *26*, 4088–4092; (c) Yamagishi, T.; Haruki, T.; Yokomatsu, T. Tetrahedron **2006**, *62*, 9210–9217; (d) Cates, L. A.; Li, V. S. Pharm. Res. **1985**, 136; (e) Lecoq, A.; Yiotakis, A.; Dive, V. Synth. Commun. **1994**, 24, 2877–2892; (f) Hamilton, R.; Walker, B.; Walker, B. J. Tetrahedron Lett. **1995**, 36, 4451–4454; (g) Kaboudin, B.; As-Habei, N. Tetrahedron Lett. **2003**, *44*, 4243– 4245; (h) Pirat, J.-L.; Monbrun, J.; Virieux, D.; Volle, J.-N.; Tillard, M.; Cristau, H.-J.J. Org. Chem. **2005**, *70*, 7035–7041; (i) Boyd, E. A.; Chan, W. C.; Loh, V. M., Jr. Tetrahedron Lett. **1996**, *37*, 1647–1650.
- 11. Jiao, X.-Y.; Verbruggen, C.; Borloo, M.; Bollaert, W.; Groot, A. D.; Dommisse, R.; Haemers, A. Synthesis **1994**, 23–24.
- (a) Lewkowski, J. J. Organomet. Chem. 2003, 681, 225–227; (b) Lewkowski, J. Heteroat. Chem. 2004, 15, 162–168.
- 13. Tyka, R.; Hagele, G. Phosphorus, Sulfur, Silicon 1989, 44, 103-107.
- (a) Saigo, K.; Kubota, N.; Takebayashi, S.; Hasegawa, M. Bull. Chem. Soc. Jpn. 1986, 59, 931–932; (b) Corey, E. J.; Kuhnle, F. N. M. Tetrahedron Lett. 1997, 38, 8631–8634; (c) Larter, M. L.; Phillips, M.; Ortega, F.; Aguirre, G.; Somanathan, R.; Walsh, P. J. Tetrahedron Lett. 1998, 39, 4785–4788; (d) Lozinskaya, N. A.; Tsybezova, V. V.; Proskurnina, M. V.; Zefirov, N. S. Russ. Chem. Bull. 2003, 52, 674–678; (e) Nishiyama, K.; Saito, M.; Oba, M. Bull. Chem. Soc. Jpn. 1988, 61, 609–611; (f) Uchida, H.; Shimizu, T.; Reddy, P. Y.; Nakamura, S.; Toru, T. Synthesis 2003, 1236–1240; (g) Uchida, H.; Tanikoshi, H.; Nakamura, S.; Reddy, P. Y.; Toru, T. Synlett 2003, 1117–1120; (h) Kaboudin, B.; Saadati, F. Heterocycles 2005, 65, 353–357.
- (a) Corey, E. J.; Grogan, M. J. Org. Lett. **1999**, *1*, 157–160; (b) Isobe, T.; Fukuda, K.; Araki, Y.; Ishikawa, T. Chem. Commun. **2001**, 243–244; (c) Anastassiadou, M.; Baziard-Mouysset, G.; Payard, M. Synthesis **2000**, 1814–1816; (d) Corey, E. J.; Huang, H. C. Tetrahedron Lett. **1989**, *30*, 5235; (e) Corey, E. J.; Imwinkelried, R.; Pikul, S. B. J. Am. Chem. Soc. **1989**, *111*, 5493–5495; (f) Corey, E. J.; Kim, S. S. J. Am. Chem. Soc. **1990**, *112*, 4976–4977.
- (a) Kaboudin, B.; Moradi, K. Synthesis 2006, 2339–2342; (b) Kaboudin, B.; Jafari, E. Synthesis 2006, 3063–3066.
- (a) Kaboudin, B. Chem. Lett. 2001, 880-881; (b) Kaboudin, B.; Nazari, R. Tetrahedron Lett. 2001, 42, 8211; (c) Kaboudin, B.; Nazari, R. Synth. Commun. 2001, 31, 2245-2250; (d) Kaboudin, B. Tetrahedron Lett. 2002, 43, 8713-8714; (e) Kaboudin, B. Tetrahedron Lett. 2003, 44, 1051-1053; (f) Kaboudin, B.; Rahmani, A. Synthesis 2003, 2705–2708; (g) Kaboudin, B.; Saadati, F. Synthesis 2004, 1249-1252; (h) Kaboudin, B.; Rahmani, A. Org. Prep. Proced. Int 2004, 36, 82-86; (i) Kaboudin, B.; Moradi, K. Tetrahedron Lett. 2005, 46, 2989-2991; (j) Kaboudin, B.; Haghighat, H. Tetrahedron Lett. 2005, 46, 7955-7957; (k) Kaboudin, B.; Haghighat, H.; Yokomatsu, T. J. Org. Chem. 2006, 71, 6604-6606; (1) Kaboudin, B.; Karimi, M. Bioorg. Med. Chem. Lett. 2006, 16, 5324-5327; (m) Kaboudin, B.; Farjadian, F. Beilstein J. Org. Chem 2006, 2, 4; (n) Yamagishi, T.; Kusano, T.; Kaboudin, B.; Yokomatsu, T.; Sakuma, C.; Shibuya, S. Tetrahedron **2003**, 59, 767–772; (o) Kaboudin, B.; Haruki, T.; Yamaghishi, T.; Yokomatsu, T. Tetrahedron 2007, 63, 8199-8205; (p) Kaboudin, B.; Haruki, T.; Yamagishi, T.; Yokomatsu, T. Synthesis 2007, 3226-3232; (q) Kaboudin, B.; Haghighat, H.; Yokomatsu, T. Tetrahedron: Asymmetry **2008**, 19, 862–866.
- 18. Matziari, M.; Georgiadis, D.; Dive, V.; Yiotakis, A. Org. Lett. 2001, 3, 659-662.
- 19. The aldehyde (3 mmol) was added to ammonium hydroxide (30%, 15 mL), and the solution was stirred for 5 h at reflux. During this time, a white precipitate formed. The precipitate was removed by filtration and dried. The solid was dissolved in 5 mL of ethanol, and anhydrous hypophosphorus acid (5 mmol) was added to this mixture and the resulting solution was stirred for 2–12 h at reflux. The solvent was evaporated and the mixture was dissolved in acetone by heating. Dropwise addition of water gave the crude product as a white solid. The crude product was washed with ethanol and dried in air at room temperature to give product 2 in 40–71% yield. The solid product was washed with ethanol/water (50 ml, 9:1) and dried in air at room temperature to give a single diastereoisomer. All products gave satisfactory spectral data in accord with the assigned structures. Analytical and spectral data for compounds 2: *Compound* 2a: white solid, mixture of two diastereoisomers; ¹H NMR (D₂O/TMS-250 MHz): 3.79, 4.16 (2H, 2d, *J* = 13.5 Hz, -CHP), 6.83, 6.88 (2H, 2d, *J*_{HP} = 552 Hz), 6.85–7.40 (10H, m); ¹³C NMR (D₂O-NaOD/TMS-62.9 MHz): 60.0–

62.1 (m, CHP), 127.0, 127.5, 127.7, 128.5, 128.7, 128.9, 134.9, 136.5; ³¹P NMR (D₂O/H₃PO₄-101.2 MHz): 17.58, 17.85 (40:60) ppm; IR (KBr): 3650-2120 (-OH), 1250 (P=O), 1055–710 (P–O) cm⁻¹; Anal. Calcd for $C_{14}H_{17}NO_4P_2$: C, 51.68; H, 5.27; N, 4.31. Found: C, 51.55; H, 5.20; N, 4.56. Compound 2b: white solid, single diastereoisomer, mp: 240–242 °C; ¹H NMR (D₂O/TMS-250 MHz): 3.79 (2H, d, *J* = 14.0 Hz, –CHP), 6.84 (2H, d, *J*_{HP} = 552 Hz), 6.91 (4H, d, *J* = 7.2 Hz), 7.21 (4H, d, *J* = 7.2 Hz); ¹³C NMR (D₂O–NaOD/TMS-62.9 MHz): 59.2–61.5 (m, CHP), (A) 128.7, 130.1 (d, J_{CP} = 5.7 Hz), 132.8 (d, J_{CP} = 3.1 Hz), 133.4 (d, J_{CP} = 3.8 Hz); ³¹P $MR (D_20/H_3PO_4-101.2 MH2): 17.37 ppm; IR (KBr): 3650-2120 (-OH), 1250 (P=0), 1080-750 (P-0) cm^{-1}; Anal. Calcd for C_{14}H_{15}Cl_2NO_4P_2. C, 42.75; H,$ (r=0), 1030-r=0, 0147 ; Ania, carte di C₁4115(2),0472, c 4273, n, 3.85; N, 3.56. Found: C, 42.56; H, 3.80; N, 3.52. Compound **2c**: white solid, single diastereoisomer, mp: 230–232 °C; ¹H NMR (D₂0/TMS-250 MHz): 3.98 (2H, d, J = 13.5 Hz, -CHP), 6.99 (2H, d, $J_{HP} = 552$ Hz), 7.01–7.15 (m, 8H); ³¹P (P=0), 1180–580 (P-0) cm⁻¹; ¹³C NMR (D₂O-NaOD/TMS-62.9 MHz): 60.1 (dd, J_{CP} = 98.5 and 14.1 Hz), 115.4 (d, J_{CP}=21.4 Hz), 130.1–130.5 (m, Ar), 162.1 (d, J_{CP} = 243.2 Hz), Anal. Calcd for $C_{14}H_{15}F_2NO_4P_2$: C, 46.53; H, 4.19; N, 3.88. Found: C, 46.56; H, 4.12; N, 3.72. Compound **2d**: white solid, single diastereoisomer, mp: 212–214 °C; ¹H NMR (D₂O/TMS-250 MHz): 3.63 (s, 6H), 3.73 (2H, d, J = 14.5 Hz, -CHP), 6.86 (2H, d, J_{HP} = 550 Hz), 6.82 (4H, d, J = 8.0 Hz), 6.90 (4H, d, J = 8.0 Hz); ³¹P NMR (D₂O/H₃PO₄-101.2 MHz): 17.81 ppm; IR (KBr): 3650–2320 (–OH), 1248 (P=O), 1150–650 (P–O) cm⁻¹; ¹³C NMR (D₂O-NaOD/ TMS-62.9 MHz): 55.3, 59.2–61.5 (m, CHP), 114.8, 120.4, 130.1 (d, J_{CP} = 5.5 Hz), 159.7; Anal. Calcd for C₁₆H₂₁NO₆P₂: C, 49.86; H, 5.50; N, 3.64. Found: C, 49.71; H, 5.43; N, 3.52. Compound 2e: white solid, mixture of two diastereoisomers; ¹H NMR (D₂O/TMS-250 MHz): 2.07, 2.10 (s, 6H), 3.72, 4.12 (2H, 2d, J = 14.0 Hz, -CHP), 6.83, 6.88 (2H, 2d, J_{HP} = 551 Hz), 6.75–7.17 (8H, m); ³¹P NMR (D₂O/ $\begin{array}{c} H_3PO_4\text{-}101.2 \text{ MHz}); & 17.86, & 18.12 \text{ ppm}; \text{ IR (KBr)}; & 3650\text{-}2220 & (\text{-}OH), & 1251 \\ (P=0), & 1185\text{-}610 & (P-0) & \text{cm}^{-1}; & ^{13}\text{C} & \text{NMR} & (D_2O\text{-}NaOD/\text{TMS-}62.9 \text{ MHz}); & 20.1, \\ \end{array}$ 60.0–62.1 (m, CHP), 128.4 (d, J_{CP} = 5.7 Hz), 128.8 (d, J_{CP} = 5.7 Hz), 129.0, 129.2, 131.6, 133.3, 137.6 (d, J_{CP} = 3.1 Hz), 137.9 (d, J_{CP} = 3.1 Hz). Anal. Calcd for C16H21NO4P2: C, 54.38; H, 5.99; N, 3.96. Found: C, 54.30; H, 5.85; N, 4.05. Compound 2f: white solid, single diastereoisomer, mp: 226-228 °C; ¹H NMR $(D_2O/TMS-250 \text{ MHz})$: 4.15 (2H, d, J = 14.7 Hz, -CHP), 6.90 (2H, d, $J_{\text{HP}} = 557.5 \text{ Hz}$), 6.75–7.85 (m, 8H); ³¹P NMR ($D_2O/\text{H}_3\text{PO}_4$ -101.2 MHz): -CHP), 6.90 (2H, d, 15.95 ppm; IR (KBr): 3650-2120 (-OH), 1240 (P=O), 1180-580 (P-O) cm⁻ ¹³C NMR (D_2O -NaOD/TMS-62.9 MHz): 60.1 (dd, J_{CP} = 98.5 and 14.1 Hz), 116.1 $(d, J_{CP} = 21.4 \text{ Hz}), 125.3 - 130.5 \text{ (m, Ar)}, 161.1 \text{ (d, } J_{CP} = 242.0 \text{ Hz}), \text{ Anal. Calcd for}$ C14H15F2NO4P2: C, 46.53; H, 4.19; N, 3.88. Found: C, 46.50; H, 4.15; N, 3.75. Compound 2g: white solid, single diastereoisomer, mp: 212-214 °C; ¹H NMR (D₂O/TM225 white 30mic, single diadected and the probability, in *p* = 12 ± 21 + 0, in the temperature of the probability of 17.80 ppm; IR (KBr): 3650-2100 (-OH), 1248 (P=O), 1050-750 (P-O) cm⁻¹; ¹³C NMR (D₂O-NaOD/TMS-62.9 MHz): 61.2 (dd, J_{CP} = 97.5 and 13.8 Hz), 126.1-Calcd for $C_{22}H_{21}NO_4P_2$: C, 62.10; H, 4.98; N, 3.29. Found: C, 61.97; H, 5.03; N, 3 2 2

20. A mixture of phosphinic acid **2a** (one diastereoisomer, 1 mmol) and 1,1,1,3,3-hexamethyldisilazane (5 mmol, 1 mL) was heated at 110 °C for 2 h under Ar. The mixture was cooled to rt. Methyl acrylate (2 mmol, 0.18 mL) was added dropwise and the resulting mixture was stirred at 60 °C for 12 h. Absolute EtOH (10 mL) was added and the mixture was cooled to rt slowly. The solvent was removed under vacuum. The residue was treated with a solution of sodium bicarbonate (5%, 10 mL) and extracted with ethyl acetate (3 × 10 mL). The aqueous solution was acidified with HCl (5%), and extracted with CHCl₃ (4 × 10 mL). The combined organic layers were washed with brine, dried over Na₂SO₄ and evaporated under reduced pressure to give **3a** as a white solid in 85% yield. Mp: 95–96 °C; ¹H NMR (DMSO/TMS-250 MHz): 1.55–1.90 (m, 4H), 2.20–2.40 (m, 4H), 3.54 (s, 8H), 5.10–5.75 (br s, NH), 7.10–7.40 (m, 10H); ³¹P NMR (DMSO/H₃PO₄–101.2 MHz): 43.68 ppm; ¹³C NMR (DMSO/TMS-62.9 MHz): 21.8 (d, J_{CP} = 93.7 Hz), 26.6, 52.0, 59.9 (dd, J_{CP} = 99.8 and 13.3 Hz), 127.2–129.5 (m, Ar), 135.6, 172.9 (d, J_{CP} = 16.1 Hz); Anal. Calcd for C₂₂H₂₉NO₈P₂: C, 53.10; H, 5.89; N, 2.82. Found: C, 53.05; H, 5.80; N, 2.65.