

PII: S0040-4020(97)00448-1

Tandem Wolff Rearrangement-"tert-Amino Effect" Sequence: Synthesis of 2-Oxoindolinium Enolate Derivatives

Françoise Léost, Bernard Chantegrel and Christian Deshayes

Laboratoire de Chimie Organique, Département de Biochimie, Bâtiment 403, Institut National des Sciences Appliquées, 20 avenue Albert Einstein, 69621 Villeurbanne, FRANCE.

Abstract: The thermolysis of 1-diazo-2-oxo-(2-N,N)-disubstituted aminophenyl)ethylphosphonates 1 gave rise to 2-oxoindolinium enolate derivatives 4 through Wolff rearrangement and interaction of the *tert*-amino moiety with the ketene functionality. Variable amounts of either ammonium ylides 5 or of products resulting from their transformations were also formed during the course of the reaction. If the amino moiety was substituted by a benzyl or an allyl group, indolinones, resulting from [1,3] or [1,2] benzylic or allylic shifts, were isolated in place of compounds 4 and 5. \odot 1997 Elsevier Science Ltd.

INTRODUCTION

The formation of heterocycles by ring closure of *ortho*-substituted *N*,*N*-dialkylanilines (the "*tert*-amino effect") has been reviewed by Meth-Cohn and Suschitzky,^{1a} Verboom and Reindhoudt^{1b} and more recently by Meth-Cohn.^{1c} According to Meth-Cohn, five types of *tert*-amino effect processes are distinguished by the size of the ring formed or by its mode of formation in the key step of the reaction. The most frequently encountered examples are type 1 and 2 reactions which are two-step processes involving first hydrogen abstraction from the α position of the *tert*-amino function followed by cyclization of a dipolar intermediate to a five or six-membered ring and type 4 reactions with initial attack of the nitrogen atom on the unsaturated *ortho* substituent X=Y.



E-mail: deshayes@insa.insa-lyon.fr. Fax: 33 (0)4 72 43 88 96.

F. LÉOST et al.

The "tert-amino effect" has been exemplified by various "ortho 2π substituents" such as C=C, C=O, C=N, NO₂, N=N, N=SO₂. However, to the best of our knowledge, there was no study about the reaction of the tert-amino moiety with a ketene functionality as the X=Y substituent.²

Examples of intermolecular interaction between ketenes and tertiary amines are known to proceed through zwitterionic species which then rearrange.³⁻⁵ A related intramolecular transformation has been reported by the Rudler group.⁶ The reaction of aminocarbene chromium complexes IV with alkynes gives rise to intermediate enamino ketenes VI which afford nitrogen ylide chromium complexes VII resulting from a through-space interaction of the tertiary amine with the carbonyl group of the ketene function; however no metal-free ylides deriving from VII could be obtained (Scheme 1).



Scheme 1

Recent results from our laboratory have demonstrated that dimethylphosphono vinyl or aryl ketenes are easily generated by the thermal Wolff rearrangement of α -diazo- β -keto- γ , δ -alkenyl(or β -aryl)phosphonates.⁷ Consequently the thermolysis of 1-diazo-2-oxo-(2-N,N-disubstituted aminophenyl)ethylphosphonates 1 should afford intermediate 2-ketenyl-N,N-disubstituted anilines 3 and thus allow the study of the interaction between the tert-amino and ketene moieties (Scheme 2). In a preliminary communication,⁸ we have investigated the thermolysis of the β -aryl- α -diazo- β -ketophosphonates **1a**, b ortho-substituted by piperidine or pyrrolidine and shown that the attack of the nitrogen lone pair onto the ketene functionality led to the formation of mesoionic compounds **4a**,**b** (*tert*-amino effect process of type 4); during the course of the reaction, no product resulting from a hydrogen shift was observed but variable amounts of stable ammonium ylides 5 were formed (Table 1). The general pathway of the Wolff rearrangement involves initial loss of N_2 to form the keto carbene 2, which can react either by migration of the aryl group to afford the ketene 3, or by a variety of other processes.⁹ Thus the formation of the ylide 5, by interaction between the tert-amino group and the carbene moiety, appears as a competing process to the Wolff rearrangement. Most often ammonium ylides are intermediate species which can evolve, depending on their structures, either by Stevens [1,2] shift or [2,3] sigmatropic rearrrangement.¹⁰⁻¹³ Only a few examples of stable ammonium ylides are known, resulting from either intermolecular^{14,15} or intramolecular^{6,16} reactions. In particular the reaction of the 5-nitro-2-(piperidin-1-yl)benzoyl chloride with diazomethane was found to give a mixture of the expected diazoketone together with the 5-nitro-3-oxo-1spiropiperidinoindolinium enolate,¹⁶ resulting presumably from a similar amino-carbene interaction as for the analogous 5a.

As compared to the thermolysis of 2-dialkylaminobenzoyldiazophosphonates 1, a very similar chemistry has been observed in the thermolysis of 2-dialkylaminobenzoylazides VIII (Scheme 3). Martin, Meth-Cohn and Suschitzky¹⁶ and very recently Waldron and co-workers¹⁷ have reported that the decomposition of VIII produced a mixture of isocyanates IX resulting from the Curtius rearrangement and 1,1-dialkylindazol-1-ium-3-

olates X through direct interaction of the *tert*-amino moiety with a nitrene intermediate (or by a concerted mechanism from the azide), the ratios of the two products depending upon the nature of the R, R^1 and R^2 substituents. However no further evolution of the isocyanates IX to mesoionic compounds was observed.

The present report describes full results concerning the influence of the *tert*-amino moiety on the decomposition pathways followed during the thermal decomposition of the α -diazo- β -ketophosphonates 1.



Scheme 2



Scheme 3

RESULTS AND DISCUSSION

The ethyl N,N-disubstituted anthranilates 8a-b,d-i were prepared from ethyl 2-fluorobenzoate 6 by fluorine substitution with the requisite secondary amine in the presence of potassium carbonate,¹⁸ whereas compound 8c was obtained by reaction of ethyl anthranilate 7 with (Z)-2-butene-1,4-diol dimesylate¹⁹ (Scheme 4). Subsequent condensation of esters 8 with dimethyl lithiomethylphosphonate followed by diazotransfer with tosyl azide yielded the α -diazo- β -ketophosphonates 1a-i.



a) R1R2NH, K2CO3; b) (Z)-MsOCH2CH=CHCH2OMs, C6H6; c) LiCH2P(O)(OMe)2, THF; d) TsN3, K2CO3, CH3CN.

Scheme 4

As reported in our preliminary communication,⁸ the thermolysis of 1a in refluxing toluene gave rise to a 49:51-mixture of 4a and 5a whose structures were established by X-ray crystallography.²⁰ The structure of the oxoindolinium enolate 4a²¹ deserves comment. Examination of the four N(1)-C bonds showed normal lenghts of 1,508 Å for N(1)-C(8) and N(1)-C(12) (C_{sp} 3-N⁺ = 1,510 Å)²² and of 1,458 Å for N(1)-C(7a) (C_{ar} -N⁺ = 1,465 Å) and a very long N(1)-C(2) bond of 1.645 Å. This remarkable value together with a rather short C(2)-C(3) bond of 1.382 Å and an O(1)-C(2)-C(3) bond angle of 140.0° suggested a significant contribution of the open valence tautomer A as a canonical form to the structure 4a (Scheme 5). An other important contributor to the overall structure must be the cyclic valence tautomer B in relation with the C(2)-O(1) bond of 1,210 Å (typical C=O double bond; 1.20 Å) coupled with the very strong infrared carbonyl absorption at 1750 cm⁻¹. Previous to this work, the same structural analysis has been reported first by Thiessen and Hope for the molecular geometry and bonding in the sydnone ring $(1,2,3-\text{oxadiazol-}5-\text{ones})^{23}$ and then by other groups for isoxazolium enolates²⁴ and imidazo[1,2-c]pyrimidin-3-ones.²⁵ It appeared that the characteristic bond lenghts and bond angles in 4a compared well with the literature values for the related spiro mesoionic compounds XI6a and XII24a (Table 2) but, except for the C(2)-O(1) bond, are quite different from those reported for the mesoionic compound XIII.²⁵ Thus the N(1)-C(2) bond length in XIII (nearly that of a C-N single bond at 1.47 Å) is shorter than in 4a, XI and XII and this is presumably due to the delocalisation of the positive charge to the nitrogen atoms N(1) and N(4) in XIII which may stabilize and thus favour the cyclic valence tautomers with respect to the open valence tautomer. Such a stabilization is impossible for 4a, XI and XII, the positive charge being located on the tetrasubstituted nitrogen atom. This analysis is confirmed by a similar comparison between the C(2)-C(3) bond

lengths and N(1)-C(2)-O(1) and O(1)-C(2)-C(3) bond angles in 4a, XI, XII and XIII. As previously observed by the Rudler^{6a} and Edstrom²⁵ groups, the weakness of the N(1)-C(2) bond is illustrated by the facile acidcatalysed ring opening of 4a by water to give the acid 10 which decarboxylated immediately to yield methylenephosphonate 11 (Scheme 6).

The ¹³C-NMR chemical shifts for the mesoionic compound **4a** with C-2 at δ 170.2 and C-3 at δ 62.3 are close to those reported by Rudler^{6a} for XI, respectively at δ 169.1 and δ 67.3, but very different from those for the ammonium ylide **5a**, with C-2 at δ 105.8 (¹J_{CP} = 226.8 Hz) and C-3 at δ 172.1 (²J_{CP} = 16.8 Hz). Compounds **4a** and **5a** are also easily differentiated by their infrared carbonyl absorption, respectively at 1750 cm⁻¹ and 1620 cm⁻¹. These characteristic spectral parameters were used in the sequel to support the structures of the other mesoionic compounds **4** (Table 3) and ylides **5**.

Starting Compounds	Ratios of Products	Total yield (%)
la	4a:5a = 49:51	87
1b	4b:5b = 98:2	88
1 c	4c:5c = 100:0	66
1 d	4d:5d = 94:6	72
1 e	4e:(5e+12) = 58:42	91
1 f	4f:(13+14) = 51:49	77
18	19:20 = 87:13	60
1 g	4g:21 = 90:10	74
1 h	22h:23h = 53:47	82
1i	22i:23i = 73:27	96

Table (1): Thermal Decomposition of the Diazophosphonates 1a-i and 18

Table (2):	Characteristic I	Bond Lenghts	[Å] and Angle	Values [°]	l in	some M	lesoionic	Compounds*
---------	-----	------------------	--------------	----	-------------	------------	------	--------	-----------	------------

Bonds or angles	$ \begin{array}{c} $	Ph-Cr(CO) ₃ Ph 3 + O 1 X1 (ref 6a)	XII (ref 24a)	Bn C1 + 1 N + 1 N + N +
N(1)-C(2)	1.645	1.590	1.545	1.457
C(2)-C(3)	1.382	1.368	1.387	1.408
C(2)-O(1)	1.210	1.225	1.218	1.219
N(1)-C(2)-O(1)	114.8	115.7	116.0	121.4
O(1)-C(2)-C(3)	140.0	138.6	137,7	135.3
N(1)-C(2)-C(3)	105.1	105.7	106,4	-

* The atoms are numbered for an easy comparison between the different compounds.



The replacement of the piperidino substituent by a pyrrolidino substituent resulted in the formation of the mesoionic compound **4b** together with a minute amount of the ylide **5b**. Similar results were observed when an insaturation was introduced in the five membered ring (pyrrolino: c) or in the case of a saturated seven membered ring (perhydroazepino: d) (Table 1).

The decomposition of the diazo 1e, bearing a morpholino substituent, yielded a 58:13:29-mixture of 4e, 5e and oxazepinoindolinone 12 (Scheme 7). Refluxing the ylide 5e in toluene for 24 h afforded quite quantitatively the indolinone 12 which should therefore be the result of a Stevens [1,2] shift. It has been shown that the Stevens rearrangement normally involves a radical pair mechanism.^{26,27} Thus the facile formation of the oxazepinoindolinone 12 from 5e can be rationalized by the intermediate formation of the stabilized biradical 5'e in which the cyclization either to the α - or β -position gives rise to the same product. The decomposition of the diazo 1f, bearing a tetrahydropyridino substituent, afforded a 51:49-mixture of mesoionic compound 4f and tricyclic compounds (13+14) resulting from rearrangement of the ylide 5f which, in this case, was not isolated. In a similar manner as for 12, the formation of 13 and 14 can be explained by the intermediate formation of the stabilized biradical 5'f, in which radical recombination either to the α - or β -position leads respectively to 13 or 14. Pure samples of tricyclic compounds 12 and 13 were isolated and their structures were characterized by their ¹³C-NMR spectra showing resonances for a ketonic carbonyl respectively at δ 197.6 or 197.9 and for a carbon atom bonded to the phosphorus atom respectively at δ 72.8 or 75.3 (¹J_{CP} = 154.4 or 152.7 Hz); furthermore the resonances of the two methoxy groups were splitted in two doublets, due to the 2-bonds C-P coupling and to the presence of an asymetric carbon in these structures. We were not able to isolate a pure sample of 14; its structure was inferred from the presence of characteristic vinylic protons resonances at δ 6.15 (m, 1H) and δ 5.55 (m, 2H) in the ¹H-NMR spectrum of a mixture constituted by 13 and 14.

Compound	C-2	C-3	C-3a	C-5, C-6 ,	C-7a	R ₁ , R ₂	PO(OMe)2
(vCO)				C-4** , C-7			
4a	170.2	62.3	139.3	130.1, 120.0,	137.7	55.8, 21.0,	52.3
1750	$^{2}J_{CP} = 24.8$	${}^{1}J_{CP} = 219.3$	$^{2}J_{CP} = 15.0$	119.9, 118.7	${}^{3}J_{CP} = 13.0$	20.3	${}^{2}J_{CP} = 5.3$
4b	170.5	61.2	139.5	130.1, 120.8,	140.4	61.6, 25.1	52.2
1735	$2_{J_{CP}} = 25.0$	${}^{1}JCP = 220.2$	${}^{2}J_{CP} = 15.0$	119.4, 116.0	${}^{3}J_{CP} = 13.2$		${}^{2}J_{CP} = 5.4$
4c	169.6	63.4	138.9	130.3, 120.9,	141.6	125.0, 67.9	52.0
1750	${}^{2}J_{CP} = 25.6$	${}^{1}J_{CP} = 219.9$	${}^{2}J_{CP} = 14.9$	119.1, 114.8	${}^{3}J_{CP} = 13.3$		${}^{2}J_{CP} = 5.3$
4d	173.0	61.4	138.0	130.1, 120.6,	139.8	61.5, 29.7,	52.2
1750	${}^{2}J_{CP} = 24.7$	${}^{1}J_{CP} = 219.7$	${}^{2}J_{CP} = 15.0$	119.6, 117.0	${}^{3}J_{CP} = 13.4$	23.5	${}^{2}J_{CP} = 5.3$
4e	170.7	61.2	139.1	130.6, 120.6,	137.4	61.1, 54.7	52.4
1760	${}^{2}J_{CP} = 25.6$	${}^{1}J_{CP} = 219.4$	$^{2}J_{CP} = 14.8$	120.2, 117.9	${}^{3}J_{CP} = 12.9$		${}^{2}J_{CP} = 5.4$
4f	169.8	62.4	139.1	130.4, 120.6,	137.1	124.8, 122.1,	52.29 (d) and
1750	${}^{2}J_{CP} = 25.2$	${}^{1}J_{CP} = 219.5$	${}^{2}J_{CP} = 14.8$	119.7, 117.9	${}^{3}J_{CP} = 13.0$	52.6, 52.4,	52.25 (d)
						21.2	${}^{2}J_{CP} = 5.3$
4g	165.5	68.3	142.1	130.5, 120.9,	131.1	55.2, 8.1	52.0
1735	$^{2}J_{CP} = 24.8$	$1_{J_{CP}} = 217.3$	$J_{CP} = 15.1$	119.3, 116.7	${}^{3}J_{CP} = 13.3$		${}^{2}J_{CP} = 5.1$
19	167.8	62.2	133.7	126.8, 138.6,	151.4	55.1, 21.6,	52.3
1740	${}^{2}J_{CP} = 22.7$	$^{1}J_{CP} = 219.3$	$J_{CP} = 14.5$	125.6	${}^{3}J_{CP} = 14.1$	19.9	${}^{2}J_{CP} = 5.4$

Table (3): v_{CO} (cm⁻¹) and ¹³C-NMR Data* of Mesoionic Compounds 4 and 19 [δ (ppm) and J (Hz)]

*The ¹³C-NMR chemical shifts and coupling constants of compound 4a were attributed on the basis of H,H-COSY, H,C-COSY and H,C-Heteronuclear Multi Bond Correlation. Chemical shifts and coupling constants of other compounds 4 were deduced on the basis of their spectral similarities with 4a. ** No 3-bonds coupling was observed between P and C(4).



F. LÉOST et al.

It appeared that the ring size of the amino functionality strongly influenced the decomposition towards the Wolff rearrangement path or the ylide path, but that the presence of an insaturation or of an heteroatom in the ring had a minor influence. Thus five or seven membered rings (b,c,d) gave rise to mixtures containing mainly compounds 4 resulting from the Wolff rearrangement whereas six membered rings (a,e,f) afforded mixtures containing nearly equal amounts of 4 and products resulting from the ylide path (Table 1). The Wolff rearrangement as well as the ylide formation²⁸ are irreversible processes and therefore both compounds 4 and 5 must be considered as kinetic products. Hence the orientation of the thermolysis must be related to the competition between the Wolff rearrangement and the attack of the nitrogen atom lone pair onto the carbene moiety. A similar analysis has been reported by Waldron and Raza^{17b} to explain the results in the thermal decomposition of 2-dialkylaminobenzoylazides VIII^{16,17} (Scheme 3). The orientation of the reaction resulted from a competition between the Curtius rearrangement leading to the isocyanate IX and the cyclisation to the indazolium olate X, arising probably from the trapping of the intermediate nitrene by the lone pair of electrons on the nearby amino group. The availability and the orientation of the lone pair on the nitrogen were determined to be the key factors and the (IX/X) ratios were interpreted by electronic and steric effects of the R, R¹ and R² substituents. Thus in the decomposition of compounds VIIIa ($R^1 = R^2 = H, R = Me$) and VIIIb ($R^1 = R^2 = H$, R = Et), the (IXa/Xa) and (IXb/Xb) ratios were found to be respectively 56/31 and 9/79; the prevailing orientation directed towards the zwitterionic product X in the case of the ethyl groups was explained by a greater electron availability on the nitrogen atom and a greater steric hindrance. Unfortunately in our case, we found no evident steric, electronic or stability factors in relation with the ring size of the amino functionality to rationalize the observed products ratios.

Although the present work was centered on the nature of the amino moiety, we investigated the effect of the replacement of the benzene ring by a pyridine ring. Starting from ethyl 2-chloronicotinate 15, we prepared the diazophosphonate 18 in a similar manner as described above (Scheme 8); its thermolysis afforded a 87:13-mixture of mesoionic compound 19 and ammonium ylide 20, showing that the orientation of the reaction was dependent on the aromatic moiety. As compared with the decomposition of 1a, this result could be related, for electronic and steric reasons, to an unfavorable orientation of the nitrogen atom lone pair lowering its interaction with the carbene moiety and thus favoring the Wolff rearrangement product 19. A possible influence of the pyridine ring on the rate of Wolff rearrangement could be also considered. In conclusion, we think that the decomposition paths of the diazophosphonates 1a-f or 18 are governed by fine parameters which are difficult to evidence in view of our present results.

We next turned our attention to the thermolysis of diazophosphonates 1 bearing an acyclic *tert*-amino moiety. The diazophosphonate 1g possessing two ethyl groups as the amino substituents yielded the mesoionic compound 4g as the major product together with a small amount of the hydroxyindole 21 formed by Hofmann elimination from the non-isolated ylide 5g (Scheme 9). The structure of the hydroxyindole 21 was supported by the presence of an OH stretching band at 3260 cm⁻¹ and by the resonances at δ 100.1 (${}^{1}J_{CP}$ = 218.6 Hz) for C-2 and δ 151.6 (${}^{2}J_{CP}$ = 15.8 Hz) for C-3.



a) Piperidine, K₂CO₃; b) LiCH₂P(O)(OMe)₂, THF; c) TsN₃, K₂CO₃, CH₃CN.

Scheme 8





Neither mesoionic compound 4 nor ammonium ylide 5 were isolated in the case of diazophosphonates 1 bearing an alkyl group of strong migratory aptitude on the nitrogen atom. Thus the thermolysis of compounds 1 possessing an allyl group (h) or a benzyl group (i) as the R₂ amino substituent gave rise to mixtures of indolinones 22 and 23 (Scheme 10). The 2-indolinones 22 were formed through the corresponding mesoionic compounds 4 by a [1,3] shift of the allyl or benzyl group. It should be noticed that no [1,3] shift had occured in the case of cyclic *tert*-amino moieties (a-f), even when an allyl system was present in the ring (c,f); these results are different from the [1,3] nitrogen-to-carbon migrations of various alkyl groups observed by Rudler^{6a,b} when ylide complexes VII, XI or related compounds were refluxed in toluene.²⁹ The formation of 3-indolinones 23 resulted from a Stevens rearrangement of the ylides 5h,i which showed a similar behaviour as the ylides 5e,f. The lactam structures of 22h,i were in agreement with resonances at δ 173.0-172.8 (²J_{CP} = 2.8-2.7 Hz) for C-2 and at δ 54.7-56.3 (¹J_{CP} = 148.4-142.6 Hz) for C-2 and at δ 196.2-196.1 (²J_{CP} = 2.4-3.0 Hz) for C-3. The resonances of the two methoxy groups in 22 or 23 were splitted in two doublets, due to the 2-bonds C-P coupling and to the presence of an asymetric carbon, respectively C-3 or C-2.



We have also synthesized the diazophosphonate of type 1 bearing a methyl group and a phenyl group as the amino substituents. Its decomposition gave a complex mixture of unidentified products and therefore these experiments are not included in the experimental section. As compared with dialkyl substituents, this negative result can be explained by the lowering of the nitrogen nucleophilicity, resulting from the introduction of the phenyl substituent, which prevents the interaction of nitrogen with the intermediate carbene or ketene moities to take place.

CONCLUSION

The nature of the products obtained by thermal decomposition of 1-diazo-2-oxo-(2-N,N-disubstituted aminophenyl)ethylphosphonates 1 depends strongly on the structure of the *tert*-amino substituent. The reaction allows the synthesis of 2-oxoindolinium enolate derivatives 4 if the amino moiety is part of a ring or bears substituents of low migratory aptitude. Ammonium ylides 5 or products resulting from their transformations are also formed during the course of the decompositions, especially when the amino moiety is part of a 6-membered ring. If the amino moiety is acyclic and bears a substituent of strong migratory aptitude, indolinones 22 or 23, resulting from [1,3] or [1,2] benzylic or allylic shifts, were directly obtained in place of compounds 4 and 5.

EXPERIMENTAL

General.

Diethyl ether was distilled from potassium hydroxide, pentane from phosphorus pentoxide, tetrahydrofuran from sodium benzophenone ketyl. Benzene and toluene were dried over sodium. Organic solutions were dried over anhydrous sodium sulfate. Column chromatographies were performed using Merck Silica gel 60 (70-230 mesh) and TLC were carried out using Merck Kieselgel 60 F254 plates. Melting points were determined on a Kofler block apparatus. IR spectra were recorded on a Perkin Elmer 1310 infrared spectrophotometer. Nuclear magnetic resonance spectra were recorded in CDCl₃ on a Brucker AC200 (200/50 MHz) spectrometer. All NMR recordings were referenced to CHCl₃ resonances (7.26 and 77.0 ppm). Splitting patterns abbreviations are: s, singulet; d, doublet; t, triplet; m, multiplet; b, broad; p, pseudo. Multiplicity (¹³C NMR) was determined by DEPT sequencies. Elemental analyses were performed by Service Central d'Analyse, Centre National de la Recherche Scientifique, 69300 Vernaison, France.

SYNTHESIS OF AMINOESTERS (8a-i) and (16)

The ethyl 2-fluorobenzoate 6 was obtained by reaction of 2-fluorobenzoyl chloride with absolute ethanol in refluxing ethanol (yield: 91 %). The ethyl 2-chloronicotinate 15 was obtained by reaction of 2-chloronicotinic acid with thionyl chloride in refluxing benzene followed by treatment of the intermediate 2-chloronicotinoyl chloride with absolute ethanol at room temperature (yield: 95 %).

General Procedure for the Preparation of Aminoesters 8a,b,d,e,f,h,i and 16: a mixture of 2-fluorobenzoate 6 (3 g, 17.8 mmol) or ethyl 2-chloronicotinate 15 (3.3 g, 17.8 mmol), potassium carbonate (2.9 g, 21.4 mmol) and of the requisite secondary amine (21.4 mmol) in 10 ml of DMF (a,b,d,e,f,i) or toluene (h) was refluxed for 24 h. After cooling, the resulting mixture was poured into water (120 ml) and extracted with ethyl acetate (3 x 30 ml). The organic layers were dried, filtered and concentrated to afford a residue which was purified by chromatography eluting with pentane-ethyl acetate 10:90 to afford 8a,b,d,e,f,h,i and 16.

Ethyl 2-piperidinobenzoate (8a)

Yield 3.45 g (83 %). Oil. IR (CHCl₃): 1720, 1595. ¹H-NMR: δ 7.58 (dd, 1H, J = 7.7, 1.8); 7.35 (ddd, 1H, J = 8.3, J = 7.3, J = 1.8); 7.00 (dd, 1H, J = 8.3, J = 1.1); 6.92 (td, 1H, J = 7.5, J = 1.1); 4.31 (q, 2H, J = 7.1); 2.97 (t, 4H, J = 5.2); 1.74-1.43 (m, 6H); 1.36 (t, 3H, J = 7.1). ¹³C-NMR: δ 168.5; 153.1; 132.2; 131.1; 124.7; 120.7; 118.8; 60.8; 53.8; 26.2; 24.3; 14.3. Anal Calcd for C₁₄H₁₉NO₂: C, 72.07; H, 8.21; N, 6.00. Found: C, 71.85; H, 8.36; N, 5.96.

Ethyl 2-pyrrolidinobenzoate (8b)

Yield 2.77 g (71 %). Oil. IR (CHCl₃): 1725, 1595. ¹H-NMR : δ 7.57 (dd, 1H, J = 7.8); 7.30 (ddd, 1H, J = 8.2, J = 7.2, J = 1.7); 6.77 (d, 1H, J = 8.2); 6.70 (pdd, 1H, J = 7.7, J = 7.2); 4.35 (q, 2H, J = 7.1); 3.27-3.20 (m, 4H); 1.97-1.90 (m, 4H); 1.37 (t, 3H, J = 7.1). ¹³C-NMR: δ 169.1; 147.9; 131.7; 131.0; 117.5; 115.6; 114.0; 60.7; 50.8; 25.9; 14.4. Anal Calcd for C₁₃H₁₇NO₂: C, 71.21; H, 7.81; N, 6.39. Found: C, 71.19; H, 7.54; N, 6.25.

Ethyl 2-perhydroazepinobenzoate (8d)

Yield 3.21 g (73 %). Oil. IR (neat): 1705, 1595. ¹H-NMR: δ 7.52 (dd, 1H, J = 7.7, J = 1.8); 7.28 (ddd, 1H, J = 7.7, J = 8.1, J = 1.8); 6.95 (d, 1H, J = 8,1); 6.73 (pt, 1H, J = 7.7); 4.33 (q, 2H, J = 7.1); 3.33 (m, 4H); 1.76 (m, 4H); 1.59 (m, 4H); 1.45 (t, 3H, J = 7.1). ¹³C-NMR: δ 169.4; 151.2; 131.5; 130.9; 120.4; 117.1 (2C); 60.7; 52.9; 28.5; 28.1; 14.3. Anal Calcd for C₁₅H₂₁NO₂: C, 72.84; H, 8.56; N, 5.66. Found: C, 72.89; H, 8.47; N, 5.61.

Ethyl 2-morpholinobenzoate (8e)

Yield 1.60 g (41 %). Oil. IR (CHCl₃): 1720, 1595. ¹H-NMR: δ 7.75 (dd, 1H, J = 7.9, J = 1.9); 7.42 (m, 1H); 7.04 (d, 1H, J = 7.8); 7.02 (m, 1H); 4.35 (q, 2H, J = 7.1); 3.91-3.80 (m, 4H); 3.10-3.00 (m, 4H); 1.39 (t, 3H, J = 7.1). ¹³C-NMR : δ 167.6; 151.9; 132.3; 131.2; 124.8; 121.6; 118.7; 66.9; 60.7; 52.6; 14.1. Anal Calcd for C₁₃H₁₇NO₃: C, 66.36; H, 7.28; N, 5.95. Found: C, 66.16; H, 7.68; N, 5.92.

Ethyl 2-(2-(1,2,3,6-tetrahydropyridino)benzoate (8f)

Yield 3.00 g (73 %). Oil. IR (CHCl₃): 1715, 1595. ¹H-NMR; δ 7.68 (dd, 1H, J = 7.7, J = 1.6); 7.37 (t, 1H, J = 7.7); 7.02 (d, 1H, J = 8.3); 6.96 (ddd, 1H, J = 8.3, J = 7.7, J = 1.6); 5.87 (dm, 1H, J = 10.0); 5.76 (dm, 1H, J = 10.0); 4.35 (q, 2H, J = 7.1); 3.61 (m, 2H); 3.21 (t, 2H, J = 5.5); 2.30 (tm, 2H, J = 5.5); 1.37 (t, 3H, J = 7.1). ¹³C-NMR: δ 168.4; 151.9; 132.3; 131.4; 125.8; 125.3; 123.6; 120.3; 118.1; 60.8; 50.5; 50.4; 26.1; 14.3. No correct microanalysis could be obtained for this compound.

Ethyl 2-(N-allyl, N-methylamino)benzoate (8h)

Yield 3.04 g (78 %). Oil. IR (neat): 1710, 1595. ¹H-NMR: δ 7.64 (dd, 1H, J = 7.7, J = 1.7); 7.34 (ddd, J = 8.3, J = 7.4, J = 1.7); 6.98 (d, 1H, J = 8.3); 6.87 (td, 1H, J = 7.4, J = 0.9); 5.93 (ddt, 1H, J = 17.1, J = 10.2, J = 5.9); 5.30-5.10 (m, 2H); 4.37 (q, 2H, J = 7.1); 3.71 (d, 2H, J = 5.9); 2.80 (s, 3H); 1.39 (t, 3H, J = 7.1). ¹³C-NMR : δ 168.5; 151.6; 134.8; 131.9; 131.2; 122.5; 119.3; 118.2; 117.3; 60.8; 59.4; 39.9; 14.3. Anal Calcd for C₁₃H₁₇NO₂: C, 71.21; H, 7.81; N, 6.39. Found: C, 71.38; H, 7.80; N, 6.53.

Ethyl 2-(N-benzyl, N-methylamino)benzoate (8i)

Yield 2.97 g (62 %). Oil. IR (CHCl₃): 1725, 1595. ¹H-NMR: δ 7.65 (dd, 1H, J = 7.7, J = 1.7); 7.36-7.21 (m, 6H); 6.97 (dd, 1H, J = 8.4, J = 1.0); 6.88 (td, 1H, J = 7.4, J = 1.0); 4.32 (q, 2H, J = 7.1); 4.31 (s, 2H); 2.74 (s, 3H); 1.34 (t, 3H, J = 7.1). ¹³C-NMR: δ 168.7; 151.7; 138.2; 131.9; 131.2; 128.3; 127.9; 127.0; 123.1; 119.6; 118.7; 60.9; 59.8; 40.9; 14.3. Anal Calcd for C₁₇H₁₉NO₂: C, 75.81; H, 7.11; N, 5.20. Found: C, 75.61; H, 7.21; N, 5.15.

Ethyl 2-piperidinonicotinate (16)

Yield 3.25 g (78 %). Oil. IR (CHCl₃): 1705, 1580. ¹H-NMR: δ 8.25 (dd, 1H, J = 4.7, J = 2.0); 7.92 (dd, 1H, J = 7.6, J = 2.0); 6.67 (dd, 1H, J = 7.6, J = 4.7); 4.35 (q, 2H, J = 7.1); 3.36 (br s, 4H); 1.65 (br s, 6H); 1.38 (t, 3H, J = 7.1). ¹³C-NMR: δ 167.7; 159.6; 150.3; 140.4; 113.9; 113.2; 60.9; 50.4; 25.9; 24.6; 14.3. Anal Calcd for C₁₃H₁₈N₂O₂: C, 66.64; H, 7.74; N, 11.96. Found: C, 66.77; H, 7.72; N, 11.72.

Procedure for the Preparation of Aminoester 8c: the ethyl anthranilate 7 (2.7 g, 16.3 mmol) was added slowly to a mixture of (Z)-2-butene-1,4-diol dimesylate (8 g, 32.7 mmol) and ethyldiisopropylamine (4.2 g, 32.7 mmol) in benzene (160 ml). After refluxing for 24 h, the benzene was evaporated in vacuum. To the residue was added water (50 ml) and then a saturated aqueous ammoniac solution up to pH 10. After extraction with ethyl acetate (3 x 50 ml), the organic layers were dried, filtered and concentrated to afford a residue which was purified by chromatography eluting with pentane-ethyl ether 90:10.

Ethyl 2-pyrrolinobenzoate (8c)

Yield 2.97 g (84 %). Mp 41-43°C. IR (CHCl₃): 1700, 1600. ¹H-NMR : δ 7.51 (dd, 1H, J = 7.5, J = 1.7); 7.31 (dt, 1H, J = 7.5, J = 1.8); 6.73 (m, 2H); 5.88 (s, 2H); 4.36 (q, 2H, J = 7.1); 4.12 (s, 4H); 1.39 (t, 3H, J = 7.1). ¹³C-NMR: δ 169.6; 146.4; 131.6; 130.8; 125.7; 117.4; 115.6; 114.2; 60.9; 56.8; 14.3. Anal Calcd for C₁₃H₁₅NO₂: C, 71.87; H, 6.96; N, 6.45. Found: C, 71.76; H, 6.97; N, 6.54.

Procedure for the Preparation of Aminoester 8g: a mixture of 2-fluorobenzoate 6 (17.8 mmol), potassium carbonate (2.9 g, 21.4 mmol) and diethylamine (1.56 g, 21.4 mmol) was introduced into a high-pressure autoclave and heated at 145 °C for 24 h. After cooling the residue was poured into water (120 ml) and extracted with ethyl acetate (3 x 30 ml). The organic layers were dried, filtered and concentrated to afford a residue which was purified by chromatography eluting with pentane-ethyl acetate 10:90 to afford 8g.

Ethyl 2-(N,N-diethylamino)benzoate (8g)

Yield 2.28 g (58 %). Oil. IR (CHCl₃): 1700, 1590. ¹H-NMR: δ 7.55 (dd, 1H, J = 7.7, J = 1.7); 7.34 (ddd, 1H, J = 8.3, J = 7.3, J = 7.3, J = 1.7); 7.05 (dd, 1H, J = 8.3, J = 0.9; 6.93 (ptd, 1H, J = 7.6, 0.9); 4.34 (q, 2H, J = 7.1); 3.15 (q, 4H, J = 7.1); 1.37 (t, 2H, J = 7.1); 1.05 (t, 4H, J = 7.1). ¹³C-NMR : δ 168.9; 150.3; 131.3; 130.4; 127.0; 121.2; 120.7; 60.7; 47.0; 14.3; 12.4. Anal Calcd for C₁₃H₁₉NO₂: C, 70.56; H, 8.65; N, 6.33. Found: C, 70.10; H, 8.53; N, 6.18.

SYNTHESIS OF β -KETOPHOSPHONATES (9a-i) and (17)

General Procedure: to a stirred solution of dimethyl methylphosphonate (3.5 g, 28.2 mmol) in anhydrous THF (70 ml) cooled at -80°C, was added dropwise, under nitrogen, 17.6 ml (28.2 mmol) of *n*-butyllithium 1.6 M in hexanes. The mixture was kept for 45 mn at -60 °C and then a solution of amino ester 8a-i or 16 (14.1 mmol) in anhydrous THF (30 ml) was added slowly. The reaction mixture was allowed to react at room temperature for 3.5 h, and was then quenched with a saturated ammonium chloride solution (200 ml). The aqueous phase was extracted with ethyl acetate (5 x 60 ml), the organic layers were washed with brine, dried, filtered and concentrated to afford a residue which was purified by chromatography eluting with ethyl acetate. The samples obtained after this purification contained variable amounts of residual dimethyl methylphosphonate and were characterized by their IR spectra [v_{CO} : 1680-1650 cm⁻¹] and by their ¹H-NMR spectra [CH₂P δ 4.04-3.68 ($^{2}_{JHP} = 22.7-21.6$) and PO(OMe)₂: δ 3.79-3.52 ($^{3}_{JHP} = 11.2$)].

SYNTHESIS of α -DIAZO- β -KETOPHOSPHONATES (1a-i) and (18)

General Procedure: to a mixture of β -ketophosphonate 9a-i or 17 (10 mmol), obtained as described above, and potassium carbonate (1.72 g, 12.5 mmol) in acetonitrile (40 ml) cooled in a water-ice bath, under nitrogen, was added dropwise with stirring a solution of tosyl azide^{30,31} (2.46 g, 12.5 mmol) in acetonitrile (30 ml). The cooling bath was removed and the mixture was stirred at room temperature. The disappearance of compound 9 or 17 was monitored by tlc. After 1-2.5 h, potassium carbonate was filtered off and acetonitrile was evaporated *in vacuo* to afford a residue which was purified by chromatography eluting with ethyl acetate (1a,b,g,i) or ethyl acetate:pentane 70:30 (1c-f,h and 18).

Dimethyl 1-diazo-2-oxo-2-(2-piperidinophenyl)ethylphosphonate (1a)

Yield 2.73 g (81 %). Mp 93-94°C. IR (CHCl₃): 2125, 1625, 1265, 1055, 1030. ¹H-NMR: δ 7.32 (td, 1H, J = 7.8, J = 1.7); 7.23 (dd, 1H, J = 7.8, J = 1.7); 7.10-7.02 (m, 2H); 3.86 (d, 6H, ³J_{HP} = 11.9); 2.96 (t, 4H, J = 5.2); 1.78-1.68 (m, 4H); 1.60-1.52 (m, 2H). ¹³C-NMR: δ 189.8 (d, ²J_{CP} = 7.0); 132.2; 131.8 (d, ³J_{CP} = 5.9); 129.5; 122.6; 118.4; 63.3 (d, ¹J_{CP} = 224.1); 54.3; 54.2 (d, ²J_{CP} = 6.3); 26.4; 23.9. Anal Calcd for C₁₅H₂₀N₃O₄P: C, 53.41; H, 5.98; N, 12.46; P, 9.18. Found: C, 53.17; H, 5.95; N, 12.48; P, 8.65.

Dimethyl 1-diazo-2-oxo-2-(2-pyrrolidinophenyl)ethylphosphonate (1b)

Yield 2.17 g (67 %). Mp 95-96°C. IR (CHCl₃): 2110, 1620, 1250, 1050, 1030. ¹H-NMR: δ 7.34-7.24 (m, 2H); 6.79-6.70 (m, 2H); 3.84 (d, 6H, ³J_{HP} = 11.9); 3.25-3.18 (m, 4H); 1.98-1.91 (m, 4H). ¹³C-NMR: δ 189.1 (d, ²J_{CP} = 8.2); 146.9; 132.0; 129.1; 123.4 (d, ³J_{CP} = 4.7); 116.5; 114.4; 63.8 (d, ¹J_{CP} = 220.7); 54.1 (d, ²J_{CP} = 6.2); 51.0; 25.8. Anal Calcd for C₁₄H₁₈N₃O₄P: C, 52.01; H, 5.61; N, 13.00; P, 9.58. Found: C, 51.78; H, 5.43; N, 13.00; P, 9.36.

Dimethyl 1-diazo-2-oxo-2-(2-pyrrolinophenyl)ethylphosphonate (1c)

Yield 2.76 g (86 %). Mp Oil. IR (CHCl₃): 2110, 1620, 1250, 1060, 1030. ¹H-NMR: δ 7.40-7.20 (m, 2H); 6.80-6.70 (m, 2H); 5.90 (s, 2H); 4.11 (s,4H); 3.85 (d, 6H, ³J_{HP} = 11.8). ¹³C-NMR: δ 189.1 (d, ²J_{CP} = 8.8); 145.4; 131.9; 128.4; 125.9; 122.4 (d, ³J_{CP} = 4.1); 116.0; 114.6; 64.3 (d, ¹J_{CP} = 220.4); 57.1; 54.1 (d, ²J_{CP} = 6.2). Anal Calcd for C₁₄H₁₆N₃O₄P: C, 52.34; H, 5.02; N, 13.08; P, 9.64. Found: C, 52.33; H, 5.05; N*, 12.38; P, 9.08. *No better analysis could be obtained.

Dimethyl 1-diazo-2-oxo-2-(2-perhydroazepinophenyl)ethylphosphonate (1d)

Yield 3.27 g (93 %). Oil. IR (CHCl₃): 2120, 1630, 1260, 1050. ¹H-NMR: δ 7.30-7.20 (m, 2H); 6.94 (d, 1H, J = 8.1); 6.85 (br t, 1H, J = 7.5); 3.82 (d, 6H, ³J_{HP} = 11.9); 3.28-3.22 (m, 4H); 1.80-1.50 (m, 8H). ¹³C-NMR: δ 190.0 (d, ²J_{CP} = 7.3); 151.3; 131.7; 129.3; 128.3 (d, ³J_{CP} = 5.4); 119.9; 117.9; 63.3 (d, ¹J_{CP} = 221.4); 54.7; 54.1 (d, ²J_{CP} = 6.2); 28.5; 27.7. Anal Calcd for C₁₆H₂₂N₃O₄P: C, 54.70; H, 6.31; N, 11.96; P, 8.82. Found: C, 54.87; H, 6.30; N, 11.60; P, 8.31.

Dimethyl 1-diazo-2-oxo-2-(2-morpholinophenyl)ethylphosphonate (1e)

Yield 3.02 g (89 %). Mp 145-147°C. IR (CHCl₃): 2120, 1625, 1270, 1050, 1030. ¹H-NMR: δ 7.52-7.35 (m, 2H); 7.20-7.00 (m, 2H,); 3.91 (d, 6H, ³J_{HP} = 11.9); 3.85 (m, 4H); 3.03 (m, 4H). ¹³C-NMR: δ 189.4 (d, ²J_{CP} = 7.2); 149.8; 132.2; 131.5 (d, ³J_{CP} = 5.7); 129.3; 123.3; 118.0; 66.9; 63.2 (d, ¹J_{CP} = 222.8); 54.0 (d, ²J_{CP} = 6.3); 52.9. Anal Calcd for C₁₄H₁₈N₃O₅P: C, 49.56; H, 5.35; N, 12.38; P, 9.13. Found: C, 49.51; H, 5.53; N, 12.29; P, 8.81.

Dimethyl 1-diazo-2-oxo-2-(2-(1,2,3,6-tetrahydropyridino)phenyl)ethylphosphonate (1f) Yield 3.32 g (99 %). Mp 101-103°C. IR (CHCl₃): 2120, 1620, 1250, 1050, 1030. ¹H-NMR: δ 7.47-7.34 (m, 2H); 7.10-7.06 (m, 2H); 5.86 (dm, 1H, J = 10.1); 5.75 (dm, 1H, J = 10.1); 3.88 (d, 6H, ³J_{HP} = 12.0); 3.54 (m, 2H); 3.20 (t, 2H, J = 5.6); 2.40-2.30 (m, 2H). ¹³C-NMR: δ 189.7 (d, ²J_{CP} = 7.2); 150.5; 132.2; 132.1 (d, ³J_{CP} = 5.8); 129.3; 125.7; 124.9; 123.0; 118.7; 63.2 (d, ¹J_{CP} = 220.7); 54.1 (d, ²J_{CP} = 6.0); 51.9; 49.9; 26.1. Anal Calcd for C₁₅H₁₈N₃O₄P: C, 53.73; H, 5.41; N, 12.53; P, 9.24. Found: C, 53.36; H, 5.46; N, 12.70; P, 8.62.

Dimethyl 1-diazo-2-(2-(N,N-diethylamino)phenyl)-2-oxoethylphosphonate (1g) Yield 2.77 g (85 %). Oil. IR (CHCl₃): 2120, 1635, 1290, 1265, 1055, 1035. ¹H-NMR: δ 7.42-7.27 (m, 2H); 7.08-7.00 (m, 2H); 3.91 (d, 6H, ³J_{HP} = 12.0); 3.17 (q, 4H, J = 7.1); 1.05 (t, 6H, J = 7.1). ¹³C-NMR: δ 190.2 (d, ²J_{CP} = 7.0); 149.3; 132.5 (d, ³J_{CP} = 5.9); 131.5; 129.1; 122.3; 120.7; 63.1 (d, ¹J_{CP} = 220.0); 54.1 (d, ²J_{CP} = 6.0); 46.4; 11.5. Anal Calcd for C₁₄H₂₀N₃O₄P: C, 51.69; H, 6.20; N, 12.92; P, 9.52. Found: C, 51.66; H, 6.34; N, 12.94; P, 9.10.

Dimethyl 1-diazo-2-oxo-2-(2-(N-allyl,N-methylamino)phenyl)ethylphosphonate (1h) Yield 2.75 g (85 %). Oil. IR (neat): 2120, 1635, 1270, 1060, 1030. ¹H-NMR: δ 7.42-7.30 (m, 2H); 7.06-6.99 (m, 2H); 5.84 (ddt, 1H, J =17.4, J = 9.8, J = 6.3); 5.21 (dd, 1H, J =17.4, J = 1.4); 5.19 (dd, 1H, J = 9.8, J = 1.4); 3.89 (d, 6H, ³J_{HP} = 11.9); 3.64 (d, 2H, J = 6.3); 2.76 (s, 3H). ¹³C-NMR: δ 189.7 (d, ²J_{CP} = 7.3); 150.4; 133.9; 131.9; 130.7 (d, ³J_{CP} = 5.5); 129.3; 123.1; 118.7; 118.3; 63.4 (d, ¹J_{CP} = 220.8); 60.5; 54.1 (d, ²J_{CP} = 6.2); 40.1. Anal Calcd for C₁₄H₁₈N₃O₄P: C, 52.01; H, 5.61; N, 13.00; P, 9.58. Found: C, 51.46; H, 5.70; N, 12.71; P, 9.24.

Dimethyl 1-diazo-2-oxo-2-(2-(N-benzyl,N-methylamino)phenyl)ethylphosphonate (1i) Yield 3.58 g (96 %). Mp 75-77°C. IR (CHCl₃): 2120, 1625, 1260, 1050, 1030. ¹H-NMR: δ 7.43-6.97 (m, 9H); 4.23 (s, 2H); 3.87 (d, 6H, ³J_{HP} = 11.9); 2.69 (s, 3H). ¹³C-NMR: δ 189.8 (d, ²J_{CP} = 7.3); 150.4; 137.0; 132.0; 130.8 (d, ³J_{CP} = 5.3); 129.1; 128.4; 128.3; 127.5; 122.2; 119.2; 63.4 (d, ¹J_{CP} = 236.5); 61.6; 54.1 (d, ²J_{CP} = 6.1); 40.1. Anal Calcd for C₁₈H₂₀N₃O₄P: C, 57.91; H, 5.40; N, 11.25; P, 8.30. Found: C, 57.67; H, 5.41; N, 11.24; P, 8.23.

Dimethyl 1-diazo-2-oxo-2-(3-(2-piperidinopyridinyl)) ethylphosphonate (18) Yield 2.98 g (88 %). Mp 66-68°C. IR (CHCl₃): 2100, 1620, 1260, 1060, 1030. ¹H-NMR: δ 8.32 (dd, 1H, J = 4.8, J = 2.0); 7.67 (dd, 1H, J = 7.5, J = 2.0); 6.86 (dd, 1H, J = 7.5, J = 4.8); 3.90 (d, 6H, ³J_{HP} = 11.9); 3.32 (m, 4H); 1.80-1.60 (m, 6H). ¹³C-NMR: δ 188.0 (d, ²J_{CP} = 7.5); 159.1; 150.4; 138.9; 122.4 (d, ³J_{CP} = 5.7); 115.7; 63.1 (d, ¹J_{CP} = 222.5); 54.2 (d, ²J_{CP} = 6.3); 51.4; 26.0; 24.3. Anal Calcd for C₁₄H₁₉N₄O₄P: C, 49.71; H, 5.66; N, 16.56; P, 9.16. Found: C, 49.92; H, 5.60; N, 16.97; P, 8.98.

THERMOLYSIS OF α -DIAZO- β -KETOPHOSPHONATES (1a-i) and 18

General Procedure: to the α -diazo- β -ketophosphonate 1 or 18 (500 mg) was added anhydrous benzene (15 ml) and benzene was evaporated *in vacuo*. Compounds 1 or 18 were then dissolved in anhydrous toluene (50 ml) and the solution was refluxed under nitrogen until the disappearance of compound 1 or 18 was completed as judged by TLC. The toluene was then evaporated *in vacuo* and the residue was purified. For ¹³C-NMR data of mesoionic compounds 4 and 19, see Table (2). No microanalysis was made for the ylides 5b,d and 20 which were isolated only in small quantities.

Thermolysis of 1a: Time of decomposition, 3.5 h. Purification of the residue by chromatography on silica gel yielded 4a after elution with acetone and 5a after elution with ethyl acetate-methanol 50:50.

3-Dimethylphosphono-2-oxo-1-spiropiperidinoindolinium enolate (4a) Yield 197 mg (43 %). Mp = 170-172 °C. IR (CHCl₃): 1750, 1600, 1580, 1055, 1025 cm⁻¹. ¹H-RMN: δ 7.57 (d, 1H, J = 8.0); 7.52 (d, 1H, J = 7.9); 7.30 (pt, 1H, J = 7.7); 6.94 (pt, 1H, J = 7.8); 3.69 (d, 6H, ³J_{HP} = 11.5); 3.55 (td, 2H, J = 13.2, J = 3.8); 2.88 (pd, 2H, J = 13.2); 2.33-2.18 (m, 2H); 2.07-1.98 (m, 3H); 1.76-1.71 (m, 1H). Anal Calcd for C₁₅H₂₀NO₄P: C, 58.25; H, 6.52; N, 4.53; P, 10.01. Found: C, 58.10; H, 6.79; N, 4.58; P, 9.77.

2-Dimethylphosphono-3-oxo-1-spiropiperidinoindolinium enolate (5a)

Yield 202 mg (44 %). Mp = 219-221 °C. IR (CHCl₃) : 1620, 1585, 1035 cm⁻¹. ¹H-RMN: δ 7.95 (d, 1H, J = 8.2); 7.82 (d, 1H, J = 6.4); 7.65 (pt, 1H, J = 7.4); 7.52 (pt, 1H, J = 6.8); 4.87 (td, 2H, J = 13.4, J = 4.0); 3.74 (d, 6H, ³J_{HP} = 12.1) ; 3.22 (pd, 2H, J = 13.4); 2.47-2.31 (m, 2H); 2.12-1.97 (m, 3H); 1.89-1.77 (m, 1H). ¹³C-RMN: δ 172.11 (d, ²J_{CP} = 16.8); 150.88 (d, ³J_{CP} = 8.6); 135.44 (d, ³J_{CP} = 15.7); 130.85; 129.10; 123.79 (d, J_{CP} = 0.7); 119.27 (d, J_{CP} = 1.2); 105.75 (d, ¹J_{CP} = 226.8); 64.40; 52.77 (d, ²J_{CP} = 5.6); 22.72; 19.50. Anal Calcd for C₁₅H₂₀NO₄P: C, 58.25; H, 6.52; N, 4.53; P, 10.01. Found: C, 58.13; H, 6.39; N, 4.50; P, 9.95.

Thermolysis of 1b: Time of decomposition, 1 h. Purification of the residue by chromatography on silica gel yielded 4b after elution with acetone and a small amount of 5b after elution with ethyl acetate-methanol 50:50.

3-Dimethylphosphono-2-oxo-1-spiropiperidinoindolinium enolate (4b)

Yield 402 mg (88 %). Mp 140-141°C. IR (CHCl₃): 1735, 1600, 1575, 1050, 1020. ¹H-NMR: δ 7.47 (dd,1H, J = 7.8, J = 1.2); 7.23 (td, 1H, J = 7.8, J = 1.1); 7.05 (pd, 1H, J = 8.0); 6.88 (td, 1H, J = 7.7, J = 1.2); 4.05-3.70 (m, 8H with a doublet at 3.76, ³J_{HP} = 11.4); 3.40-3.25 (m, 2H); 2.50-2.35 (m, 2H). Anal Calcd for C₁₄H₁₈NO₄P: C, 56.95; H, 6.14; N, 4.74; P, 10.49. Found: C, 56.69; H, 6.09; N, 4.73; P, 10.20.

2-Dimethylphosphono-3-oxo-1-spiropiperidinoindolinium enolate (5b) Yield 9 mg (2 %). Oil. IR (CHCl₃): 1600, 1580, 1030. ¹H-NMR: δ 7.80 (dd, 1H, J = 8.4, J = 1.5); 7.60-7.32 (m, 3H); 4.80-4.64 (m, 2H); 3.80-3.60 (m, 8H with a doublet at 3.75, ³J_{HP} = 11.9); 2.80-2.55 (m, 2H); 2.55-2.30 (m, 2H).

Thermolysis of 1c: Time of decomposition, 1 h. Purification of the residue by recrystallisation from toluene yielded 4c.

3-Dimethylphosphono-2-oxo-1-spiro(3-pyrrolino)indolinium enolate (4c)

Yield 301 mg (66 %). Mp 173-175°C. IR (CHCl₃): 1750, 1605, 1580, 1055, 1020. ¹H-NMR: δ 7.47 (dd,1H, J = 7.8, J = 0.9); 7.26 (ddd, 1H, J = 7.8, J = 7.6; J = 0.9); 7.14 (d, 1H, J = 7.9); 6.88 (m, 1H); 6.09 (m, 2H); 4.78 (d, 2H, J = 13.7); 4.09 (d, 2H, J = 13.1); 3.80 (d, 6H, ³J_{HP} = 11.5). Anal Calcd for C₁₄H₁₆NO₄P: C, 57.34; H, 5.50; N, 4.78; P, 10.56. Found: C, 57.41; H, 5.51; N, 4.80; P, 10.68.

Thermolysis of 1d: Time of decomposition, 2.5 h. Purification of the residue by chromatography on silica gel yielded 4d after elution with acetone and a small amount of 5d after elution with ethyl acetate-methanol 50:50.

3-Dimethylphosphono-2-oxo-1-spiroperhydroazepinoindolinium enolate (4d)

Yield 304 mg (66 %). Mp 193-195°C. IR (CHCl₃): 1750, 1600, 1575, 1055, 1025. ¹H-NMR: δ 7.52 (dd, 1H, J = 8.2, J = 1.4); 7.25 (m, 2H); 6.92 (td, 1H, J = 7.7, J = 1.3); 3.82 (m, 2H); 3.80 (d, 6H, ³J_{HP} = 11.4); 3.24 (dd, 2H, J = 13.8, J = 8.6); 2.50-2.30 (m, 2H); 2.20-1.80 (m, 6H). Anal Calcd for C₁₆H₂₂NO₄P: C, 59.44; H, 6.86; N, 4.33; P, 9.58. Found: C, 59.14; H, 6.90; N, 4.31; P, 9.11.

2-Dimethylphosphono-3-oxo-1-spiroperhydroazepinoindolinium enolate (5d) Yield 28 mg (6%). Mp 197-199°C. IR (CHCl₃): 1620, 1600, 1580, 1055, 1035. ¹H-NMR: δ 7.79 (dd, 1H, J = 8.8, J = 6.5); 7.65 (dd, 1H, J = 6.9, J = 1.9); 7.51 (m, 2H); 4.59 (dd, 2H, J = 13.6, J = 9.4); 3.72 (d, 6H, ³J_{HP} = 11.8); 3.64 (dd, 2H, J = 13.6, J = 6.7); 2.60-2.40 (m, 2H); 2.20-1.70 (m, 6H).¹³C-NMR: δ 171.7 (d, ²J_{CP} = 17.3); 153.2 (d, ³J_{CP} = 8.8); 134.3 (d, ³J_{CP} = 15.6); 130.4; 129.8; 123.5; 117.1; 107.4 (d, ¹J_{CP} = 227.9); 71.6; 52.4 (d, ²J_{CP} = 5.7); 28.6; 25.7.

Thermolysis of 1e: Time of decomposition, 3 h. Purification of the residue by chromatography on silica gel yielded 12 and then 4e after elution with acetone and 5e after elution with ethyl acetate-methanol 50:50. A second purification of the sample of 5e by chromatography with ethyl acetate-methanol 50:50 as eluent was necessary to obtain pure 5e.

3-Dimethylphosphono-2-oxo-1-spiromorpholinoindolinium enolate (4e)

Yield 243 mg (53 %). Mp 203-205°C. IR (CHCl₃): 1760, 1595, 1575, 1050, 1025. ¹H-NMR: δ 7.62-7.46 (m, 2H); 7.31 (dt, 1H, J = 7.7, J = 1.0); 6.96 (td, 1H, J = 7.7, J = 1.2); 4.49 (td, 2H, J = 13.1, J = 4.3); 4.22 (ddd, 2H, J = 13.1, J = 9.0, J = 3.0); 3.89-3.61 (m, 8 H, with a doublet at 3.78, ³J_{HP} = 11.5); 3.04 (dt, 2H, J = 13.1, J = 3.0). HRMS (EI) m/z calcd for C₁₄H₁₈NO₅P (M⁺) 311.0923, found 311.0912. LRMS (EI) m/z (rel.int.): 311 (37.1); 280 (48.1); 268 (44.6); 253 (68.7); 238 (21.6); 225 (25.9); 174 (100); 91 (37.7); 77 (24.7).

2-Dimethylphosphono-3-oxo-1-spiromorpholinoindolinium enolate (5e)

Yield 55 mg (12%). Mp 218-220°C (dec.). IR (CHCl₃): 1620, 1585, 1135, 1040. ¹H-NMR: δ 8.10 (d, 1H, J = 8.1); 7.93 (d, 1H, J = 7.3); 7.66 (pt, 1H, J = 7.4); 7.53 (dd 1H, J = 8.1; J = 7.3); 5.19 (td, 2H, J = 13.0, J = 4.5); 4.47 (td, 2H, J = 13.0, J = 2.0); 4.15 (dd, 2H, J = 13.0, J = 4.5); 3.81 (d, 6 H, ³J_{HP} = 12.0); 3.20 (dd, 2H, J = 13.0, J = 2.0). ¹³C-NMR: δ 171.7 (d, ²J_{CP} = 16.8); 150.9 (³J_{CP} = 8.4); 135.4 (³J_{CP} = 15.4); 131.2; 129.4; 124.0; 118.7; 105.7 (d, ¹J_{CP} = 226.9); 62.6; 62.5; 52.7 (d, ²J_{CP} = 5.7). HRMS (EI) m/z calcd for C₁₄H₁₈NO₅P (M⁺) 311.0923, found 311.0915. LRMS (EI) m/z (rel.int.): 311 (34.1); 268 (14.6); 202 (100); 172 (26.1); 110 (34.1); 77 (18.1).

5a-Dimethylphosphono-6-oxo-1,4-oxazepano[4,5-a]indoline (12)

Yield 119 mg (26 %). Oil. IR (CHCl₃): 1690, 1610, 1580, 1250, 1060, 1030. ¹H-NMR: δ 7.58 (d, 1H, J = 7.8); 7.47 (pt, 1H, J = 7.8); 6.86-6.75 (m, 2H); 3.95-3.92 (m, 2H); 3.85-3.70 (m, 5 H with a doublet at 3.77, ³J_{HP} = 10.7); 3.60 (d, 3 H, ³J_{HP} = 10.4); 3.51-3.30 (m, 1 H); 2.86-2.51 (m, 3 H). ¹³C-NMR: δ 197.6 (d,

 ${}^{2}J_{CP} = 1.3$; 160.8 (d, ${}^{3}J_{CP} = 7.3$); 137.8; 125.2; 121.9 (d, ${}^{3}J_{CP} = 1.3$); 119.1; 110.4; 72.8 (d, ${}^{1}J_{CP} = 154.4$); 68.2 (d, ${}^{4}J_{CP} = 2.5$); 66.8 (d, ${}^{3}J_{CP} = 14.6$); 54.9 (d, ${}^{2}J_{CP} = 7.2$); 53.9 (d, ${}^{2}J_{CP} = 7.7$); 47.1; 35.9. HRMS (EI) m/z calcd for C₁₄H₁₈NO₅P (M⁺) 311.0923, found 311.0920. LRMS (EI) m/z (rel.int.): 311 (17.2); 202 (100); 172 (14.0); 144 (5.1); 105 (3.7); 77 (6.8).

Thermolysis of 1f: Time of decomposition, 3 h. The ¹H-NMR spectrum of the crude residue showed that it contained of 51:40:9-mixture (determined by ¹H-NMR) of 4f, 13 and 14. The purification of the residue by chromatography on silica gel yielded 57 mg of 13 and 175 mg of a mixture of 13+14 after elution with ethyl acetate-acetone 80:20 and then 119 mg of 4f after elution with acetone, the total yield being 77 %.

3-Dimethylphosphono-2-oxo-1-spiro-(1',2',3',6'-tetrahydropyridino)indolinium enolate (4f) Isolated yield 119 mg (26 %). Mp 142-144°C. IR (CHCl₃): 1750, 1600, 1575, 1050, 1025. ¹H-NMR: δ 7.57 (dd, 1H, J = 7.3, J = 1.0); 7.34-7.24 (m, 2H); 6.89 (td, 1H, J = 7.3, J = 1.3); 6.18 (m, 1H); 6.02 (m, 1H); 4.38 (m, 1H); 3.79 (d, 6H, ³J_{HP} = 11.4); 3.67 (m, 1H); 3.23 (m, 1H); 2.93 (m, 1H); 2.74-2.48 (m, 2H). Anal Calcd for C₁₅H₁₈NO₄P: C, 58.63; H, 5.90; N, 4.56; P, 10.08. Found: C, 58.19; H, 5.88; N, 4.68; P, 9.71.

5a-Dimethylphosphono-6-oxo-(2, 3, 6, 7-tetrahydro-1H-azepino)[1,2-a]indoline (13) Isolated yield 57 mg (12%). Mp 113-115°C. IR (CHCl₃): 1690, 1610, 1580, 1060, 1030. ¹H-NMR: δ 7.56 (d, 1H, J = 7.7); 7.43 (ddd, 1H, J = 8.3, J = 7.5, J = 1.3); 6.90 (d, 1H, J = 8.3); 6.77 (dd, 1H, J = 7.7, J = 7.5); 5.23 (br s, 2H); 4.12-3.55 (m, 2H); 3.75 (d, 3H, ³J_{HP} = 10.7); 3.57 (d, 3H, ³J_{HP} = 10.5); 3.25-3.07 (m, 1H); 2.89-2.76 (m, 1H); 2.69-2.52 (m, 1H); 2.35-2.26 (m, 1H). ¹³C-NMR: δ 197.9; 161.0 (d, ⁴J_{CP} = 7.4); 137.0; 130.5; 124.9; 123.0 (d, ³J_{CP} = 1.3); 121.5 (d, ³J_{CP} = 16.4); 118.9; 111.2; 75.3 (d, ¹J_{CP} = 152.7); 54.8 (d, ²J_{CP} = 7.2); 53.8 (d, ²J_{CP} = 7.7); 40.3; 28.6 (d, ²J_{CP} = 2.4); 28.0. Anal Calcd for C₁₅H₁₈NO₄P: C, 58.63; H, 5.90; N, 4.56; P, 10.08. Found: C, 58.47; H, 6.00; N, 4.36; P, 9.02.

Thermolysis of 1g: Time of decomposition, 3.5 h. Purification of the residue by chromatography on silica gel yielded 21 after elution with ethyl acetate-pentane 75:25 and 4g after elution with ethyl acetate-methanol 80:20.

3-Dimethylphosphono-1,1-diethylindolium-2-olate (4g)

Yield 306 mg (67 %). Mp 163-165°C. IR (CHCl₃): 1735, 1605, 1575, 1050, 1020. ¹H-NMR: δ 7.55 (d, 1H, J = 7.8); 7.30 (td, 1H, J = 7.6, 1.9); 7.02-6.94 (m, 2H); 3.77 (d, 6H, ³J_{HP} = 11.5); 3.66-3.48 (m, 2H); 3.34-3.21 (m, 2H); 0.93 (t, 6H, J = 7.1). ¹³C-NMR: δ 165.5 (d, ³J_{CP} = 24.8); 142.1 (d, J_{CP} = 15.1); 131.1 (d, J_{CP} = 13.3); 130.5; 120.9; 119.3; 116.7; 68.3 (d, ¹J_{CP} = 217.3); 55.2; 52.0 (d, ²J_{CP} = 5.1); 8.1. Anal Calcd for C₁₄H₂₀NO₄P: C, 56.56; H, 6.78; N, 4.71; P, 10.42. Found: C, 56.27; H, 6.56; N, 4.79; P, 10.38.

Dimethyl 1-ethyl-3-hydroxy-2-indolephosphonate (21)

Yield 29 mg (7%). Oil. IR (CHCl₃): 3260, 1605, 1045, 1020. ¹H-NMR: δ 9.00-8.60 (brs, 1H); 7.76 (d, 1H, J = 8.1); 7.59-7.22 (m, 2H); 7.07 (td, 1H, J = 7.3, 0.9 Hz); 4.11 (q, 2H, J = 7.1); 3.76 (d, 6H, ³J_{HP} = 11.6); 1.31 (t, 3H, J = 7.1). ¹³C-NMR: δ 151.6 (d, ²J_{CP} = 15.8); 137.2 (d, ³J_{CP} = 12.4); 126.4; 120.1; 118.8; 117.7 (d, ³J_{CP} = 15.3); 109.6 (d, ⁴J_{CP} = 2.0); 100.1 (d, ¹J_{CP} = 218.6); 52.84 (d, ²J_{CP} = 4.5); 40.0; 14.3. Anal Calcd for C₁₂H₁₆NO₄P: C, 53.53; H, 5.99; N, 5.20; P, 11.50. Found: C, 53.68; H, 5.83; N, 5.44; P, 11.34. **Thermolysis of 1h**: Time of decomposition 2.5 h. Purification of the residue by chromatography on silica gel, eluting with ethyl acetate, yielded 23h and then 22h.

Dimethyl 3-allyl-1-methyl-2-oxo-3-indolinephosphonate (22h)

Yield 196 mg (43 %). Mp 75-77°C. IR (CHCl₃): 1705, 1605, 1490, 1050, 1030. ¹H-NMR: δ 7.48 (dt, 1H, J = 7.4, J = 1.5); 7.32 (tt, 1H, J = 7.7, J = 1.5, ⁶J_{HP} = 1.5); 7.10 (br t, 1H, J = 7.4); 6.84 (d, 1H, J = 7.7); 5.38-5.15 (m, 1H); 5.01 (br d, 1H, J = 16.9); 4.87 (br d, 1H, J = 9.8); 3.84 (d, 3H, ³J_{HP} = 10.9); 3.56 (d, 3H, ³J_{HP}); 5.15 (m, 1H); 5.01 (br d, 1H, J = 16.9); 4.87 (br d, 1H, J = 9.8); 3.84 (d, 3H, ³J_{HP}); 5.16 (d, 3H, ³J_{HP}); 5.16 (d, 3H, ³J_{HP}); 5.17 (d, 3H, ³J_{HP}); 5.17 (d, 3H, ³J_{HP}); 5.18 (d, 3H, ³J_{HP}

= 10.7); 3.23 (s, 3H); 3,03 (m, 2H). ¹³C-NMR: δ 173.0 (d, ²J_{CP} = 2.8); 144.2 (d, ³J_{CP} = 7.8); 130.7 (d, ³J_{CP} = 14.4); 128.5 (d, J_{CP} = 2.8); 125.6 (d, J_{CP} = 3.7); 124.9 (d, J_{CP} = 6.0); 122.7 (d, J_{CP} = 3.1); 119.7; 108.1 (d, ²J_{CP} = 1.8); 54.7 (d, ¹J_{CP} = 137.4); 54.7 (d, ²J_{CP} = 6.8); 53.8 (d, ²J_{CP} = 7.6); 36.3 (d, ²J_{CP} = 4.4); 26.6. Anal Calcd for C₁₄H₁₈NO₄P: C, 56.95; H, 6.14; N, 4.74; P, 10.49. Found: C, 56.65; H, 6.32; N, 4.71; P, 10.31.

Dimethyl 2-allyl-1-methyl-3-oxo-2-indolinephosphonate (23h)

Yield 178 mg (39 %). Oil. IR (CHCl₃): 1690, 1610, 1580, 1060, 1030. ¹H-NMR: δ 7.59 (br d, 1H, J = 7.7); 7.54-7.43 (m, 1H); 6.82 (d, 1H, J = 8.3); 6.72 (t, 1H, J = 7.1); 5.40-5.10 (m, 2H); 5.00-4.90 (m, 1H); 3.84 (d, 3H, ³J_{HP} = 10.8); 3.63 (d, 3H, ³J_{HP} = 10.7); 3.20 (s, 3H); 3.10-2.90 (m, 2H). ¹³C-NMR: δ 196.2 (d, ²J_{CP} = 2.4); 161.0 (d, ³J_{CP} = 6.0); 137.7; 130.2 (d, ³J_{CP} = 5.6); 124.7; 120.6 (d, ⁴J_{CP} = 1.6); 120.0; 117.8; 108.6; 73.1 (d, ¹J_{CP} = 148.4); 54.4 (d, ²J_{CP} = 7.0); 53.7 (d, ²J_{CP} = 7.7); 34.9; 29.7. Anal Calcd for C₁₄H₁₈NO₄P: C, 56.95; H, 6.14; N, 4.74; P, 10.49. Found: C, 56.83; H, 6.23; N, 4.84; P, 9.84.

Thermolysis of 1i: Time of decomposition 3.5 h. The purification of the residue by chromatography on silica gel, eluting with ethyl acetate, yielded 69 mg of 23i, 232 mg of a 22:78-mixture of 23i+22i and 144 mg of 22i, the total yield being of 96 %.

Dimethyl 3-benzyl-1-methyl-2-oxo-3-indolinephosphonate (22i)

Isolated yield 69 mg (15 %). Mp 116-117°C. IR (CHCl₃): 1705, 1610, 1055, 1035. ¹H-NMR: δ 7.61 (dt, 1H, J = 7.5, 1.7); 7.21 (tt, 1H, J = 7.7, 1.5); 7.10 (pd, 1H, J = 7.0); 7.05-6.93 (m, 3H); 6.90-6.83 (m, 2H); 6.57 (d, 1H, J = 7.7); 3.91 (d, 3H, ³J_{HP} = 10.9); 3.71 (A part of a H_AH_BP system, 1H, J_{AB} = 13.2, ³J_{AP} = 6.9); 3.61 (d, 3H, ³J_{HP} = 10.7); 3.46 (B part of a H_AH_BP system, 1H, J_{AB} = 13.2, ³J_{BP} = 8.0); 2.99 (s, 3H). ¹³C-NMR: δ 172.8 (d, ²J_{CP} = 2.7); 144.0 (d, ³J_{CP} = 8.1); 134.4 (d, ²J_{CP} = 15.7); 129.8; 128.9 (d, J_{CP} = 2.7); 127.6; 126.7; 126.0 (d, J_{CP} = 3.5); 124.6 (d, ²_{CP} = 5.6); 122.4 (d, J_{CP} = 2.9); 108.0 (d, J_{CP} = 1.6); 56.3 (d, ¹J_{CP} = 137.5); 54.7 (d, ²J_{CP} = 6.8); 53.8 (d, ²J_{CP} = 7.5); 37.9 (d, ²J_{CP} = 4.2); 26.3. Anal Calcd for C₁₈H₂₀NO₄P: C, 62.61; H, 5.84; N, 4.06; P, 8.97. Found: C, 62.79; H, 6.10; N, 4.03; P, 8.46.

Dimethyl 2-benzyl-1-methyl-3-oxo-2-indolinephosphonate (23i)

Isolated yield 144 mg (31 %). Oil. IR (CHCl₃): 1690, 1615, 1055, 1035. ¹H-NMR: δ 7.50 (pd, 1H, J = 7.6); 7.35 (td, 1H, J = 7.7, 1.4); 7.06 (s, 5H); 6.69-6.60 (m, 2H); 3.89 (d, 3H, ³J_{HP} = 10.8); 3.76 (A part of a H_AH_BP system, 1H, J_{AB} = 14.2, ³J_{AP} = 11.1); 3.64 (d, 3H, ³J_{HP} = 10.7); 3.45 (B part of a H_AH_BP system, 1H, J_{AB} = 14.2, ³J_{BP} = 10.5); 3.22 (s, 3H). ¹³C-NMR: δ 196.1 (d, ²J_{CP} = 3.0); 160.7 (d, ³J_{CP} = 5.0); 137.5; 133.9 (d, ³J_{CP} = 15.5); 129.6; 128.1; 127.0; 124.5; 120.4 (d, J_{CP} = 1.3); 117.6; 108.2; 74.5 (d, ¹J_{CP} = 142.6); 54.3 (d, ³J_{CP} = 7.2); 53.9 (d, ³J_{CP} = 7.8); 35.9; 30.4. Anal Calcd for C₁₈H₂₀NO₄P: C, 62.61; H, 5.84; N, 4.06; P, 8.97. Found: C, 62.25; H, 5.92; N, 4.16; P, 8.83.

Thermolysis of 18: Time of decomposition 1.5 h. Purification of the residue by chromatography on silica gel yielded 19 after elution with acetone and a small amount of 20 after elution with ethyl acetate-methanol 50:50.

7-Aza-3-dimethylphosphono-2-oxo-1-spiropiperidinoindolinium enolate (19) Yield 238 mg (52 %). Mp 179-181°C. IR (CHCl₃): 1740, 1590, 1575, 1260, 1060, 1035. ¹H-NMR: δ 7.88 (d, 1H, J = 4.9); 7.82 (d, 1H, J = 7.8); 7.19 (dd, 1H, J = 7.8, J = 4.9); 3.78 (d, 6H, ³J_{HP} = 11.5); 3.54 (m, 2H); 3.02 (m, 2H); 2.75 (m, 2H); 2.10-1.90 (m, 3H); 1.90-1.60 (m, 1H). Anal Calcd for C₁₄H₁₉N₂O₄P: C, 54.19; H, 6.17; N, 9.03; P, 9.98. Found: C, 53.83; H, 6.19; N, 9.16; P, 9.70.

7-Aza-2-dimethylphosphono-3-oxo-1-spiropiperidinoindolinium enolate (20)

Yield 37 mg (8 %). Mp 193-195°C. IR (CHCl₃): 1620, 1595, 1570, 1035. ¹H-NMR: δ 8.41 (dd, 1H, J = 4.9, J = 1.7); 8.15 (dd, 1H, J = 7.5, J = 1.7); 7.51 (dd, 1H, J = 7.5, J = 4.9); 4.78 (m, 2H); 3.75 (d, 6H, ³J_{HP} = 11.9); 3.17 (m, 2H); 3.10-2.80 (m, 2H); 2.10-1.90 (m, 1H); 1.90-1.60 (m, 3H). ¹³C-NMR: δ 170.0 (d, ²J_{CP} = 15.9); 163.7; 147.2; 132.5; 127.8 (d, ³J_{CP} = 14.5); 125.9; 105.5 (d, ¹J_{CP} = 225.9); 64.1; 52.6 (d, ²J_{CP} = 5.6); 22.4; 20.3.

REACTION OF WATER WITH THE 3-DIMETHYLPHOSPHONO-2-OXO-1-SPIROPIPERIDINOINDOLINIUM ENOLATE (4a)

A mixture of 4a (187 mg, 0.6 mmol), water (4 ml) and silica gel (\sim 5 mg) was refluxed for 4 h. After cooling, and extraction with ethyl acetate (3 x 15 ml), the organic layer was dried, filtered and concentrated to give a residue which was purified by chromatography eluting with ethyl acetate to afford 11.

Dimethyl (2-piperidinophenyl)methylphosphonate (11)

Yield 156 mg (89 %). Oil. IR (CHCl₃): 1595,1250,1055,1030. ¹H-NMR: δ 7.47 (dt, 1H, J = 7.5, J = 2.0); 7.23-7.02 (m, 3H); 3.70 (d, 6H, ³J_{HP} = 10.8); 3.41 (d, 2H, ²J_{HP} = 22,2); 2.96-2.71 (m, 4H); 1.83-1.48 (m, 6H). ¹³C-NMR: δ 153.3 (d, ³J_{CP} = 8.4); 130.6 (d, ³J_{CP} = 5.0); 127.8 (d, J_{CP} = 3.2); 127.4 (d, ²J_{CP} = 8.0); 124.0 (d, J_{CP} = 3.2); 121.1 (d, J_{CP} = 2.4); 54.4; 53.0 (d, ²J_{CP} = 6.7); 26.7; 26.2 (d, ¹J_{CP} = 138.7); 24.2. Anal Calcd for [C₁₄H₂₂NO₃P,0.5 H₂O]*: C, 57.52; H, 7.93; N, 4.79; P, 10.60. Found: C, 57.43; H, 7.76; N, 4.68; P, 10.38. *Identical microanalyses were obtained with samples of 11 prepared by independent experiments.

REFERENCES AND NOTES

- a) Meth-Cohn, O.; Suschitzky, H. Adv. Heterocycl. Chem., 1972, 14, 211-278. b) Verboom, W.; Reinhoudt, D. N. Recl. Trav. Chim. Pays-Bas, 1990, 109, 311-324. c) Meth-Cohn, O. Adv. Heterocycl. Chem. 1996, 65, 1-37.
- 2. A 1,4-shift of a NCH₂ proton to a highly basic central allenic carbon atom has been observed in the thermal isomerization of 1-morpholino-3-phenyl(or vinyl) allenes. See: Mayer, T.; Maas, G. *Tetrahedron Lett.* **1992**, *33*, 205-208.
- 3. Edstrom, E.D. J. Am. Chem. Soc. 1991, 113, 6690-6692.
- 4. Roberts, S.M.; Smith, C.; Thomas, R.J. J. Chem. Soc., Perkin Trans. 1 1990, 1493-1495.
- 5. Maurya, R.; Pittol, C.A.; Pryce, R.J.; Roberts, S.M.; Thomas, R.J.; Williams J.O. J. Chem. Soc., Perkin Trans. 1 1992, 1617-1621.
- a) Chelain, E.; Goumont, R.; Hamon, L.; Parlier, A.; Rudler, M.; Rudler, H.; Daran, J.-C.; Vaissermann, J. J. Am. Chem. Soc. 1992, 114, 8088-8098. b) Chelain, E.; Parlier, A.; Audouin, M.; Rudler, H.; Daran, J.-C.; Vaissermann, J. J. Am. Chem. Soc. 1993, 115, 10568-10580.
- a) Andriamiadanarivo, R.; Pujol, B.; Chantegrel, B.; Deshayes, C.; Doutheau, A. Tetrahedron Lett. 1993, 34, 7923-7924. b) Chen, Y. P.; Chantegrel, B.; Deshayes, C. Heterocycles, 1995, 41, 175-186. c) Collomb, D.; Deshayes, C.; Doutheau, A. Tetrahedron 1996, 52, 6665-6684. d) Collomb, D.; Chantegrel, B.; Deshayes, C. Tetrahedron 1996, 52, 10455-10472.
- 8. Chantegrel, B.; Deshayes, C.; Faure, R. Tetrahedron Lett. 1995, 36, 7859-7862.

- 9. Tidwell, T.T. Ketenes, Wiley: New York, 1995, p. 78.
- For reviews concerning ammonium ylides, see: a) Padwa, A., Hornbuckle, S.F. Chem. Rev. 1991, 91, 263-309. b) Padwa, A., Weingarten, M.D. Chem. Rev. 1996, 96, 223-269.
- a) West, F.G.; Naidu, B.N. J. Am. Chem. Soc. 1993, 115, 1177-1178. b) West, F.G.; Naidu, B.N. J. Org. Chem. 1994, 59, 6051-6056. c) West, F.G.; Naidu, B.N.; Tester, R.W. J. Org. Chem. 1994, 59, 6892-6894.
- 12. Zaragoza, F. Synlett, 1995, 237-238.
- a) Clark, J.S.; Hodgson, P.B. J. Chem. Soc., Chem. Commun. 1994, 2701-2702. b) Clark, J.S.; Hodgson, P.B. Tetrahedron Lett. 1995, 36, 2519-2522.
- 14. Jemison, R.W.; Mageswaran, S.; Ollis, W.D.; Potter, S.E.; Pretty, A.J.; Sutherland, I.O.; Thebtaranonth, Y. J. Chem. Soc., Chem. Comm. 1970, 1201-1202.
- 15. Bailey, N.A.; Hull, S.E.; Kersting, G.F.; Morrison, J. J. Chem. Soc., Chem. Comm. 1971, 1429-1430.
- 16. Martin, J.; Meth-Cohn, O.; Suschitzky, H. J. Chem. Soc. Perkin Trans. 1 1974, 2451-2458.
- a) Waldron, N.M.; Montevalli, M.; Azam, S.; Dasopolous, P.C. J. Chem. Soc., Chem. Commun. 1995, 81-82. b) Waldron, N.M.; Raza, M. J. Chem. Soc. Perkin Trans.1 1996, 271-276.
- a) Niewiadomski, K.B.; Suschitzky, H. J. Chem. Soc., Perkin Trans. 1 1975, 1679-1682. b) Nijhuis,
 W.H.N.; Verboom, W.; Reinhoudt, D.N. Synthesis 1987, 641-645.
- 19. Ding, Z.; Tufariello, J.J. Synth. Commun. 1990, 20, 227-230.
- 20. The X ray data of 4a and 5a are registered at the Cambridge Cristallographic Data Centre, Cambridge (U.K.).
- 21. The N(1)C(2)C(3)C(3a)C(7a) ring is almost planar, the C(7a) atom being at a distance of 0.027 Å from the N(1)C(2)C(3)C(3a) mean plane. The dihedral angle between the C(3a)C(4)C(5)C(6)C(7)C(7a) and N(1)C(2)C(3)C(3a)C(7a) planes is 1.10°. The piperidinium ring adopts a perfect chair conformation; the dihedral angle between his mean plane and the N(1)C(2)C(3)C(3a)C(7a) mean plane is 88.84°.
- Handbook of Chemistry and Physics; Lide, D.R., Editor-in-Chief; 72nd Edition 1991-1992, CRC Press. Bond lenghts in crystalline organic compounds 9-3 to 9-17.
- 23. Thiessen, W.E.; Hope, H. J. Am. Chem. Soc. 1967, 89, 5977-5978.
- a) Zvilichovsky, G.; David, M. Heterocycles, 1987, 26, 961-968. b) Zvilichovsky, G.; Gurvich, V.;
 Cohen, S. J. Chem. Research (S) 1994, 335; J. Chem. Research (M) 1994, 1801-1812.
- 25. Edstrom, E.D.; Wei, Y.; Gordon, M. J. Org. Chem. 1994, 59, 2473-2481.
- 26. Ollis, W.D.; Rey, M.; Sutherland, I.O. J. Chem. Soc. Perkin Trans. 1 1983, 1009-1027.
- a) Heard, G.L.; Yates, B.F. Aust. J. Chem. 1994, 47, 1685-1694 and 1995, 48, 1413-1423. b)
 Heard, G.L.; Yates, B.F. J. Org. Chem. 1996, 61, 7276-7284.
- 28. When a toluenic solution of the ylide 5a was refluxed for 24 h, no transformation into the mesoionic compound 4a was observed.
- 29. For instance as described in ref. 6a, refluxing the mesoionic compound XI (see structure in Table 2) in toluene yielded a bridgehead lactam through a [1,3] shift.
- 30. Regitz, M.; J. Hocker, J.; Liedhegener, A. Org. Synth. 1973, Coll. Vol. 5, 179.
- 31. For explosive hazard of diazotransfer reagents, see: Bollinger, F.W.; Tuma, L.D. Synlett, 1996, 407-413.

(Received in Belgium 15 January 1997; accepted 17 April 1997)