

The mechanism of the reaction may in agreement with the usual thermal (as opposed to photochemical) decomposition of Δ^2 -triazolines be formulated according to eq. 3 [3]. The major pathway of the betaine **7** is the *Wagner-Meerwein* (or *Demjanov-Tiffeneau* type) rearrangement to the imidate ester **3d**. Side products are probably derived from the intermediate aziridine **8**, which being unstable undergoes further rearrangements.

We wish to thank the *Rulgers Research Council* for financial assistance.

REFERENCES

- [1] Parts I and II, in press.
- [2] *L. Wolff*, Liebigs Ann. Chem., **394**, 23 (1912).
- [3] *G. L'Abbé*, Chem. Rev. **69**, 345 (1969); *T. Sheradsky* in 'The Chemistry of the Azido Group', *S. Patai*, ed. Interscience, New York, 1971, p. 359ff; 373ff; *W. Lwowski*, *ibid.*, p. 529; and references cited therein.
- [4] *R. Fusco*, *G. Bianchetti* & *D. Pocar*, Gazz. chim. ital., **91**, 933 (1961); *Cmp. R. M. Scribner*, Tetrahedron Letters 1967, 4737.
- [5] *H. O. House*, *L. J. Czuba*, *M. Gall* & *H. D. Olmstead*, J. org. Chemistry **34**, 2324 (1969), and references cited therein.

Fourth Hungarian Bioflavonoid Symposium

Keszthely (Lake Balaton), September 20–22, 1973

Further informations: Prof. Lorant Farkas, Institute of Organic Chemistry, Technical University, 1111 Budapest, Gellert tér 4.

The Third International Symposium on Polyhalogen Compounds

Barcelona, October 22–26, 1973

Further informations: Dr. David Ellis, Instituto de Química Orgánica, Patronato Juan de la Cierva, Barcelona 17

Vortragstagung Linz 1973

Chemie für das Leben von heute und morgen

Linz, 26.–29. September 1973

Organisation: Verein Österreichischer Chemiker, Sekretariat: A-1010 Wien, Eschenbachgasse 9. Gesellschaft für Chemiewirtschaft, Sekretariat: A-1010 Wien, Eschenbachgasse 11.