# **Green Chemistry**

# PAPER

**Cite this:** *Green Chem.*, 2014, **16**, 1633

Received 24th November 2013, Accepted 19th December 2013 DOI: 10.1039/c3qc42406e

www.rsc.org/greenchem

## Introduction

With increasing awareness of the ever-growing  $CO_2$  level in the aerosphere, great efforts have been devoted to development of strategies towards the reduction of CO<sub>2</sub> emissions.<sup>1</sup> The chemical fixation of CO<sub>2</sub> is highly attractive and effective as a straightforward method to recycle CO2 into value-added products due to the great potential of CO<sub>2</sub> as an abundant, nontoxic, easily available and renewable C1 building block in organic synthesis.<sup>2</sup> However, only a small proportion of the total abundance of CO<sub>2</sub> is currently being consumed for chemical production due to its low reactivity as a result of its high thermodynamic stability and kinetic inertness. Consequently, vigorous nucleophiles like Grignard reagents and organolithium reagents, high-energy starting materials such as small ring compounds and organometallic reagents, or drastic reaction conditions are commonly required for reactions involving CO<sub>2</sub>.<sup>3</sup> Despite the significant advances in the past few decades, converting CO2 into readily usable chemicals still

<sup>a</sup>State Key Laboratory and Institute of Elemento-Organic Chemistry, Collaborative Innovation Center of Chemical Science and Engineering (Tianjin), Nankai University, Tianjin, 300071, P. R. China. E-mail: heln@nankai.edu.cn

<sup>b</sup>Key Laboratory of Advanced Energy Materials Chemistry (Ministry of Education), College of Chemistry, Nankai University, Tianjin, 300071, P. R. China.



bifunctional Ag<sub>2</sub>WO<sub>4</sub>/Ph<sub>3</sub>P system<sup>†</sup>

An efficient heterogeneous silver-catalyzed reaction for construction of the  $\alpha$ -methylene cyclic carbonate motif was developed through carboxylative assembly of propargyl alcohols and CO<sub>2</sub>. Such a CO<sub>2</sub> fixation protocol proceeded smoothly with only 1 mol% of Ag<sub>2</sub>WO<sub>4</sub> and 2 mol% of PPh<sub>3</sub> as well as atmospheric CO<sub>2</sub> at room temperature under solvent-free conditions, in an environmentally benign and low energy manner along with an easy operating procedure. Notably, up to 98% isolated yields of carbonates could be attained with exclusive chemo-selectivity. In addition, the dual activation capacity of Ag<sub>2</sub>WO<sub>4</sub> towards both the propargylic substrate and CO<sub>2</sub> is based on which cooperative catalytic mechanism by the silver cation and the tungstate anion is proposed. Recycling trials on carboxylative cyclization of propargyl alcohols and CO<sub>2</sub> illustrate that the catalyst can be reused at least 4 times with retention of high catalytic activity and selectivity. Especially, it allows the direct and effective application in the one-pot synthesis of various oxazolidinones bearing exocyclic alkenes and carbamates in moderate to high yields upon the alternative introduction of primary or secondary amines.

Efficient chemical fixation of CO<sub>2</sub> promoted by a

remains a daunting challenge presumably due to limitations existing in those known processes such as tedious reaction procedures, harsh reaction conditions, or low yields.<sup>4</sup> Therefore, development of effective methodologies using  $CO_2$  as a feedstock under mild conditions particularly low pressure of  $CO_2$  (ideally at 1 atm) could be still highly desirable.

One of the most promising examples of  $CO_2$  fixation is carboxylative cyclization of propargyl alcohols with  $CO_2$  to access  $\alpha$ -alkylidene cyclic carbonates, which stands for an important class of heterocyclic compounds with a wide range of applications in organic synthesis (Scheme 1).<sup>5</sup> In this context, numerous catalytic systems including Ru,<sup>6</sup> Pd,<sup>7</sup> Co,<sup>8</sup> Cu,<sup>9</sup> Ag,<sup>10</sup> Fe,<sup>11</sup> phosphines,<sup>12</sup> N-heterocyclic carbenes (NHC),<sup>13</sup> K<sub>2</sub>CO<sub>3</sub>/crown ethers,<sup>14</sup> bicyclic guanidines,<sup>15</sup> silver-NHC complexes,<sup>16</sup> or N-heterocyclic olefin/CO<sub>2</sub> adducts<sup>17</sup> have been developed. Moreover, reactions conducted in ionic liquids<sup>18</sup> or using the electrochemical method<sup>19</sup> have also been reported. However, those procedures pose a number of drawbacks, in particular, use of complicated catalysts,



Scheme 1 Chemical fixation of  $\text{CO}_2$  into  $\alpha$ -alkylidene cyclic carbonates.



View Article Online

E-mail: weili@nankai.edu.cn

 $<sup>\</sup>dagger\, Electronic$  supplementary information (ESI) available. See DOI: 10.1039/ c3gc42406e

<sup>‡</sup>Authors with equal contributions.



 $\mbox{Scheme 2}$  Chemical fixation of  $\mbox{CO}_2$  through the dual activation pathway.

requirement of higher  $CO_2$  pressure and avoidance of moisture. These challenges can be surmounted, to our delight, through the rational design of a high performance catalytic system. Consequently, a robust catalytic system is believed to be essential to find novel protocols for chemical fixation of  $CO_2$  to value-added products in a cost-effective, practically viable, and environmentally benign manner.

Silver salts are widely employed as activators of the C=C triple bond in organic synthesis.<sup>20</sup> For the reaction of propargyl alcohols with CO<sub>2</sub>, the Yamada group<sup>10</sup> and the Jiang group<sup>16</sup> have developed a silver-based procedure for the synthesis of the  $\alpha$ -alkylidene cyclic carbonates from CO<sub>2</sub>. However, a stoichiometric amount of the base or high CO<sub>2</sub> pressure was required in those Ag-catalyzed systems. On the other hand, Mizuno et al.<sup>21</sup> reported that  $[WO_4]^{2-}$  could activate a  $CO_2$  molecule by forming the  $[WO_4]^{2-}/CO_2$  adduct, which thus could allow the CO2 transformation to proceed smoothly under mild conditions. In this context, we envisaged that a novel bifunctional catalytic species containing Ag<sup>+</sup> and  $[WO_4]^{2-}$ , *i.e.* Ag<sub>2</sub>WO<sub>4</sub>, could be particularly efficient for the reaction of propargylic alcohols with  $CO_2$ . The  $[WO_4]^{2-}$  anion would activate both CO<sub>2</sub> and the O-H bond of propargylic alcohol,<sup>21a</sup> which could favor the generation of the carboxylate intermediate. The carboxylate intermediate would then be trapped by the silver-activated C=C triple bond to afford the stable α-alkylidene cyclic carbonate as shown in Scheme 2.

Herein, we would like to report  $Ag_2WO_4/Ph_3P$  as a highly efficient and robust bifunctional catalyst system for chemical fixation of atmospheric CO<sub>2</sub> with propargyl alcohols to provide  $\alpha$ -alkylidene cyclic carbonates under neutral solvent-free conditions. Notably, this catalytic system was also proved to be effective for the three-component coupling reaction of propargylic alcohols, amines and CO<sub>2</sub>. The present system can eliminate the need for large quantities of caustic base or flammable organic solvents.

#### **Results and discussion**

Several metal compounds were initially investigated for the reaction of 2-methylbut-3-yn-2-ol (1a), as the model substrate under a CO<sub>2</sub> balloon without a solvent at 50 °C for 6 h (Table 1). The reaction without a catalyst and an additive was carried out only to recover the starting material quantitatively (entry 1, Table 1). The catalysts such as DBU, Ph<sub>3</sub>P, and CuCl were demonstrated to be ineffective under the given conditions (entries 2–4). Different silver salts like AgNO<sub>3</sub>, AgOAc, Ag<sub>2</sub>WO<sub>4</sub>,

Table 1 Ag(I)-catalyzed reaction of propargyl alcohol 1a with CO<sub>2</sub><sup>a</sup>



Entry	Catalyst	Additive	Time/h	Yield <sup>b</sup> /%
1	_	_	6	0
2	DBU	_	6	0
3	Ph <sub>2</sub> P	_	6	0
4	CuCl	_	6	0
5	AgNO <sub>3</sub>	Ph <sub>3</sub> P	6	0
6	AgOAc	Ph <sub>3</sub> P	6	<1
7	$Ag_2WO_4$	Ph <sub>3</sub> P	6	45
8	AgI	Ph <sub>3</sub> P	6	0
9	$Ag_2WO_4$	DBU	6	2
10	$Ag_2WO_4$	2,2'-Bipyridine	6	0
11	$Ag_2WO_4$	TMEDA	6	0
$12^{c}$	$Ag_2WO_4$	Ph <sub>3</sub> P	3	85
13 <sup>c</sup>	$Ag_2WO_4$	L1	3	0
$14^c$	$Ag_2WO_4$	L2	3	<1
$15^{c}$	$Ag_2WO_4$	L3	3	82
16	$Ag_2WO_4$	Ph <sub>3</sub> P	12	>99
$17^d$	$Ag_2WO_4$	Ph <sub>3</sub> P	12	98
$18^e$	$Ag_2WO_4$	Ph <sub>3</sub> P	12	82
$19^{f}$	$Ag_2WO_4$	Ph <sub>3</sub> P	6	>99
$20^g$	$Ag_2WO_4$	Ph <sub>3</sub> P	6	>99
	-			

<sup>*a*</sup> Unless otherwise specified, the reactions were performed with 2-methylbut-3-yn-2-ol (0.421 g, 5.0 mmol),  $[Ag^+]$  or other catalysts (2 mol%), additive (2 mol%), CO<sub>2</sub> balloon, 50 °C. DBU = 1,8-diazabicyclo[5.4.0]undec-7-ene; TMEDA = tetramethylethylene-diamine. <sup>*b*</sup> GC yields were given using biphenyl as the internal standard. <sup>*c*</sup> Additive (1 mol%), 0.3 MPa. <sup>*d*</sup> 25 °C. <sup>*c*</sup> Ag<sub>2</sub>WO<sub>4</sub> (16.3 mg, 0.7 mol%), Ph<sub>3</sub>P (18.4 mg, 1.4 mol%). <sup>*f*</sup> 0.5 MPa. <sup>*g*</sup> Ag<sub>2</sub>WO<sub>4</sub> (58.0 mg, 2.5 mol%), Ph<sub>3</sub>P (65.5 mg, 5 mol%).

and AgI were also screened in the presence of Ph<sub>3</sub>P as an additive (entries 5–8). Gratifyingly, the  $\alpha$ -alkylidene cyclic carbonates 2a were obtained in a yield of 45% by employing 1 mol% of Ag<sub>2</sub>WO<sub>4</sub> in conjunction with 2 mol% of Ph<sub>3</sub>P at 50 °C for 6 h (entry 7). The results encouraged us to further optimize the reaction conditions by using Ag<sub>2</sub>WO<sub>4</sub> as a catalyst. The reaction did not occur using DBU, 2,2'-bipyridine, or TMEDA as an additive (entries 9-11). Then several bidentate phosphine ligands were screened under 0.3 MPa of  $CO_2$  (entries 13–15). Those results show that L1 and L2 were not effective, whereas L3 gave 82% yield of the target product. Gratifyingly, a simple Ag<sub>2</sub>WO<sub>4</sub>/Ph<sub>3</sub>P catalytic system gave 85% yield of 2a by increasing CO<sub>2</sub> to 0.3 MPa (entry 12). Based on this result, a quantitative yield was successfully reached in the presence of 0.1 MPa CO<sub>2</sub> at 50 °C or 25 °C by further prolonging the reaction time to 12 h (entries 16 and 17). With lower loading of the catalyst, the reaction at 50 °C for 12 h afforded 2a in a yield of 82% (entry 18). Furthermore, it was found that higher  $CO_2$  pressure or catalyst loading was needed for the full conversion in 6 h (entries 19 and 20).

Table 2 Carboxylative cyclization of propargyl alcohols with CO<sub>2</sub><sup>a</sup>



<sup>*a*</sup> Unless otherwise specified, the reactions were performed with 1 (5.0 mmol),  $Ag_2WO_4$  (23.2 mg, 1 mol%),  $Ph_3P$  (26.2 mg, 2 mol%), 25 °C. <sup>*b*</sup> Isolated yield. <sup>*c*</sup> 50 °C. <sup>*d*</sup> 1 MPa of CO<sub>2</sub> and 80 °C were used.

Subsequently, the substrate scope of this reaction was investigated under 0.1 MPa of CO<sub>2</sub> at 25 °C (Table 2). Propargyl alcohols with a variety of alkyl and aryl substituents at the propargylic position (**1a–1e**) were effective substrates to give the corresponding carbonates **2a–2e** in good to excellent yields even under the conditions of atmospheric CO<sub>2</sub> (entries 1–5, Table 2), whereas 1-isopropyl propargyl alcohol lost reactivity



Fig. 1 Catalyst recycling for the carboxylation of 1a with CO $_2$  (0.5 MPa) at 50 °C for 6 h.

(entry 6), and 1,1-cyclopentylene showed low activity (entry 7), probably due to the steric hindrance and ring strain. The internal propargyl alcohol **1h** required higher  $CO_2$  pressure (1 MPa) and temperature (80 °C) to afford the corresponding carbonate in 82% yield after 36 h (entry 8). However, the secondary alcohol **1i** failed in giving the cyclic carbonate but the subsequent alcoholysis product **2i** was formed (entry 9).

Notably, the reaction of propargyl alcohol 1a with CO<sub>2</sub> (0.5 MPa) at 50 °C was continuously run to investigate the constancy of the catalyst activity. In each cycle, the heterogeneous catalyst could be easily separated by extraction with hexane. After drying with nitrogen, the catalyst was reused for the next run. As shown in Fig. 1, the catalyst can be reused at least 4 times without significant loss of catalytic activity.

The successful transformation of propargyl alcohols to  $\alpha$ -alkylidene cyclic carbonates through CO<sub>2</sub> fixation under extremely mild conditions prompted us to expand potential applications of this Ag<sub>2</sub>WO<sub>4</sub>/Ph<sub>3</sub>P system to the synthesis of  $\alpha$ -alkylidene cyclic carbamates *via* the three-component reaction<sup>15,22</sup> of propargylic alcohols, primary amines and CO<sub>2</sub> under mild conditions as listed in Table 3. Pleasingly, the transformation was performed smoothly with the catalysis of Ag<sub>2</sub>WO<sub>4</sub> in the presence of 0.5 MPa CO<sub>2</sub> at 50 °C for 12 h. When 1,1-dimethylpropargylic alcohol was used, and amines bearing n-propyl, n-butyl, cyclohexyl, and benzyl groups respectively were subjected to the reaction conditions, the 5-methyleneoxazo-lidin-2-one derivatives (4a-4d)were obtained in good to excellent yields. In the case of aniline, the undehydrated product 4e was afforded in 31% yield. Interestingly, another five-membered ring was constructed via intramolecular dehydration to furnish 4f in 91% yield by employing ethanolamine as a substrate. Reactions of propargylic alcohols with a range of substituents including 1-methyl, 1-ethyl, 1-phenyl, and 1-n-pentyl groups gave the desired products (4g-4i). Moreover, the steric hindrance of the tertiary butyl group of the amine significantly diminishes the reactivity of the carbamate species, which failed to provide the intramolecular cycloaddition product but β-oxopropylcarbamate 4k was formed. Disappointingly, the corresponding carbamate 4l was not obtained with internal propargyl alcohol as a starting material.





<sup>*a*</sup> Unless otherwise stated, the reactions were conducted on a 5 mmol scale of propargylic alcohol and amine with  $Ag_2WO_4$  (23.2 mg, 1 mol%) and  $Ph_3P$  (26.2 mg, 2 mol%) under 0.5 MPa of  $CO_2$  at 50 °C for 12 h. Isolated yield was given.



Scheme 3 The  $Ag_2WO_4/Ph_3P$ -catalyzed three-component reaction of propargylic alcohols, secondary amines and  $CO_2$ . Isolated yield was given.

Similarly, secondary amines were also applicable to the three-component reaction under identical reaction conditions, which afforded  $\beta$ -oxopropylcarbamates as products.<sup>11,22*a*,23</sup> Various representative  $\beta$ -oxopropylcarbamates synthesized are depicted in Scheme 3. In most cases, the reaction was performed with an approximately equimolar amount of reactants and proceeded to completion within 12 h at 50 °C. Aliphatic propargylic alcohols, including 1,1-dimethyl propargylic



Scheme 4 Stepwise synthesis of typical oxazolidinone and  $\beta\text{-}oxopropylcarbamate.}$ 



Scheme 5 Proposed mechanism for the  ${\rm Ag_2WO_4/Ph_3P}\xspace$  three-component reactions.

alcohol, 1-methyl-1-phenyl propargylic alcohol, 1-ethyl-1methyl propargylic alcohol, and 1-ethynylcyclohexanol, were also found to be good substrates. Both cyclic and linear secondary aliphatic amines afforded the corresponding product with good to excellent yields. Unfortunately, the reaction using the aromatic amines cannot furnish the desired products.

Although we are quite confident that the oxazolidinone generated from the three-component coupling reaction between propargylic alcohols, primary amines and CO<sub>2</sub> proceeds *via* a tandem aminolysis of  $\alpha$ -alkylidene cyclic carbonates/intramolecular cyclization,<sup>22c,23b</sup> the individual steps of this process were further examined as depicted in Scheme 4. In the absence of the amine, the expected carbonates **2a** were produced in 98% GC yield under the reaction conditions. Subsequently, equimolar primary and secondary amines were added into the reaction mixture, respectively. The corresponding products **4b** and **6a** were isolated in reasonable yields after 0.5 h.

Consequently, a plausible mechanism of the  $Ag_2WO_4/Ph_3P$ catalyzed three-component reactions is proposed as illustrated in Scheme 5. The carboxylative cyclization of the propargyl alcohol with  $CO_2$  affords the carbonate 2 as an intermediate, which then undergoes a nucleophilic attack by a primary or secondary amine to generate the carbamate species. In the presence of a secondary amine, the corresponding

Paper

 $\beta$ -oxopropylcarbamate **6** can be obtained *via* tautomerization. In the case of a primary amine, further intramolecular nucleophilic attack followed by dehydration affords the oxazolidinone **4**.

### Conclusions

In summary, we have described a sustainable and cost-competitive catalytic process promoted by the Ag<sub>2</sub>WO<sub>4</sub>/Ph<sub>3</sub>P system for efficient chemical fixation of  $CO_2$  to produce  $\alpha$ -alkylidene cyclic carbonates,  $\alpha$ -alkylidene cyclic carbamates, and  $\beta$ -oxopropylcarbamates under mild conditions, respectively. The scope of the reaction is broad, with a variety of propargylic alcohols undergoing carboxylative cyclization. Moreover, the three-component reactions of propargylic alcohols, amines and CO<sub>2</sub> are also well performed with this protocol. In addition, this procedure can eliminate the need for large quantities of caustic base and flammable organic solvents, yielding the strategy to prepare value-added  $\alpha$ -alkylidene cyclic carbonates,  $\alpha$ -alkylidene cyclic carbamates, and  $\beta$ -oxopropylcarbamates from both environmental and economical points of view. Importantly, the catalytic system is not sensitive to air and minor moisture, which may enhance the possibility of fixation of CO<sub>2</sub>. Such findings may be of great interest for developing alternative access to chemical fixation of CO2 with the preparation of the CO<sub>2</sub>-based products under mild conditions.

#### **Experimental section**

General procedure for the synthesis of α-alkylidene cyclic carbonates 2a: Ag<sub>2</sub>WO<sub>4</sub> (23.2 mg, 1 mol%), PPh<sub>3</sub> (26.2 mg, 2 mol%), and propargylic alcohols (5 mmol) were added to a Schlenk tube equipped with a magnetic stir bar. A CO<sub>2</sub> balloon was consecutively introduced. Then the mixture was stirred at 25 °C for the desired time. Upon completion, the mixture was diluted with *n*-hexane  $(3 \times 5 \text{ mL})$ . The organic phase was collected and then purified by column chromatography on silica gel using petroleum ether-ethyl acetate as an eluent to give the desired products. The solid residue in the tube was Ag<sub>2</sub>WO<sub>4</sub>, which could be used directly in the next run. 2a was obtained as a colourless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz)  $\delta$  4.75 (d, J = 4.0 Hz, 1H), 4.31 (d, J = 4.0 Hz, 1H), 1.59 (s, 6H) ppm. <sup>13</sup>C NMR  $(CDCl_3, 100.6 \text{ MHz}) \delta$  158.6 (C=O), 151.2, 85.2, 84.6, 27.5 ppm. MS (EI, 70 eV) m/z (%) = 128.10 (2.81), 85.10 (6.49), 84.10 (100), 83.10 (3.54), 69.10 (48.16).

## Acknowledgements

We are grateful to the National Natural Science Foundation of China (no. 21172125 and 21376123), Specialized Research Fund for the Doctoral Program of Higher Education (20130031110013), and Tianjin Co-Innovation Center of Chemical Science and Engineering.

# Notes and references

- (a) P. H. Abelson, Science, 2000, 289, 1293; (b) D. W. Keith, Science, 2009, 325, 1654–1655; (c) C. Federsel, R. Jackstell and M. Beller, Angew. Chem., Int. Ed., 2010, 49, 6254–6257; (d) A.-H. Liu, R. Ma, C. Song, Z.-Z. Yang, A. Yu, Y. Cai, L.-N. He, Y.-N. Zhao, B. Yu and Q.-W. Song, Angew. Chem., Int. Ed., 2012, 51, 11306–11310.
- 2 (a) D. J. Darensbourg, Chem. Rev., 2007, 107, 2388-2410;
  (b) J. Louie, Curr. Org. Chem., 2005, 9, 605-623;
  (c) S. N. Riduan and Y. Zhang, Dalton Trans., 2010, 39, 3347-3357; (d) X. Yin and J. R. Moss, Coord. Chem. Rev., 1999, 181, 27-60; (e) Z.-Z. Yang, Y.-N. Zhao and L.-N. He, RSC Adv., 2011, 1, 545-567; (f) L.-N. He, J.-Q. Wang and J.-L. Wang, Pure Appl. Chem., 2009, 81, 2069-2080.
- 3 For reviews on chemical fixation of CO<sub>2</sub>, see:
  (a) T. Sakakura, J.-C. Choi and H. Yasuda, *Chem. Rev.*, 2007, 107, 2365–2387; (b) M. Cokoja, C. Bruckmeier, B. Rieger, W. A. Herrmann and F. E. Kühn, *Angew. Chem., Int. Ed.*, 2011, 50, 8510–8537; (c) K. Huang, C.-L. Sun and Z.-J. Shi, *Chem. Soc. Rev.*, 2011, 40, 2435–2452; (d) Y. Tsuji and T. Fujihara, *Chem. Commun.*, 2012, 48, 9956–9964; (e) W.-Z. Zhang and X.-B. Lu, *Chin. J. Catal.*, 2012, 33, 745–756; (f) Z.-Z. Yang, L.-N. He, J. Gao, A.-H. Liu and B. Yu, *Energy Environ. Sci.*, 2012, 5, 6602–6639.
- 4 (a) G. Centi, E. A. Quadrelli and S. Perathoner, *Energy Environ. Sci.*, 2013, 6, 1711–1731; (b) P. Markewitz, W. Kuckshinrichs, W. Leitner, J. Linssen, P. Zapp, R. Bongartz, A. Schreiber and T. E. Muller, *Energy Environ. Sci.*, 2012, 5, 7281–7305.
- 5 A few reports on the transformation of  $\alpha$ -alkylidene cyclic carbonates, see: (a) Y. Inoue, K. Matsushita, I. F. Yen and S. Imaizumi, *Chem. Lett.*, 1991, **20**, 1377–1378; (b) K. Ohe, H. Matsuda, T. Ishihara, S. Ogoshi, N. Chatani and S. Murai, *J. Org. Chem.*, 1993, **58**, 1173–1177; (c) K. Ohe, H. Matsuda, T. Morimoto, S. Ogoshi, N. Chatani and S. Murai, *J. Am. Chem. Soc.*, 1994, **116**, 4125–4126; (d) M. Yoshida and M. Ihara, *Angew. Chem., Int. Ed.*, 2001, **40**, 616–619.
- 6 C. Bruneau and P. H. Dixneuf, J. Mol. Catal., 1992, 74, 97–107.
- 7 K. Iritani, N. Yanagihara and K. Utimoto, *J. Org. Chem.*, 1986, **51**, 5499–5501.
- 8 Y. Inoue, J. Ishikawa, M. Taniguchi and H. Hashimoto, *Bull. Chem. Soc. Jpn.*, 1987, **60**, 1204–1206.
- 9 (a) H.-S. Kim, J.-W. Kim, S.-C. Kwon, S.-C. Shim and T.-J. Kim, J. Organomet. Chem., 1997, 545-546, 337-344;
  (b) H.-F. Jiang, A.-Z. Wang, H.-L. Liu and C.-R. Qi, Eur. J. Org. Chem., 2008, 2309-2312.
- 10 (a) S. Kikuchi, S. Yoshida, Y. Sugawara, W. Yamada, K. Sekine, I. Iwakura, T. Ikeno, T. Yamada and H.-M. Cheng, *Bull. Chem. Soc. Jpn.*, 2011, 84, 698–717; (b) W. Yamada, Y. Sugawara, H. M. Cheng, T. Ikeno and T. Yamada, *Eur. J. Org. Chem.*, 2007, 2604–2607.
- 11 T.-J. Kim, K.-H. Kwon, S.-C. Kwon, J.-O. Baeg, S.-C. Shim and D.-H. Lee, *J. Organomet. Chem.*, 1990, **389**, 205–217.

- 12 (a) J. M. Joumier, J. Fournier, C. Bruneau and P. H. Dixneuf, J. Chem. Soc., Perkin Trans. 1, 1991, 3271– 3274; (b) Y. Kayaki, M. Yamamoto and T. Ikariya, J. Org. Chem., 2007, 72, 647–649.
- 13 Y. Kayaki, M. Yamamoto and T. Ikariya, *Angew. Chem., Int. Ed.*, 2009, **48**, 4194–4197.
- 14 K. Uemura, T. Kawaguchi, H. Takayama, A. Nakamura and Y. Inoue, *J. Mol. Catal. A: Chem.*, 1999, **139**, 1–9.
- 15 N. D. Ca, B. Gabriele, G. Ruffolo, L. Veltri, T. Zanetta and M. Costa, *Adv. Synth. Catal.*, 2011, **353**, 133–146.
- 16 X. Tang, C. Qi, H. He, H. Jiang, Y. Ren and G. Yuan, *Adv. Synth. Catal.*, 2013, **355**, 2019–2028.
- 17 Y.-B. Wang, Y.-M. Wang, W.-Z. Zhang and X.-B. Lu, J. Am. Chem. Soc., 2013, 135, 11996–12003.
- 18 Y. L. Gu, F. Shi and Y. Q. Deng, J. Org. Chem., 2004, 69, 391-394.
- 19 G.-Q. Yuan, G.-J. Zhu, X.-Y. Chang, C.-R. Qi and H.-F. Jiang, *Tetrahedron*, 2010, **66**, 9981–9985.
- 20 For examples of silver-promoted activation of alkynes, see:
  (*a*) U. Halbes-Letinois, J.-M. Weibel and P. Pale, *Chem. Soc. Rev.*, 2007, 36, 759–769; (*b*) X.-Q. Yao and C.-J. Li, *Org. Lett.*, 2005, 7, 4395–4398; (*c*) J. Liu, Z. Fang, Q. Zhang, Q. Liu and X. Bi, *Angew. Chem., Int. Ed.*, 2013, 52, 6953–6957;

(d) M. Gao, C. He, H. Chen, R. Bai, B. Cheng and A. Lei, Angew. Chem., Int. Ed., 2013, 52, 6958–6961; (e) F. Li, J. Nie, L. Sun, Y. Zheng and J.-A. Ma, Angew. Chem., Int. Ed., 2013, 52, 6255–6258; (f) C. He, S. Guo, J. Ke, J. Hao, H. Xu, H. Chen and A. Lei, J. Am. Chem. Soc., 2012, 134, 5766–5769; (g) X. Zhang, W. Z. Zhang, X. Ren, L. L. Zhang and X. B. Lu, Org. Lett., 2011, 13, 2402–2405.

- 21 (a) T. Kimura, K. Kamata and N. Mizuno, Angew. Chem., Int. Ed., 2012, 51, 6700–6703; (b) T. Kimura, H. Sunaba, K. Kamata and N. Mizuno, Inorg. Chem., 2012, 51, 13001–13008.
- 22 For selected examples of the synthesis of N-substituted 4-methylene-2-oxazolidinones, see: (a) Y. L. Gu, Q. H. Zhang, Z. Y. Duan, J. Zhang, S. G. Zhang and Y. Q. Deng, *J. Org. Chem.*, 2005, **70**, 7376–7380; (b) H. Jiang, J. Zhao and A. Wang, *Synthesis*, 2008, 763–769; (c) Y. Sasaki and P. H. Dixneuf, *J. Org. Chem.*, 1987, **52**, 4389–4391.
- 23 Chemical fixation of CO<sub>2</sub> into β-oxopropylcarbamates, see:
  (a) Q. Zhang, F. Shi, Y. Gu, J. Yang and Y. Deng, *Tetrahedron Lett.*, 2005, 46, 5907–5911;
  (b) C.-R. Qi and H.-F. Jiang, *Green Chem.*, 2007, 9, 1284–1286;
  (c) H.-S. Kim, J.-W. Kim, S.-C. Kwon, S.-C. Shim and T.-J. Kim, *J. Organomet. Chem.*, 1997, 545–546, 337–344.